

Increased periostin associates with greater airflow limitation in patients receiving inhaled corticosteroids

Yoshihiro Kanemitsu MD^{*1,2}, Hisako Matsumoto MD, PhD^{*1,2}, Kenji Izuhara MD, PhD³, Yuji Tohda MD, PhD^{2,4}, Hideo Kita MD, PhD^{2,5}, Takahiko Horiguchi MD, PhD^{2,6}, Kazunobu Kuwabara MD, PhD^{2,6}, Keisuke Tomii MD, PhD^{2,7}, Kojiro Otsuka MD, PhD^{1,2,7}, Masaki Fujimura MD, PhD^{2,8}, Noriyuki Ohkura MD^{2,8}, Katsuyuki Tomita MD, PhD^{2,4}, Akihito Yokoyama MD, PhD^{2,9}, Hiroshi Ohnishi MD, PhD^{2,9}, Yasutaka Nakano MD, PhD^{2,10}, Tetsuya Oguma MD, PhD^{2,10}, Soichiro Hozawa MD, PhD^{2,11}, Tadao Nagasaki MD¹, Isao Ito MD, PhD¹, Tsuyoshi Oguma MD¹, Hideki Inoue MD¹, Tomoko Tajiri MD¹, Toshiyuki Iwata MD¹, Yumi Izuhara MD¹, Junya Ono BS¹², Shoichiro Ohta MD, PhD¹³, Mayumi Tamari MD, PhD¹⁴, Tomomitsu Hirota DDS, PhD¹⁴, Tetsuji Yokoyama MD, PhD¹⁵, Akio Niimi MD, PhD^{1,2,16} and Michiaki Mishima MD, PhD^{1,2}

¹Department of Respiratory Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan

²Kinki Hokuriku Airway disease Conference (KiHAC)

³Division of Medical Biochemistry, Department of Biomolecular Sciences, Saga Medical School, Saga, Japan

⁴Department of Respiratory Medicine and Allergology, Faculty of Medicine, Kinki University, Osaka, Japan

⁵Department of Respiratory Medicine, Takatsuki Red Cross Hospital, Osaka, Japan

⁶Department of Respiratory Internal Medicine, Fujita Health University Second Educational Hospital, Aichi, Japan

⁷Department of Respiratory Medicine, Kobe City Medical Center General Hospital, Hyogo, Japan

⁸Department of Respiratory Medicine, Cellular Transplantation Biology, Kanazawa

University Graduate School of Medicine, Kanazawa, Japan

⁹Department of Hematology and Respiratory Medicine, Kochi University, Kochi, Japan

¹⁰Division of Respiratory Medicine, Department of Internal Medicine, Shiga University of Medical Science, Shiga, Japan

¹¹Hiroshima Allergy and Respiratory Clinic, Hiroshima, Japan

¹²Shino-Test Corporation, Kanagawa, Japan

¹³Department of Laboratory Medicine, Saga Medical School, Saga, Japan

¹⁴Laboratory for Respiratory Diseases, Center for Genomic Medicine, RIKEN, Yokohama, Kanagawa, Japan

¹⁵Department of Health Promotion, National Institute of Public Health, Wako, Saitama, Japan

¹⁶Department of Medical Oncology and Immunology, Nagoya City University School of Medical Sciences, Aichi, Japan

*YK and HM contributed equally to this study.

Correspondence should be addressed to Hisako Matsumoto MD, PhD

Department of Respiratory Medicine, Graduate School of Medicine, Kyoto University

53 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

E-mail: hmatsumo@kuhp.kyoto-u.ac.jp

Funded by KiHAC, as a project of 2009 KiHAC Respiratory Medicine Group and the

Adaptable and Seamless Technology Transfer Program through target-driven R&D, JST.

Abstract

Background: Periostin, an extracellular matrix protein, contributes to subepithelial thickening in asthmatic airways, and its serum levels reflect airway eosinophilic inflammation. However, the relationship between periostin and the development of airflow limitation, a functional consequence of airway remodeling, remains unknown.

Objective: To determine the relationship between serum periostin levels and pulmonary function decline in asthmatic patients on inhaled corticosteroid (ICS) treatment.

Methods: 224 asthmatic patients (average age 62.3 years) treated with ICS for at least 4 years were enrolled. Annual changes in forced expiratory volume in one second (FEV₁), from at least one year after the initiation of ICS treatment to the time of enrollment or later (average 16.2 measurements over 8 years per individual), were assessed. At enrollment, clinical indices, biomarkers including serum periostin, and periostin gene polymorphisms were examined. Associations between clinical indices or biomarkers and a decline in FEV₁ of 30 mL·yr⁻¹ or greater were analyzed.

Results: High serum periostin levels (≥ 95 ng/mL) at enrollment, the highest treatment step, higher ICS daily doses, a history of admission due to asthma exacerbation, comorbid or a history of sinusitis, and ex-smoking were associated with a decline in FEV₁ of 30 mL·yr⁻¹ or greater. Multivariate analysis revealed that high serum periostin, the highest treatment step, and ex-smoking were independent risk factors for the decline. Polymorphisms of periostin gene were related to higher serum periostin levels (rs3829365) and a decline in FEV₁ of 30 mL·yr⁻¹ or greater (rs9603226).

Conclusions: Serum periostin appears to be a useful biomarker for the development of airflow limitation in asthmatic patients on ICS.

Clinical implications (25 words)

Serum periostin levels reflect greater FEV₁ decline in asthmatic patients on inhaled

74 corticosteroid treatment. *POSTN* gene polymorphisms may also be helpful for identifying
75 rapid FEV₁ decliners.

76 **Key words**

77 Asthma, inhaled corticosteroids, lung function decline, periostin, *POSTN* gene polymorphism,
78 sinusitis, treatment step

79

80 **Abbreviations**

81 ACT: asthma control test

82 ECP: eosinophil cationic protein

83 FAS I: fasciclin I

84 FEV₁: forced expiratory volume in one second

85 FVC: forced vital capacity

86 hsCRP: high sensitivity C-reactive protein

87 ICS: inhaled corticosteroids

88 IgE: immunoglobulin E

89 IL: interleukin

90 ROC: receiver operating characteristic

91 SNP: single-nucleotide polymorphism

92 TGF- β : transforming growth factor beta

93

94 Total word counts for the text and the abstract are 3800 and 258 words, respectively.

95 **Capsule summary (32 words)**

96 This is the first study to identify a relationship between high serum periostin and greater
97 annual decline in FEV₁, which sheds new light on serum periostin as a useful biomarker in
98 asthma.

99 Introduction

100 Airway inflammation and remodeling are key features of asthma that have been
 101 demonstrated by pathological¹ and radiological findings^{2,3}. Physiologically, patients with
 102 asthma show a greater decline in pulmonary function than subjects without asthma⁴. Studies
 103 that were mostly conducted in the era before inhaled corticosteroids (ICS) demonstrated that
 104 more severe symptoms or severe exacerbations⁵⁻⁷, long-standing asthma⁸, and smoking
 105 history^{4,8} were moderate to strong risk factors for greater decline in pulmonary function⁵.
 106 Blood and sputum eosinophilia^{9,10} and genetic predisposition¹¹⁻¹³ were also potential risk
 107 factors. Owing to early intervention with ICS, however, airway inflammation and the degree
 108 of annual decline in pulmonary function have been attenuated in a majority of asthmatic
 109 patients¹⁴⁻¹⁶. Meanwhile, a subset of patients still show accelerated decline in FEV₁ and
 110 develop irreversible airway obstruction despite adequate treatment^{17,18}. van Veen et al. found
 111 that exhaled nitric oxide of 20 ppb or higher is a predictor of accelerated decline in
 112 pulmonary function in patients with difficult-to-treat asthma¹⁸. However, other biomarkers for
 113 greater decline in FEV₁ despite treatment with ICS remain unknown.

114 The airway inflammation of asthma is classically characterized by infiltration and
 115 activation of eosinophils, mast cells, and Th2 cells with several mediators and Th2 cytokines,
 116 such as interleukin (IL)-4, IL-5, and IL-13^{19,20}. Periostin, a secreted, 90-kDa, extracellular
 117 matrix protein that is induced by IL-4 and IL-13, was originally isolated as an osteoblast-
 118 specific factor; it shares structural homology to the insect cell adhesion molecule fasciclin I
 119 (FAS I) and binds to fibronectin, tenascin-C, and collagen^{21,22}. In airway epithelial cells
 120 collected from patients with asthma, periostin is one of the up-regulated genes²³, and its
 121 expression is correlated with thickness of the airway basement membrane²⁴. Takayama et al.
 122 clearly demonstrated that periostin is deposited in the airway subepithelial layer in asthmatic
 123 patients. Moreover, serum periostin is identified as the single best predictor of airway
 124 eosinophilia in patients with severe asthma who remain symptomatic despite maximal ICS

125 treatment²⁵. Therefore, we hypothesized that periostin would be a novel biomarker of
126 Th2/eosinophil-driven airway inflammation and greater decline in pulmonary function, a
127 functional consequence of airway remodeling in patients with asthma.

128 In this study, the effects of biomarkers and clinical indices on greater annual decline
129 in pulmonary function in asthmatic patients on ICS treatment were examined, with the
130 specific aim of determining the association between serum periostin levels and pulmonary
131 function decline. Polymorphisms of the *POSTN* gene, which encodes periostin, were also
132 examined on the hypothesis that *POSTN* gene polymorphisms may affect serum periostin
133 levels.

Methods

For full details see Online Repository

Patients

Patients with asthma were recruited from nine institutions belonging to the Kinki Hokuriku Airway disease Conference where asthma specialists manage patients. Asthma was diagnosed according to the American Thoracic Society criteria²⁶. From September 2009 to December 2011, patients were enrolled if they had received ICS treatment for 4 years or more, undergone three or more pulmonary function tests when they were stable, and were free from exacerbations for at least one month. The first pulmonary function test was performed at least one year after the commencement of ICS treatment and at 25 years of age or older. Patients who had smoked more than 10 pack-years, smoked in the past one year, or had other pulmonary diseases were excluded.

This study was approved by the ethics committee of each participant institution and was registered in the UMIN Clinical Trials Registry (Registry ID UMIN000002414). Written informed consent was obtained from all participants.

Measurements

At enrollment, patients underwent a work-up that included answering a self-completed questionnaire, spirometry, and blood tests. After enrollment, spirometry was repeated at least 6 months later for up to 12 months.

Self-completed questionnaire and clinical indices

The self-completed questionnaire was composed of 4 major items, as presented in Table 1. The Asthma Control Test (ACT)TM was also scored. The treatment step at enrollment was determined according to the Global Initiative for Asthma 2010 guideline²⁷.

Pulmonary function

Spirometry was performed using an electrical spirometer, which was calibrated once a week, at each institution. Spirometry data were obtained only when patients were stable. To determine pulmonary function on daily medications, ICS and other controllers, including long-acting β_2 agonists, leukotriene receptor antagonists, or slow-release theophylline, were not withdrawn before spirometry.

Measurement of systemic biomarkers

Blood eosinophil and neutrophil counts, and serum levels of total immunoglobulin E (IgE), specific IgE against common inhaled allergens, eosinophil cationic protein (ECP), high sensitivity C-reactive protein (hsCRP), and periostin were determined.

Serum periostin levels were measured using an enzyme-linked immunosorbent assay at Shino-test (Kanagawa, Japan), as described previously²⁸. Pooled serum periostin level data from 66 healthy subjects [mean (SD), 60.7 (16.7) years old, 40 males]^{28,29} were used for comparison with those of asthmatic patients.

Haplotype analysis, DNA extraction, and genotyping of the *POSTN* gene

A total of 47 single-nucleotide polymorphisms (SNPs) in the region of the *POSTN* gene and its upstream, total 39 kb, was captured in the HapMap Japanese data set. Haplotype analysis identified 4 major haplotypes and 2 minor haplotypes. Two minor haplotypes were grouped into the closest major haplotype, and 3 tag SNPs that determined the 4 haplotypes were identified (Figure 1).

Genomic DNA was isolated from blood cells using a QIAamp DNA Blood Mini Kit (Qiagen, Tokyo, Japan). SNPs were genotyped using a Taqman genotyping assay according to the manufacturer's instructions (Applied Biosystems, Tokyo, Japan) and analyzed using an Applied Biosystems 7300 Real-Time PCR System (Applied Biosystems).

Statistical analysis

Statistical analyses were performed using JMP version 9.0 (SAS Institute Inc., Tokyo, Japan). Annual changes in FEV₁ (Δ FEV₁) were estimated for each subject by fitting a least-square regression line to all of his/her all available data points. Receiver operating characteristic (ROC) curve analysis was performed to determine a serum periostin cut-off value for asthmatic patients. The effects of serum biomarkers or other indices on Δ FEV₁ were estimated using a generalized linear mixed model with adjustment for sex, height, age at enrollment, and FEV₁ at the first measurement. The institutions were included as random effects in this model. On univariate analysis of Δ FEV₁, the adjusted p value, i.e., q value, which was a measure of significance in terms of the false discovery rate, was obtained using R and QVALUE software³⁰ to determine spurious significance in multiple testing. The effects on the dichotomous data for a decline in FEV₁ of -30 mL·yr⁻¹ or greater³¹ were similarly estimated using a generalized linear mixed model by IBM SPSS Advanced Statistics 19 (SPSS Inc., Tokyo, Japan). Multivariate analysis was performed using variables with $p < 0.10$ on univariate analysis, except for ICS daily maintenance dose because of its strong correlation with treatment step. On multivariate analysis, the periostin level was considered as a dichotomous variable (high or low) instead of a continuous variable. Correlation coefficients between serum periostin levels and clinical indices were estimated by fitting least-square regression lines to data, in which institutions were included as random effects. Unpaired *t*- and Chi-square tests were performed for comparisons of continuous and dichotomous variables, respectively. When data were not normally distributed, they were log-transformed. Data are presented as means (SD). P values ≤ 0.05 were considered significant.

Results

Patients' characteristics

Initially, 233 patients were enrolled in this study, but 9 patients were excluded: 5 with a smoking history of more than 10 pack-years and 4 who did not have enough pulmonary function data available. The demographic data of the remaining 224 patients are presented in Table 2. The mean age at enrollment was 62.3 (13.7) years. Overall, 130 (58%) had onset of asthma at 40 years or older. The average number of measurements of FEV₁, follow-up period, and Δ FEV₁ of 224 patients were 16.2 (13.9) times, 8.0 (4.5) years, and -7.8 (34.6) mL · yr⁻¹, respectively. The distribution of Δ FEV₁ in this population is shown in Figure E1 in the Online Repository. Within 2 years after diagnosis, 46% of patients started ICS treatment. At enrollment, 82% of patients took controllers such as long-acting β_2 agonists, leukotriene receptor antagonists, or sustained release theophylline to achieve adequate asthma control. Based on a questionnaire, adherence to medication was satisfactory; 49% of the participants never and 38% seldom forgot to take ICS or other medications. Based on ACT scores, 50% was totally controlled, and 38% scored from 20 to 24, indicating that they were well controlled at enrollment.

Serum periostin levels of asthmatic patients [92.8 (38.4) ng/mL] were significantly higher than those of healthy subjects [39.1 (24.5) ng/mL, $p < 0.001$]. The ROC curve analysis was performed to discriminate patients with asthma who were thought to have refractory Th2 inflammation despite long-term ICS treatment from healthy subjects. The highest specificity among the 4 cut-off values tested was achieved at 95 ng/mL (0.985) in the comparison study of 224 asthmatic patients and 66 healthy subjects. Therefore a cut-off value of 95 ng/mL was used to define a high serum periostin group, although it had relatively lesser sensitivity (0.379) (see Figure E2 in the Online Repository). In asthmatic patients, 85 patients (38%) had high serum periostin levels (≥ 95 ng/mL). Of the 85 patients, 40 patients (47%) were on

treatment step 4, according to the treatment step classification²⁷, and 9 patients (11%) were on treatment step 5.

Associations between serum periostin levels and greater annual decline in FEV₁ and a decline in FEV₁ of 30 mL·yr⁻¹ or greater

In an analysis of continuous values of Δ FEV₁, greater decline in FEV₁ was associated with higher serum periostin levels at enrollment, treatment step 5, lower ACT scores, incomplete adherence to medications, comorbid or a history of sinusitis, and comorbid diabetes mellitus (Table 3). When patients were stratified into two groups according to their serum periostin levels, high serum periostin (≥ 95 ng/mL) was also associated with greater decline in FEV₁ (Table 3). Of these, high serum periostin was significant after controlling for multiple testing using the false discovery rate ($q = 0.03$, data not shown in Table 3).³⁰ Multivariate analysis revealed that greater decline of FEV₁ was solely associated with high serum periostin (≥ 95 ng/mL) (estimated effect -5.39, 95% confidence interval -10.0 to -0.77, $p = 0.02$).

Fifty-two patients (23%) showed a decline in FEV₁ of 30 mL·yr⁻¹ or greater [mean -51.8 (18.4) mL·yr⁻¹] and were considered rapid decliners³¹. When adjusted by confounders, higher serum periostin levels at enrollment, treatment step 5, a history of admission due to asthma exacerbation, higher ICS daily doses, comorbid or a history of sinusitis, and ex-smoking were associated with a decline in FEV₁ of 30 mL·yr⁻¹ or greater. High serum periostin (≥ 95 ng/mL) was also associated with a decline in FEV₁ of 30 mL·yr⁻¹ or greater (Table 4). On multivariate analysis, high serum periostin (≥ 95 ng/mL), treatment step 5, and ex-smoking were independent risk factors for a decline in FEV₁ of 30 mL·yr⁻¹ or greater (Table 4).

Of the 224 patients, 19 patients were on treatment step 5, and 36 patients took high-dose ICS (1,000 μ g or higher doses of ICS equivalent to fluticasone propionate daily). When

patients were stratified into the high periostin group, the average ΔFEV_1 of patients on treatment step 5 ($n = 9$) was -41.0 (49.3) $\text{mL} \cdot \text{yr}^{-1}$, and 7 of them (78%) had excess decline; the average ΔFEV_1 of patients on high-dose ICS ($n=18$) was -34.3 (39.4) $\text{mL} \cdot \text{yr}^{-1}$, and 11 of them (61%) had a decline in FEV_1 of $30 \text{ mL} \cdot \text{yr}^{-1}$ or greater.

Serum periostin levels and clinical indices

In 224 patients, serum periostin levels were weakly associated with blood eosinophil counts (Figure 2), serum IgE (Figure 2) and ECP levels ($r = 0.25$, $p = 0.0005$), ICS-untreated period, i.e. period between onset of asthma and the initiation of ICS therapy ($r = 0.16$, $p = 0.01$), daily maintenance doses of ICS at enrollment ($r = 0.13$, $p = 0.05$), and a history of admission due to asthma exacerbation ($r = 0.15$, $p = 0.03$). Serum periostin levels were significantly higher in patients on high-dose ICS ($\geq 1,000 \mu\text{g}$ daily) than in the remaining patients (110.3 ng/mL vs. 89.5 ng/mL , $p = 0.003$). Lastly, serum periostin levels were higher in patients with sinusitis than in those without sinusitis (103.9 ng/mL vs. 88.3 ng/mL , $p = 0.007$). Serum periostin levels did not show any seasonal variability or association with age at onset of asthma (data not shown).

***POSTN* gene polymorphisms**

Associations between polymorphisms of the *POSTN* gene, which encodes periostin, and both serum periostin levels and pulmonary function decline were then investigated. In one patient, DNA quality was insufficient for genotyping; thus, 3 tag SNPs of the *POSTN* gene were analyzed in 223 patients. All genotyped data were in Hardy-Weinberg equilibrium. The frequencies of the 3 tag SNPs and analysis results using dominant and recessive models for serum periostin levels and a decline in FEV_1 of $30 \text{ mL} \cdot \text{yr}^{-1}$ or greater are presented in Table 5.

Serum periostin levels were higher in patients with the GG genotype of rs3829365 than

286 in those with the GC/CC genotype (GG 98.7 ng/mL vs. GC/CC 86.1 ng/mL, $p = 0.003$).
287 rs1028728 was not associated with serum periostin levels or with the frequency of rapid
288 decliners, but patients with the TT genotype of rs1028728, 4 patients only, showed no
289 significant decline compared with the AA/AT genotype (AA/AT $-8.6 \text{ mL} \cdot \text{yr}^{-1}$ vs. TT 29.3
290 $\text{mL} \cdot \text{yr}^{-1}$, $p = 0.03$). Rapid decliners were more frequently observed in patients with the minor
291 A allele of rs9603226 than in the GG genotype (GG 16% vs. AG/AA 30%, $p = 0.02$). A
292 marked difference in the frequency of rapid decliners was observed when patients were
293 stratified into the high periostin group [GG of rs9630226 ($n = 37$) 19% vs. AG/AA ($n = 47$)
294 45%, $p = 0.01$].

295 Discussion

296 To the best of our knowledge, this is the first study to identify a relationship between
297 greater decline in FEV₁ and higher serum periostin levels, particularly if they were 95 ng/mL
298 or more, in asthmatic patients on ICS treatment. It was also shown that high serum periostin,
299 together with treatment step 5 and light ex-smoking, was an independent risk factor for a
300 decline in FEV₁ of 30 mL·yr⁻¹ or greater. In addition, polymorphisms of the *POSTN* gene,
301 which encodes periostin, were associated with serum periostin levels and a decline in FEV₁
302 of 30 mL·yr⁻¹ or greater in asthmatic patients. These findings suggest that serum periostin
303 may be a useful biomarker for the development of airflow limitation in asthmatic patients on
304 ICS.

305 In this study, despite long-term treatment with ICS with or without other controllers,
306 23% of asthmatic patients were rapid decliners who showed a decline in FEV₁ of 30 mL·yr⁻¹
307 or greater, for which treatment step 5 was an independent risk factor. Adherence to ICS
308 treatment and the frequency of early intervention with ICS did not differ between rapid
309 decliners and non-decliners, although long-term adherence to ICS was undetermined in the
310 present study. In previous studies of patients who were not treated with ICS, severe
311 exacerbation of asthma contributed to greater annual decline of pulmonary function^{6,7}, but the
312 exacerbation-related greater annual decline disappeared in an early intervention group with
313 ICS treatment in the START study⁶, which might be interpreted to mean that asthmatic
314 patients on ICS treatment have little risk of accelerated FEV₁ decline. However, since the
315 START study originally recruited mild persistent asthmatic patients, its results cannot simply
316 be applied to severe asthmatic patients. As observed in the present study, there would be a
317 subset of asthmatic patients still at risk of greater annual decline of pulmonary function
318 despite intensive treatment for asthma.

319 Persistent eosinophilic airway inflammation is a key process in irreversible airway
320 obstruction¹⁰. Indeed, exhaled nitric oxide of 20 ppb or higher is a risk factor for accelerated

FEV₁ decline in patients with difficult-to-treat asthma¹⁸. Studies on novel therapies for refractory eosinophilic asthma, i.e., anti-IL-5 therapy³² and anti-IL-13 therapy³³, revealed that these treatments may reverse airway remodeling when patients are adequately targeted, suggesting the necessity of establishing “companion diagnostics” for this population. According to the most recent study, serum periostin is the single best biomarker reflecting sputum and tissue eosinophilia among several biomarkers, including blood eosinophils and exhaled nitric oxide²⁵. In the current study, the serum periostin level, which was associated with the blood eosinophil count, was the sole biomarker that reflected greater decline in FEV₁. Periostin is secreted by airway epithelial cells^{23, 24} and lung fibroblasts²¹ in response to IL-4 and IL-13 and is thought to be secreted into the capillary vessels. Downstream of IL-13, which plays a pivotal role in subepithelial airway fibrosis³⁴, airway remodeling³⁵, and steroid insensitivity³⁶, periostin mediates collagen synthesis²⁴ and fibrillogenesis^{24, 37} by binding to collagen³⁷ and activates TGF- β ²⁴. In the asthmatic airway, periostin is deposited in the subepithelial layer, colocalizing with collagens I, III, and V, fibronectin, tenascin-C, and periostin itself²¹, which indicates involvement of periostin in airway remodeling in asthma. Collectively, periostin may be a key molecule that links eosinophilic inflammation and remodeling *via* IL-13 in asthmatic airways. Further roles of periostin in allergic inflammation and remodeling in the airways remain undetermined because studies using periostin-deficient mice with acute allergen exposure have yielded conflicting findings³⁸⁻⁴⁰; one study showed that periostin facilitates eosinophil infiltration into the lung³⁸, whereas two other studies^{39, 40} suggested protective roles of periostin. Meanwhile, a recent study of a chronic mouse model of atopic dermatitis demonstrated periostin’s role in the chronicity of Th2 inflammation²⁹.

In the present study, patients on high-dose ICS showed higher serum periostin levels than the other patients. Although a longitudinal study is needed to determine responses of serum periostin levels to ICS treatment, we do not think that the high serum periostin levels in patients on high-dose ICS were induced by ICS treatment, because periostin expression in

the airway epithelium was decreased with ICS treatment²³. Rather, the elevation of serum periostin in this population may reflect IL-13-mediated inflammation that is partly refractory to ICS, as was reported in a recent study by Jia and colleagues²⁵. They showed that, in patients with severe asthma who were treated with high doses ICS (> 1000 µg daily), elevation of serum periostin levels was associated with persistent airway tissue eosinophilia, concluding that serum periostin is a systemic biomarker of airway eosinophilia refractory to high-dose ICS²⁵. Providing further support, among patients with moderate to severe asthma who are inadequately controlled despite ICS treatment, patients with high serum periostin levels are likely to benefit from anti-IL-13 antibody, lebrikizumab, treatment³³. The novelty of the present finding is that high serum periostin is an independent risk factor for greater decline in FEV₁, providing the first evidence for the potential association between persistent Th2- or IL-13-driven inflammation refractory to ICS treatment and greater decline in FEV₁, a functional consequence of airway remodeling.

Needless to say, current smokers with asthma have more accelerated FEV₁ decline⁴ than those not smoking, and current smoking impairs the therapeutic response to ICS or oral corticosteroids⁴¹. Meanwhile, smoking cessation improves their FEV₁ levels⁴², and ex-smokers with asthma with 10 pack-years or more show an intermediate response to short-term oral corticosteroid treatment, between current smokers and never-smokers⁴¹. In the present study, rather unexpectedly, ex-smoking with 10 pack-years or less was still an independent risk factor for a decline in FEV₁ of 30 mL·yr⁻¹ or greater. It should be recognized that even light ex-smoking increases the risk of airway remodeling in asthmatic patients on ICS, and its underlying mechanisms should be clarified.

Chronic sinusitis is a well-known comorbidity with severe asthma^{43, 44}. In the present study, rapid decliners were more frequently observed in asthmatic patients with sinusitis than those without sinusitis on univariate analysis, and their periostin levels were higher than in patients without sinusitis. In the present study, polypoid lesions in the sinuses were not

evaluated by otolaryngologists at enrollment. However, considering that periostin is up-regulated in nasal polyp tissue in patients with chronic rhinosinusitis⁴⁵, asthmatic patients with sinusitis may have had severe upper and lower airway inflammation with persistent increases in periostin expression, which may have resulted in a decline in FEV₁ of 30 mL·yr⁻¹ or greater. Periostin is a potential molecule that unifies sinusitis and severe asthma.

Periostin is encoded on the *POSTN* gene, which is located on chromosome 13q13.3. rs3829365, which is located at the 5'UTR region that may contain sequences to regulate translation efficiency or mRNA stability, was associated with serum periostin levels. This finding suggests that, besides IL-13, a master regulator of periostin, genetic background partly determines periostin levels, although a replication study would be necessary to confirm this. The minor A allele of rs9603226, located 66 bp upstream of exon 21 in the C-terminal region, was associated with a decline in FEV₁ of 30 mL·yr⁻¹ or greater. In periostin, FAS I domains are thought to be primary binding sites to fibronectin, tenascin-C, and collagen V²¹, whereas the C-terminal region in its intact form may down-regulate the binding activity of periostin to these extracellular matrix proteins²¹. We therefore speculate that the minor A allele of rs9603226 might modify the binding activity at the C-terminal region and facilitate airway remodeling, particularly if the airway is in periostin enriched milieu. Further studies are needed to clarify if these SNPs are functional variants.

The age of patients in this study appears to be older than in other Euro-American studies^{6,7,14,18,20,23,25}. One reason for the age distribution would be the entry criteria of this study. Another reason would be explained by population aging including population with asthma in Japan. According to a patient survey by the Japanese Ministry of Health, Labour and Welfare in 2008, patients aged 70 to 74 years were the most frequent age group of adult patients with asthma⁴⁶, which is still older than the average age of patients in this study.

There are several limitations to the present study. First, since this study was observational in nature, ICS doses and numbers or types of controllers were not fixed during

the follow-up period. Controllers such as long-acting β_2 agonists were not withdrawn at pulmonary function testing to evaluate function on daily medications, which may have resulted in the small average ΔFEV_1 , $-7.8 \text{ mL} \cdot \text{yr}^{-1}$. Meanwhile, averages of 16.2 measurements of FEV_1 and 8.0 years of follow-up were satisfactory for a longitudinal analysis of pulmonary function⁴⁷, and ΔFEV_1 was normally distributed. Secondly, serum biomarkers were measured only once at enrollment, but the significant associations between *POSTN* gene polymorphisms and serum periostin levels or a decline in FEV_1 of $30 \text{ mL} \cdot \text{yr}^{-1}$ or greater may circumvent the inherent insufficiency of single measurement of serum periostin. Thirdly, most of the clinical information, including smoking history and chronic sinusitis, was based on a self-completed questionnaire, which might be biased by recall memory. Despite these limitations, the current findings may provide directions for future research.

In conclusion, serum periostin appears to be a useful biomarker that reflects the development of airflow limitation in patients on prolonged treatment with ICS. *POSTN* gene polymorphisms may also be helpful for identification of rapid decliners.

414 **Acknowledgments**

415 The authors would like to acknowledge Dr Nobuo Ohta, Department of
416 Otolaryngology, Head and Neck Surgery, Yamagata University for fruitful discussion on
417 periostin expression in nasal tissue of chronic sinusitis. The authors would also like to thank
418 Ms Maki Futamata (Saga Medical School), Dr Guergana Petrova Stoyanova, Dr Cui Shilei,
419 Ms Aya Inazumi, and Ms Yuko Maeda (Kyoto University) for their technical assistance.

Reference

1. Pascual RM, Peters SP. Airway remodeling contributes to the progressive loss of lung function in asthma: an overview. *J Allergy Clin Immunol* 2005; 116:477-86; quiz 87.
2. Niimi A, Matsumoto H, Takemura M, Ueda T, Chin K, Mishima M. Relationship of airway wall thickness to airway sensitivity and airway reactivity in asthma. *Am J Respir Crit Care Med* 2003; 168:983-8.
3. Ueda T, Niimi A, Matsumoto H, Takemura M, Hirai T, Yamaguchi M, et al. Role of small airways in asthma: investigation using high-resolution computed tomography. *J Allergy Clin Immunol* 2006; 118:1019-25.
4. Lange P, Parner J, Vestbo J, Schnohr P, Jensen G. A 15-year follow-up study of ventilatory function in adults with asthma. *N Engl J Med* 1998; 339:1194-200.
5. Ulrik CS. Outcome of asthma: longitudinal changes in lung function. *Eur Respir J* 1999; 13:904-18.
6. O'Byrne PM, Pedersen S, Lamm CJ, Tan WC, Busse WW, Group SI. Severe exacerbations and decline in lung function in asthma. *Am J Respir Crit Care Med* 2009; 179:19-24.
7. Bai TR, Vonk JM, Postma DS, Boezen HM. Severe exacerbations predict excess lung function decline in asthma. *Eur Respir J* 2007; 30:452-6.
8. Ulrik CS, Lange P. Decline of lung function in adults with bronchial asthma. *Am J Respir Crit Care Med* 1994; 150:629-34.
9. Ulrik CS, Backer V, Dirksen A. A 10 year follow up of 180 adults with bronchial asthma: factors important for the decline in lung function. *Thorax* 1992; 47:14-8.
10. ten Brinke A, Zwinderman AH, Sterk PJ, Rabe KF, Bel EH. Factors associated with persistent airflow limitation in severe asthma. *Am J Respir Crit Care Med* 2001; 164:744-8.
11. Jongepier H, Boezen HM, Dijkstra A, Howard TD, Vonk JM, Koppelman GH, et al. Polymorphisms of the ADAM33 gene are associated with accelerated lung function decline in asthma. *Clin Exp Allergy* 2004; 34:757-60.
12. Koppelman GH, Sayers I. Evidence of a genetic contribution to lung function decline in asthma. *J Allergy Clin Immunol* 2011; 128:479-84.
13. Tantisira KG, Lasky-Su J, Harada M, Murphy A, Litonjua AA, Himes BE, et al. Genomewide association between GLCCI1 and response to glucocorticoid therapy in asthma. *N Engl J Med* 2011; 365:1173-83.
14. Dijkstra A, Vonk JM, Jongepier H, Koppelman GH, Schouten JP, ten Hacken NH, et al. Lung function decline in asthma: association with inhaled corticosteroids, smoking and sex. *Thorax* 2006; 61:105-10.
15. O'Byrne PM, Pedersen S, Busse WW, Tan WC, Chen YZ, Ohlsson SV, et al. Effects of early intervention with inhaled budesonide on lung function in newly diagnosed

- asthma. *Chest* 2006; 129:1478-85.
16. Selroos O, Löfroos AB, Pietinalho A, Riska H. Asthma control and steroid doses 5 years after early or delayed introduction of inhaled corticosteroids in asthma: a real-life study. *Respir Med* 2004; 98:254-62.
17. de Marco R, Marcon A, Jarvis D, Accordini S, Bugiani M, Cazzoletti L, et al. Inhaled steroids are associated with reduced lung function decline in subjects with asthma with elevated total IgE. *J Allergy Clin Immunol* 2007; 119:611-7.
18. van Veen IH, Ten Brinke A, Sterk PJ, Sont JK, Gauw SA, Rabe KF, et al. Exhaled nitric oxide predicts lung function decline in difficult-to-treat asthma. *Eur Respir J* 2008; 32:344-9.
19. Levine SJ, Wenzel SE. Narrative review: the role of Th2 immune pathway modulation in the treatment of severe asthma and its phenotypes. *Ann Intern Med* 2010; 152:232-7.
20. Woodruff PG, Modrek B, Choy DF, Jia G, Abbas AR, Ellwanger A, et al. T-helper type 2-driven inflammation defines major subphenotypes of asthma. *Am J Respir Crit Care Med* 2009; 180:388-95.
21. Takayama G, Arima K, Kanaji T, Toda S, Tanaka H, Shoji S, et al. Periostin: a novel component of subepithelial fibrosis of bronchial asthma downstream of IL-4 and IL-13 signals. *J Allergy Clin Immunol* 2006; 118:98-104.
22. Kii I, Nishiyama T, Li M, Matsumoto K, Saito M, Amizuka N, et al. Incorporation of tenascin-C into the extracellular matrix by periostin underlies an extracellular meshwork architecture. *J Biol Chem* 2010; 285:2028-39.
23. Woodruff PG, Boushey HA, Dolganov GM, Barker CS, Yang YH, Donnelly S, et al. Genome-wide profiling identifies epithelial cell genes associated with asthma and with treatment response to corticosteroids. *Proc Natl Acad Sci U S A* 2007; 104:15858-63.
24. Sidhu SS, Yuan S, Innes AL, Kerr S, Woodruff PG, Hou L, et al. Roles of epithelial cell-derived periostin in TGF-beta activation, collagen production, and collagen gel elasticity in asthma. *Proc Natl Acad Sci U S A* 2010; 107:14170-5.
25. Jia G, Erickson RW, Choy DF, Mosesova S, Wu LC, Solberg OD, et al. Periostin is a systemic biomarker of eosinophilic airway inflammation in asthmatic patients. *J Allergy Clin Immunol* 2012; 130: 647-54.
26. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease (COPD) and asthma. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, November 1986. *Am Rev Respir Dis* 1987; 136:225-44.
27. Global Strategy for Asthma Management and Prevention, Global Initiative for Asthma (GINA) 2010. Available from: Global Strategy for Asthma Management and

Prevention, Global Initiative for Asthma (GINA) 2010. Available from:
<http://www.ginasthma.org>.

28. Okamoto M, Hoshino T, Kitasato Y, Sakazaki Y, Kawayama T, Fujimoto K, et al. Periostin, a matrix protein, is a novel biomarker for idiopathic interstitial pneumonias. *Eur Respir J* 2011; 37:1119-27.
29. Masuoka M, Shiraishi H, Ohta S, Suzuki S, Arima K, Aoki S, et al. Periostin promotes chronic allergic inflammation in response to Th2 cytokines. *J Clin Invest* 2012; 122:2590-600.
30. Storey JD, Tibshirani R. Statistical significance for genomewide studies. *Proc Natl Acad Sci U S A* 2003; 100:9440-5.
31. Broekema M, Volbeda F, Timens W, Dijkstra A, Lee NA, Lee JJ, et al. Airway eosinophilia in remission and progression of asthma: accumulation with a fast decline of FEV₁. *Respir Med* 2010; 104:1254-62.
32. Haldar P, Brightling CE, Hargadon B, Gupta S, Monteiro W, Sousa A, et al. Mepolizumab and exacerbations of refractory eosinophilic asthma. *N Engl J Med* 2009; 360:973-84.
33. Corren J, Lemanske RF, Hanania NA, Korenblat PE, Parsey MV, Arron JR, et al. Lebrikizumab treatment in adults with asthma. *N Engl J Med* 2011; 365:1088-98.
34. Zhu Z, Homer RJ, Wang Z, Chen Q, Geba GP, Wang J, et al. Pulmonary expression of interleukin-13 causes inflammation, mucus hypersecretion, subepithelial fibrosis, physiologic abnormalities, and eotaxin production. *J Clin Invest* 1999; 103:779-88.
35. Elias JA, Zhu Z, Chupp G, Homer RJ. Airway remodeling in asthma. *J Clin Invest* 1999; 104:1001-6.
36. Saha SK, Berry MA, Parker D, Siddiqui S, Morgan A, May R, et al. Increased sputum and bronchial biopsy IL-13 expression in severe asthma. *J Allergy Clin Immunol* 2008; 121:685-91.
37. Norris RA, Damon B, Mironov V, Kasyanov V, Ramamurthi A, Moreno-Rodriguez R, et al. Periostin regulates collagen fibrillogenesis and the biomechanical properties of connective tissues. *J Cell Biochem* 2007; 101:695-711.
38. Blanchard C, Mingler MK, McBride M, Putnam PE, Collins MH, Chang G, et al. Periostin facilitates eosinophil tissue infiltration in allergic lung and esophageal responses. *Mucosal Immunol* 2008; 1:289-96.
39. Gordon ED, Sidhu SS, Wang ZE, Woodruff PG, Yuan S, Solon MC, et al. A protective role for periostin and TGF- β in IgE-mediated allergy and airway hyperresponsiveness. *Clin Exp Allergy* 2012; 42:144-55.
40. Sehra S, Yao W, Nguyen ET, Ahyi AN, Tuana FM, Ahlfeld SK, et al. Periostin regulates goblet cell metaplasia in a model of allergic airway inflammation. *J Immunol* 2011; 186:4959-66.

41. Thomson NC, Chaudhuri R, Livingston E. Asthma and cigarette smoking. *Eur Respir J* 2004; 24:822-33.
42. Chaudhuri R, Livingston E, McMahon AD, Lafferty J, Fraser I, Spears M, et al. Effects of smoking cessation on lung function and airway inflammation in smokers with asthma. *Am J Respir Crit Care Med* 2006; 174:127-33.
43. Dixon AE, Kaminsky DA, Holbrook JT, Wise RA, Shade DM, Irvin CG. Allergic rhinitis and sinusitis in asthma: differential effects on symptoms and pulmonary function. *Chest* 2006; 130:429-35.
44. Mascia K, Borish L, Patrie J, Hunt J, Phillips CD, Steinke JW. Chronic hyperplastic eosinophilic sinusitis as a predictor of aspirin-exacerbated respiratory disease. *Ann Allergy Asthma Immunol* 2005; 94:652-7.
45. Ishida A, Ohta N, Suzuki Y, Kakehata S, Okubo K, Ikeda H, et al. Expression of Pendrin and Periostin in Allergic Rhinitis and Chronic Rhinosinusitis. *Allergol Int* 2012; 61: 589-95.
46. Japanese Society of Allergology, Asthma Guideline Committee. Asthma Prevention and Management Guidelines 2012. Tokyo: Kyowa Kikaku; 2012 (in Japanese)
47. Wang ML, Gunel E, Petsonk EL. Design strategies for longitudinal spirometry studies: study duration and measurement frequency. *Am J Respir Crit Care Med* 2000; 162:2134-8.

555 **Table 1. Contents of the self-completed questionnaire**

Asthma-related history ▪ family history of asthma ▪ age of asthma onset ▪ history of pediatric asthma ▪ history of admission due to asthma worsening or exacerbation ▪ aspirin hypersensitivity ▪ asthma deterioration at the working place	
Comorbidity or a history of the following diseases ▪ allergic dermatitis ▪ allergic rhinitis ▪ seasonal rhinitis ▪ allergic conjunctivitis ▪ chronic sinusitis ▪ cardiovascular diseases including ischemic heart disease ▪ gastrointestinal diseases including GERD ▪ collagen vascular diseases including rheumatoid arthritis ▪ diabetes mellitus ▪ pulmonary diseases other than asthma ▪ other diseases including malignancy	
Lifestyle and environment ▪ smoking history ▪ pet breeding ▪ type of occupation ▪ a highway near the home ▪ age at menopause	
Adherence to medication, sputum production, and exacerbations ▪ How often do you forget to take inhaled corticosteroids or other medications? 0: never, 1: seldom, 2: sometimes, 3: often, 4: always ▪ How often do you produce sputum? 0: never, 1: once in a few days, 2: every morning, 3: every morning and daytime ▪ How often did you receive systemic steroids due to asthma exacerbations during the recent 6 months? 0: never, 1: once, 2: twice or more	

556 GERD: gastro-esophageal reflux disease

557

558 **Table 2. Patients' characteristics**

Sex (males/ females), n	53 / 171
Age at enrollment, years	62.3 (13.7)
Age at asthma onset, years	42.0 (19.0)
Body mass index (kg/m ²)	23.1 (3.5)
Smoking history (never), n	181
Atopic predisposition [*] , %	70
Pediatric asthma (none/ recurrent/ persistent), %	81 / 8 / 11
Disease duration, years	20.2 (14.5)
ICS-untreated period, years	9.2 (13.1)
ICS daily maintenance dose [†] , µg	525 (318)
Number of other controller medications, n	1.4 (1.2)
Treatment step (2/ 3/ 4/ 5) [‡] , %	16 / 27 / 49 / 8
Sputum production (0/ 1/ 2/ 3) [§] , %	54 / 20 / 8 / 18
Asthma Control Test, points	22.6 (3.5)
History of admission due to asthma, n (%)	78 (35)
Allergic rhinitis, n (%)	129 (58)
Chronic sinusitis, n (%)	65 (29)
Blood neutrophils, %	60.1 (10.0)
eosinophils, %	5.2 (4.9)
Serum IgE, IU/mL	180 (0 - 16000)
periostin, ng/mL	92.8 (38.4)
high sensitivity C-reactive protein, mg/L	1341 (3147)
eosinophil cationic protein, µg/L	15.1 (29.3)
FEV ₁ at the first measurement, L [¶]	2.11 (0.69)
%predicted FEV ₁ at the first measurement, %	91.9 (19.2)
FEV ₁ / FVC at the first measurement, %	73.9 (9.8)
FEV ₁ at enrollment, L	2.04 (0.73)
%predicted FEV ₁ at enrollment, %	97.4 (22.2)
FEV ₁ / FVC at enrollment, %	72.2 (10.0)
Reversibility at enrollment, % [#]	3.8 (6.0)

559 Data at enrollment are presented unless otherwise stated. Data are expressed as means (SD) except for median
560 (range) for serum IgE. ^{*}Considered atopic when one or more specific IgE antibodies against cat or dog dander,
561 weed, grass, or Japanese cedar pollens, moulds, or house dust mite were positive. [†]Equivalent to fluticasone
562 propionate. [‡]according to the Global Initiative for Asthma 2010 guideline²⁷. [§]0 = never, the details are shown in
563 Table 1. [¶]The first pulmonary function test was performed at least one year after the commencement of ICS
564 treatment and at 25 years of age or older. [#]n = 206, airway reversibility to 200 µg of inhaled salbutamol.

565 **Table 3. Estimated effects of clinical indices and biomarkers on Δ FEV₁**

	Estimates	95% C.I.	p value
Smoking history, ex vs. never	-8.48	-20.2, 3.27	0.16
Atopic predisposition	-1.10	-6.29, 4.09	0.68
Disease duration, years	-4.79	-18.4, 8.86	0.56
ICS-untreated period, years	0.10	-0.24, 0.45	0.65
ICS daily maintenance dose, μ g	-0.01	-0.03, 0.001	0.07
Number of other controller medications, n	-0.36	-4.21, 3.49	0.86
Adherence to medication, incomplete vs. complete*	-4.56	-9.08, -0.04	0.05
Treatment step, 5 vs. 2-4†	-7.77	-15.7, 0.13	0.05
Sputum production, never vs. others‡	0.99	-3.53, 5.51	0.67
Asthma Control Test, points	1.53	0.29, 2.77	0.02
History of admission due to asthma	-4.49	-9.45, 0.46	0.08
Aspirin hypersensitivity	-6.52	-20.0, 6.98	0.34
Asthma deterioration at the working place	-12.2	-54.4, 30.0	0.57
Allergic rhinitis	-1.21	-5.88, 3.45	0.61
Allergic dermatitis	4.51	-1.51, 10.5	0.14
Chronic sinusitis	-10.1	-19.8, -0.27	0.04
Ischemic heart disease	3.41	-16.6, 23.4	0.74
Hypertension	-3.79	-9.12, 1.53	0.16
Dyslipidemia	-3.67	-9.42, -2.06	0.21
Diabetes mellitus	-8.03	-15.4, -0.67	0.03
Gastro-esophageal reflux disease	-3.85	-9.89, 2.19	0.21
Malignancy	-3.44	-26.0, 19.1	0.76
Post-menopause	5.05	-14.2, 24.3	0.60
Pet breeding	-0.28	-12.6, 12.0	0.96
Log blood neutrophils, %	-7.40	-69.1, 54.3	0.81
eosinophils, %	-0.67	-1.60, 0.27	0.16
Log serum IgE, IU/mL	-2.85	-9.74, 4.04	0.42
periostin, ng/mL	-29.1	-56.2, -1.97	0.04
high sensitivity C-reactive protein, mg/L	-1.88	-9.85, 6.10	0.64
eosinophil cationic protein, μ g/L	-4.47	-15.7, 6.81	0.44
Periostin group, high vs. low§	-6.96	-11.4, -2.51	0.002

566 Estimated effects were adjusted by sex, height, age at enrollment, and FEV₁ at the first measurement. * “Complete”, when patients answered
567 that they never forgot to take ICS or other medications; “incomplete”, the remaining cases. † according to the Global Initiative for Asthma
568 2010 guideline²⁷. ‡ The details are shown in Table 1. § Patients were stratified into two groups according to their serum periostin levels: high $\geq$
569 95 ng/mL, low < 95 ng/mL. ICS: inhaled corticosteroids, C.I.: confidence interval

570 **Table 4. Estimated effects of clinical indices and serum periostin on a decline in FEV₁ of**
 571 **30 mL·yr⁻¹ or greater**

	Univariate analysis			Multivariate analysis		
	Estimates	95% C.I.	p value	Estimates	95% C.I.	p value
Treatment step, 5 vs. 2-4*	1.63	0.51, 2.60	0.004	1.24	0.078, 2.30	0.04
History of admission due to asthma	1.09	0.37, 1.90	0.003	0.70	-0.11, 1.50	0.09
ICS daily maintenance dose, µg	0.001	0.00, 0.002	0.01	-		
Chronic sinusitis	0.82	0.11, 1.53	0.03	0.61	-0.15, 1.37	0.12
Smoking history, ex vs. never	0.87	-0.002, 1.74	0.05	0.98	0.030, 1.93	0.04
Log serum periostin, ng/mL	2.96	0.78, 5.13	0.008	-		
Periostin group, high vs. low†	1.03	0.33, 1.72	0.004	0.87	0.11, 1.63	0.03

572 Estimated effects were adjusted by sex, height, age at enrollment, and FEV₁ at the first measurement.

573 * according to the Global Initiative for Asthma 2010 guideline²⁷.

574 †Patients were stratified into two groups according to their serum periostin levels: high ≥ 95 ng/mL, low <

575 95 ng/mL. ICS: inhaled corticosteroids, C.I.: confidence interval

576 ICS daily maintenance dose was excluded from multivariate analysis because of its strong correlation with

577 treatment step.

578

Table 5. Frequencies of 3 tag SNPs and analysis results using dominant and recessive models for serum periostin levels and frequency of rapid decliners*

Tag SNP	Genotype	n (%)	Allelic	n (%)	Serum periostin levels		Frequency of rapid decliners	
					p value		p value	
					Dominant [†]	Recessive [‡]	Dominant [†]	Recessive [‡]
rs1028728	AA	164 (74)	A	383 (86)				
	AT	55 (25)	T	63 (14)	0.40	0.46	0.17	0.14
	TT	4 (2)						
rs3829365	GG	113 (51)	G	316 (71)				
	GC	90 (40)	C	130 (29)	0.003	0.70	0.40	0.33
	CC	20 (9)						
rs9603226	GG	107 (48)	G	311 (70)				
	AG	97 (44)	A	135 (30)	0.80	0.33	0.01	0.81
	AA	19 (9)						

* defined as patients who showed a decline in FEV₁ of 30 mL·yr⁻¹ or greater

[†] Assuming that heterozygotes have the same increased risk as minor homozygous genotypes.

[‡] Assuming that heterozygotes have no increased risk.

585 **Figure legends**

586 Figure 1. Three tag SNPs that determine 4 major haplotypes of the *POSTN* gene and
587 haplotype frequencies in the Japanese population are presented.


588 *at intron 66 bp upstream of exon 21

589

590 Figure 2. Relationships between serum periostin levels and blood eosinophil counts (left) or
591 serum IgE levels (right).

592 Presented in logarithmic scales on both the X- and Y-axes.

593



594 Figure 1.




Figure 2.

Online Repository

Increased periostin associates with greater airflow limitation in patients receiving inhaled corticosteroids

Yoshihiro Kanemitsu MD^{*1,2}, Hisako Matsumoto MD, PhD^{*1,2}, Kenji Izuhara MD, PhD³, Yuji Tohda MD, PhD^{2,4}, Hideo Kita MD, PhD^{2,5}, Takahiko Horiguchi MD, PhD^{2,6}, Kazunobu Kuwahara MD, PhD^{2,6}, Keisuke Tomii MD, PhD^{2,7}, Kojiro Otsuka MD, PhD^{1,2,7}, Masaki Fujimura MD, PhD^{2,8}, Noriyuki Ohkura MD^{2,8}, Katsuyuki Tomita MD, PhD^{2,4}, Akihito Yokoyama MD, PhD^{2,9}, Hiroshi Ohnishi MD, PhD^{2,9}, Yasutaka Nakano MD, PhD^{2,10}, Tetsuya Oguma MD, PhD^{2,10}, Soichiro Hozawa MD, PhD^{2,11}, Tadao Nagasaki MD¹, Isao Ito MD, PhD¹, Tsuyoshi Oguma MD¹, Hideki Inoue MD¹, Tomoko Tajiri MD¹, Toshiyuki Iwata MD¹, Yumi Izuhara MD¹, Junya Ono BS¹², Shoichiro Ohta MD, PhD¹³, Mayumi Tamari MD, PhD¹⁴, Tomomitsu Hirota DDS, PhD¹⁴, Tetsuji Yokoyama MD, PhD¹⁵, Akio Niimi MD, PhD^{1,2,16} and Michiaki Mishima MD, PhD^{1,2}.

Affiliations

¹ Department of Respiratory Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan.

² Kinki Hokuriku Airway disease Conference (KiHAC)

³ Division of Medical Biochemistry, Department of Biomolecular Sciences, Saga Medical School, Saga, Japan.

⁴ Department of Respiratory Medicine and Allergology, Faculty of Medicine, Kinki University, Osaka, Japan.

⁵ Department of Respiratory Medicine, Takatsuki Red Cross Hospital, Osaka, Japan.

⁶ Department of Respiratory Internal Medicine, Fujita Health University Second Educational Hospital, Aichi, Japan.

⁷ Department of Respiratory Medicine, Kobe City Medical Center General Hospital, Hyogo, Japan.

624 ⁸ Department of Respiratory Medicine, Cellular Transplantation Biology, Kanazawa
625 University Graduate School of Medicine, Kanazawa, Japan.

626 ⁹ Department of Hematology and Respiratory Medicine, Kochi University, Kochi, Japan.

627 ¹⁰ Division of Respiratory Medicine, Department of Internal Medicine, Shiga University of
628 Medical Science, Shiga, Japan.

629 ¹¹ Hiroshima Allergy and Respiratory Clinic, Hiroshima, Japan.

630 ¹² Shino-Test Corporation, Kanagawa, Japan.

631 ¹³ Department of Laboratory Medicine, Saga Medical School, Saga, Japan.

632 ¹⁴ Laboratory for Respiratory Diseases, Center for Genomic Medicine, RIKEN, Yokohama,
633 Kanagawa, Japan.

634 ¹⁵ Department of Health Promotion, National Institute of Public Health, Wako, Saitama,
635 Japan

636 ¹⁶ Department of Medical Oncology and Immunology, Nagoya City University School of
637 Medical Sciences, Aichi, Japan.

638

639 *YK and HM similarly contributed to this study.

640

641 Correspondence should be addressed to Hisako Matsumoto M.D., Ph.D.,

642 Department of Respiratory Medicine, Graduate School of Medicine, Kyoto University, 53

643 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan.

644 E-mail: hmatsumo@kuhp.kyoto-u.ac.jp

645

Methods

Patients

Patients with asthma were recruited from nine institutions belonging to the Kinki Hokuriku Airway disease Conference where asthma specialists manage patients, including six university hospitals, two satellite general hospitals, and one satellite clinic. Asthma was diagnosed according to the American Thoracic Society criteria^{E1} on the basis of a history of recurrent episodes of wheezing and chest tightness with or without cough and documented airway reversibility to a bronchodilator or hyper-responsiveness to inhaled methacholine. From September 2009 to December 2011, patients were enrolled if they had received ICS treatment for 4 years or more, undergone three or more pulmonary function tests when they were stable, and were free from exacerbations for at least one month. The first pulmonary function test was performed at least one year after the commencement of ICS treatment and at 25 years of age or older. Patients who had smoked more than 10 pack-years, smoked in the past one year, or had other pulmonary diseases were excluded.

Self-completed questionnaire and clinical indices

The self-completed questionnaire was composed of 4 major items, as presented in Table 1.

Adherence to ICS or other medications, frequency of sputum production, and requirement for systemic corticosteroids during the last 6 months were graded as shown in Table 1. The Asthma Control Test (ACT)TM was also scored. Duration of ICS treatment and details on medication at enrollment were recorded from medical charts by patients' physicians. The treatment step at enrollment was determined according to the Global Initiative for Asthma 2010 guideline^{E2}.

Measurement of systemic biomarkers

Blood eosinophil and neutrophil counts, and serum levels of total immunoglobulin E (IgE) (ImmunoCAP[®] total IgE, Phadia K.K., Tokyo, Japan), specific IgE against common inhaled allergens (ImmunoCAP[®] specific IgE), eosinophil cationic protein (ECP) (ImmunoCAP[®] ECP), high sensitivity C-reactive protein (hsCRP) (CardioPhase[®] hsCRP, Siemens Healthcare Diagnostics K.K., Tokyo, Japan), and periostin were determined.

Serum periostin levels were measured using an enzyme-linked immunosorbent assay at Shino-test (Kanagawa, Japan), as described previously^{E3}. Briefly, two rat anti-human periostin monoclonal antibodies (SS18A and SS17B) were used. SS18A and SS17B are antibodies against the first and fourth FAS I domains, respectively. Intra- and inter-assay coefficients of variation ranged from 1.31% to 2.54% and 1.49% to 2.01%, respectively.

Haplotype analysis, DNA extraction, and genotyping of the *POSTN* gene

A total of 47 single-nucleotide polymorphisms (SNPs) in the region of the *POSTN* gene and its upstream, total 39 kb, was captured in the HapMap Japanese data set with minor allele frequencies > 0.10. Pairwise tagging was performed at $r^2 > 0.8$ using a tagger in Haploview 4.2 software. Haplotype analysis identified 4 major haplotypes and 2 minor haplotypes. Two minor haplotypes were grouped into the closest major haplotype, and 3 tag SNPs that determined the 4 haplotypes were identified (Figure 1). These 3 tag SNPs were located at promoter region (rs1028728), 5'UTR region (rs3829365), and at intron 66 bp upstream of exon 21 (rs9603226). The frequencies of the minor alleles in the Japanese population were 0.136 (rs1028728), 0.278 (rs3829365), and 0.330 (rs9603226).

Genomic DNA was isolated from blood cells using a QIAamp DNA Blood Mini Kit (Qiagen, Tokyo, Japan). SNPs were genotyped using a Taqman genotyping assay according to the manufacturer's instructions (Applied Biosystems, Tokyo, Japan) and analyzed using an Applied Biosystems 7300 Real-Time PCR System (Applied Biosystems).

References

- E1. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease (COPD) and asthma. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, November 1986. *Am Rev Respir Dis* 1987; 136:225-44.
- E2 Global Strategy for Asthma Management and Prevention, Global Initiative for Asthma (GINA) 2010. Available from: Global Strategy for Asthma Management and Prevention, Global Initiative for Asthma (GINA) 2010. Available from: <http://www.ginasthma.org>.
- E3. Okamoto M, Hoshino T, Kitasato Y, Sakazaki Y, Kawayama T, Fujimoto K, et al. Periostin, a matrix protein, is a novel biomarker for idiopathic interstitial pneumonias. *Eur Respir J* 2011; 37:1119-27.

713 Figure legends

714

715 Figure E1. The distribution of ΔFEV_1 in the study population

716

717 Figure E2. ROC curve analysis of serum periostin levels comparing asthmatic patients and

718 healthy subjects, in which the cutoffs of 95 ng/mL, 80 ng/mL, 92 ng/mL, and 100 ng/mL are

719 presented with arrows.

720

721





Figure E1.



724 Figure E2.