

Alternative Theorems for Set-Valued Maps

新潟大学大学院 自然科学研究科 小野塚 真紀 (Onodsuka, Maki)*
Graduate School of Science and Technology, Niigata University

新潟大学大学院 自然科学研究科 西澤 正悟 (Nishizawa, Shogo)[†]
Graduate School of Science and Technology, Niigata University

新潟大学大学院 自然科学研究科 田中 環 (Tanaka, Tamaki)[‡]
Graduate School of Science and Technology, Niigata University

Abstract: Based on a comparison of each image of a set-valued map with the zero vector with respect to a given convex cone, we establish five types of alternative theorems for set-valued maps without any convexity assumption, which are proved by a nonlinear scalarization technique. As an application, we obtain optimality conditions for vector optimization problems with set-valued maps.

Key words: Alternative theorem, nonlinear scalarization, vector optimization, set-valued optimization, set-valued maps, optimality conditions.

1 Introduction

This paper is concerned with alternative theorems for set-valued maps based on a nonlinear scalarization. Alternative theorems of the Farkas and Gordan types play important roles in many applications, especially in optimization theory concerning optimality conditions for nonconvex programming problems and duality theory of these problems. A generalized Gordan alternative theorem was given for a vector-valued function by Jeyakumar [8] in 1986, and its generalization to set-valued maps was proved by Li [10] in 1999 and Yang et al. [17] in 2000. These results rely

2000 Mathematic Subject Classification. Primary: 90C29; Secondary: 90C46, 49J53.

*E-mail: maki@m.sc.niigata-u.ac.jp

[†]E-mail: shogo@m.sc.niigata-u.ac.jp

[‡]E-mail: tamaki@math.sc.niigata-u.ac.jp




Figure 1: Five types of classification for comparison between the zero vector and each image of multifunction with respect to cones.

on certain convexity assumptions like cone-subconvexlikeness in order to adopt a separation approach; see also [2, 6] for alternative theorems of set-valued maps. If we look at this approach from a different point of view, we will know that those proofs are based on a linear scalarization like an inner product. On the one hand, a nonlinear scalarization for vector-valued functions was introduced and applied to nonconvex separation theorems by Gerth (Tammer) and Weidner [5] in 1990, and similar approaches have been taken for several applications in [1, 3, 4, 15, 16] but at the same time we have researched some fundamental properties of a specific form of those nonlinear scalarizations in [13, 14]. By using special scalarizing functions under this type of nonlinear scalarization, we establish alternative theorems for set-valued maps without any convexity assumption.

In this paper, based on comparison between a vector and a set, we show five types of alternative theorems for set-valued maps; see also [9] for a comparison method between two sets. When comparing the zero vector and each image of a set-valued map (multifunction) with respect to a given dominance cone, there are five types of relationships as illustrated in Figure 1. Under this basic policy, we establish five types of alternative theorems 3.1–3.5 with respect to the interior of a convex cone in the sense of weak efficiency. Besides, we present five types of alternative theorems 3.6–3.10 with respect to the closure of a convex cone in the sense of strong efficiency.

2 Nonlinear Scalarization

In this section, we introduce a nonlinear scalarization for set-valued maps and show some properties that a characteristic function and scalarizing functions have.

Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. We assume that $C \neq Y$, which is equivalent to

$$\text{int } C \cap (-\text{cl } C) = \emptyset \quad (2.1)$$

for a convex cone with nonempty interior.

To begin with, we define a characteristic function

$$h_C(y; k) := \inf\{t : y \in tk - C\}$$

where $k \in \text{int } C$ and moreover $-h_C(-y; k) = \sup\{t : y \in tk + C\}$. This function $h_C(y; k)$ has been treated in some papers and which is regarded as a generalization of the Tchebyshev scalarization. Essentially, $h_C(y; k)$ is equivalent to the smallest strictly monotonic function defined by Luc in [11]. Note that $h_C(\cdot; k)$ is positively homogeneous and subadditive for every fixed $k \in \text{int } C$.

Now, we give some useful properties of this function h_C .

Lemma 2.1 *Let $y \in Y$, then the following statements hold:*

- (i) *If $y \in -\text{int } C$, then $h_C(y; k) < 0$ for all $k \in \text{int } C$;*
- (ii) *If there exists $k \in \text{int } C$ with $h_C(y; k) < 0$, then $y \in -\text{int } C$.*

Proof. First we prove the statement (i). Suppose that $y \in -\text{int } C$, then there exists an absorbing neighborhood V_0 of 0 in Y such that $y + V_0 \subset -\text{int } C$. Since V_0 is absorbing, for all $k \in \text{int } C$, there exists $t_0 > 0$ such that $t_0 k \in V_0$. Therefore, $y + t_0 k \in y + V_0 \subset -\text{int } C$. Hence, we have

$$\inf\{t : y \in tk - C\} \leq -t_0 < 0,$$

which shows that $h_C(y; k) < 0$.

Next we prove the statement (ii). Let $y \in Y$. Suppose that there exists $k \in \text{int } C$ such that $h_C(y; k) < 0$. Then, there exist $t_0 > 0$ and $c_0 \in C$ such that $y = -t_0 k - c_0 = -(t_0 k + c_0)$. Since $t_0 k \in \text{int } C$ and C is a convex cone, we have $y \in -\text{int } C$. ■

Remark 2.1 *By combining statements (i) and (ii) above, we have the following: there exists $k \in \text{int } C$ such that $h_C(y; k) < 0$ if and only if $y \in -\text{int } C$.*

Lemma 2.2 *Let $y \in Y$, then the following statements hold:*

- (i) *If $y \in -\text{cl } C$, then $h_C(y; k) \leq 0$ for all $k \in \text{int } C$;*
- (ii) *If there exists $k \in \text{int } C$ with $h_C(y; k) \leq 0$, then $y \in -\text{cl } C$.*

Proof. First we prove the statement (i). Suppose that $y \in -\text{cl } C$. Then, there exist a net $\{y_\lambda\} \subset -C$ such that y_λ converges to y . For each y_λ , since $y_\lambda \in 0 \cdot k - C$ for all $k \in \text{int } C$, $h_C(y_\lambda; k) \leq 0$ for all $k \in \text{int } C$. By the continuity of $h_C(\cdot; k)$, $h_C(y; k) \leq 0$ for all $k \in \text{int } C$.

Next we prove the statement (ii). Let $y \in Y$. Suppose that there exists $k \in \text{int } C$ such that $h_C(y; k) \leq 0$. In the case $h_C(y; k) < 0$, from (ii) of Lemma 2.1,

it is clear that $y \in -\text{cl } C$. So we assume that $h_C(y; k) = 0$ and show that $y \in -\text{cl } C$. By the definition of h_C , for each $n = 1, 2, \dots$, there exists $t_n \in R$ such that

$$h_C(y; k) \leq t_n < h_C(y; k) + \frac{1}{n} \quad (2.2)$$

and

$$y \in t_n k - C. \quad (2.3)$$

From (2.2), $\lim_{n \rightarrow \infty} t_n = 0$. From (2.3), there exists $c_n \in C$ such that $y = t_n k - c_n$, that is, $c_n = t_n k - y$. Since $c_n \rightarrow -y$ as $n \rightarrow \infty$, we have $y \in -\text{cl } C$. ■

Remark 2.2 By combining statements (i) and (ii) above, we have the following: there exists $k \in \text{int } C$ such that $h_C(y; k) \leq 0$ if and only if $y \in -\text{cl } C$.

Lemma 2.3 Let $y \in Y$, then the following statements hold:

- (i) If $y \in \text{int } C$, then $h_C(y; k) > 0$ for all $k \in \text{int } C$;
- (ii) If $y \in \text{cl } C$, then $h_C(y; k) \geq 0$ for all $k \in \text{int } C$.

The following lemma shows (strictly) monotone property on $h_C(\cdot; k)$, which has been investigated in [5] and [1].

Lemma 2.4 Let $y, \bar{y} \in Y$, then the following statements hold:

- (i) If $y \in \bar{y} + \text{int } C$, then $h_C(y; k) > h_C(\bar{y}; k)$ for all $k \in \text{int } C$;
- (ii) If $y \in \bar{y} + \text{cl } C$, then $h_C(y; k) \geq h_C(\bar{y}; k)$ for all $k \in \text{int } C$.

Lemma 2.5 Let $y, \bar{y} \in Y$ and $k \in \text{int } C$, then the following statements hold:

- (i) If $h_C(y; k) > h_C(\bar{y}; k)$, then $h_C(y - \bar{y}; k) > 0$;
- (ii) If $h_C(y; k) \geq h_C(\bar{y}; k)$, then $h_C(y - \bar{y}; k) \geq 0$.

Remark 2.3 In the above lemma, we note that each converse does not hold.

Now, we consider several characterizations for images of a set-valued map by the nonlinear and strictly monotone characteristic function h_C . We observe the following four types of scalarizing functions:

- (1) $\psi_C^F(x; k) := \sup \{h_C(y; k) : y \in F(x)\},$
- (2) $\varphi_C^F(x; k) := \inf \{h_C(y; k) : y \in F(x)\},$
- (3) $-\varphi_C^{-F}(x; k) = \sup \{-h_C(-y; k) : y \in F(x)\},$
- (4) $-\psi_C^{-F}(x; k) = \inf \{-h_C(-y; k) : y \in F(x)\}.$

Functions (1) and (4) have symmetric properties and then results for function (4) $-\psi_C^{-F}$ can be easily proved by those for function (1) ψ_C^F . Similarly, the results for function (3) $-\varphi_C^{-F}$ can be deduced by those for function (2) φ_C^F . By using these four functions we measure each image of set-valued map F with respect to its 4-tuple of scalars, which can be regarded as standpoints for the evaluation of the image with respect to convex cone C .

Proposition 2.1 *Let $x \in X$, then the following statements hold:*

- (i) *If $F(x) \cap (-\text{int } C) \neq \emptyset$, then $\varphi_C^F(x; k) < 0$ for all $k \in \text{int } C$;*
- (ii) *If there exists $k \in \text{int } C$ with $\varphi_C^F(x; k) < 0$, then $F(x) \cap (-\text{int } C) \neq \emptyset$.*

Proof. Let $x \in X$ be given. First we prove the statement (i). Suppose that $F(x) \cap (-\text{int } C) \neq \emptyset$. Then, there exists $y \in F(x) \cap (-\text{int } C)$. By (i) of Lemma 2.1, for all $k \in \text{int } C$, $h_C(y; k) < 0$, and hence, $\varphi_C^F(x; k) < 0$.

Next we prove the statement (ii). Suppose that there exists $k \in \text{int } C$ such that $\varphi_C^F(x; k) < 0$. Then, there exist $\varepsilon_0 > 0$ and $y_0 \in F(x)$ such that

$$h_C(y_0; k) \leq \inf_{y \in F(x)} h_C(y; k) + \varepsilon_0 < 0.$$

By (ii) of Lemma 2.1, we have $y_0 \in -\text{int } C$, which implies that $F(x) \cap (-\text{int } C) \neq \emptyset$. ■

Remark 2.4 *By combining statements (i) and (ii) above, we have the following: there exists $k \in \text{int } C$ such that $\varphi_C^F(x; k) < 0$ if and only if $F(x) \cap (-\text{int } C) \neq \emptyset$.*

Proposition 2.2 *Let $x \in X$, then the following statements hold:*

- (i) *If $F(x) \subset -\text{int } C$ and $F(x)$ is a compact set, then $\psi_C^F(x; k) < 0$ for all $k \in \text{int } C$;*
- (ii) *If there exists $k \in \text{int } C$ with $\psi_C^F(x; k) < 0$, then $F(x) \subset -\text{int } C$.*

Proof. Let $x \in X$ be given. First we prove the statement (i). Assume that $F(x)$ is a compact set and suppose that $F(x) \subset -\text{int } C$. Then, for all $k \in \text{int } C$,

$$F(x) \subset \bigcup_{t>0} (-tk - \text{int } C).$$

By the compactness of $F(x)$, there exist $t_1, \dots, t_m > 0$ such that

$$F(x) \subset \bigcup_{i=1}^m (-t_i k - \text{int } C).$$

Since $-t_q k - \text{int } C \subset -t_p k - \text{int } C$ for $t_p < t_q$, there exists $t_0 := \min\{t_1, \dots, t_m\} > 0$ such that $F(x) \subset -t_0 k - \text{int } C$. For each $y \in F(x)$, we have

$$h_C(y; k) = \inf\{t : y \in tk - C\} \leq -t_0.$$

Hence,

$$\psi_C^F(x; k) = \sup_{y \in F(x)} h_C(y; k) \leq -t_0 < 0.$$

Next, we prove the statement (ii). Suppose that there exists $k \in \text{int } C$ such that $\psi_C^F(x; k) < 0$. Then, for all $y \in F(x)$, $h_C(y; k) < 0$. By (ii) of Lemma 2.1, we have $y \in -\text{int } C$, and hence $F(x) \subset -\text{int } C$. ■

Remark 2.5 By combining statements (i) and (ii) above, we have the following: there exists $k \in \text{int } C$ such that $\psi_C^F(x; k) < 0$ if and only if $F(x) \subset -\text{int } C$. When we replace $F(x)$ in (i) of Proposition 2.2 by $\text{cl } F(x)$, the assertion still remains.

Moreover, we can replace (i) in Proposition 2.2 by another relaxed form.

Corollary 2.1 Let $x \in X$ and assume that there exists a compact set B such that $B \subset -\text{int } C$. If $F(x) \subset B - C$, then $\psi_C^F(x; k) < 0$ for all $k \in \text{int } C$.

Proof. Let $x \in X$, and assume that there exists a compact set B such that $B \subset -\text{int } C$ and $F(x) \subset B - C$. By applying (i) of Proposition 2.2 to B instead of $F(x)$, for all $k \in \text{int } C$,

$$\sup_{y \in B} h_C(y; k) < 0.$$

Since $F(x) \subset B - C$, it follows from (i) of Lemma 2.1 and the subadditivity of $h_C(\cdot; k)$ that

$$h_C(y; k) \leq \sup_{z \in B} h_C(z; k)$$

for each $y \in F(x)$. Therefore, $\psi_C^F(x; k) < 0$ for all $k \in \text{int } C$. ■

Proposition 2.3 Let $x \in X$, then the following statements hold:

- (i) If $F(x) \cap (-\text{cl } C) \neq \emptyset$, then $\varphi_C^F(x; k) \leq 0$ for all $k \in \text{int } C$;
- (ii) If $F(x)$ is a compact set and there exists $k \in \text{int } C$ with $\varphi_C^F(x; k) \leq 0$, then $F(x) \cap (-\text{cl } C) \neq \emptyset$.

Proof. Let $x \in X$ and we prove the statement (i). Suppose that $F(x) \cap (-\text{cl } C) \neq \emptyset$. Then, there exists $y \in F(x) \cap (-\text{cl } C)$. By (i) of Lemma 2.2, for all $k \in \text{int } C$, $h_C(y; k) \leq 0$, and hence $\varphi_C^F(x; k) \leq 0$.

Next, we prove the statement (ii). Suppose that there exists $k \in \text{int } C$ such that $\varphi_C^F(x; k) \leq 0$. In the case $\varphi_C^F(x; k) < 0$, from (ii) of Proposition 2.1, it is clear that $F(x) \cap (-\text{cl } C) \neq \emptyset$. So we assume that $\varphi_C^F(x; k) = 0$ and show that $F(x) \cap (-\text{cl } C) \neq \emptyset$. By the definition of φ_C^F , for each $n = 1, 2, \dots$, there exist $t_n \in R$ and $y_n \in F(x)$ such that $y_n \in t_n k - C$ and

$$\varphi_C^F(x; k) \leq t_n < \varphi_C^F(x; k) + \frac{1}{n}. \quad (2.4)$$

From (2.4), $\lim_{n \rightarrow \infty} t_n = 0$. Since $F(x)$ is compact, we may suppose that $y_n \rightarrow y_0$ for some $y_0 \in F(x)$ without loss of generality (taking subsequence). Therefore, $y_n - t_n k \rightarrow y_0$ and then $y_0 \in -\text{cl } C$, which shows that $F(x) \cap (-\text{cl } C) \neq \emptyset$. ■

Remark 2.6 By combining statements (i) and (ii) above, we have the following: under the compactness of $F(x)$, there exists $k \in \text{int } C$ such that $\varphi_C^F(x; k) \leq 0$ if and only if $F(x) \cap (-\text{cl } C) \neq \emptyset$. Otherwise, there are counter-examples violating the statement (ii) such as an unbounded set approaching $-\text{cl } C$ asymptotically or an open set whose boundary intersects $-\text{cl } C$.

Proposition 2.4 *Let $x \in X$, then the following statements hold:*

- (i) *If $F(x) \subset -\text{cl } C$, then $\psi_C^F(x; k) \leq 0$ for all $k \in \text{int } C$;*
- (ii) *If there exists $k \in \text{int } C$ with $\psi_C^F(x; k) \leq 0$, then $F(x) \subset -\text{cl } C$.*

Proof. Let $x \in X$ be given. First we prove the statement (i). Suppose that $F(x) \subset -\text{cl } C$. Then, for each $y \in F(x)$, it follows from (i) of Lemma 2.2 that $h_C(y; k) \leq 0$ for all $k \in \text{int } C$, and hence $\psi_C^F(x; k) \leq 0$ for all $k \in \text{int } C$.

Next, we prove the statement (ii). Suppose that there exists $k \in \text{int } C$ such that $\psi_C^F(x; k) \leq 0$. Then, for all $y \in F(x)$, $h_C(y; k) \leq 0$. By (ii) of Lemma 2.2, we have $y \in -\text{cl } C$, and hence $F(x) \subset -\text{cl } C$. ■

Remark 2.7 *By combining statements (i) and (ii) above, we have the following: there exists $k \in \text{int } C$ such that $\psi_C^F(x; k) \leq 0$ if and only if $F(x) \subset -\text{cl } C$.*

3 Alternative Theorems

In this section, we present various types of alternative theorems for set-valued maps without any convexity. These alternative theorems are fundamental tools to derive optimality conditions for vector optimization problems with set-valued maps. As stated in Introduction, there are five types of relationships between the zero vector and each image of a set-valued map with respect to a given dominance cone.

First, we present five types of alternative theorems for set-valued maps when we compare each image of set-valued map with the zero vector with respect to the interior of a convex cone.

Theorem 3.1 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. Then, exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \cap (-\text{int } C) \neq \emptyset$;*
- (II) *There exists $k \in \text{int } C$ such that $\varphi_C^F(x; k) \geq 0$ for all $x \in X$.*

Proof. First, we assume that system (I) holds. Then, there exists $x \in X$ such that $F(x) \cap (-\text{int } C) \neq \emptyset$. By (i) of Proposition 2.1, $\varphi_C^F(x; k) < 0$ for all $k \in \text{int } C$, which shows that system (II) does not hold.

Next, we assume that system (II) does not hold. Then, for all $k \in \text{int } C$, there exists $x \in X$ such that $\varphi_C^F(x; k) < 0$. By (ii) of Proposition 2.1, system (I) holds. ■

Theorem 3.2 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. If F is compact-valued on X , then exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \subset -\text{int } C$;*
 (II) *There exists $k \in \text{int } C$ such that $\psi_C^F(x; k) \geq 0$ for all $x \in X$.*

Proof. First, we assume that system (I) holds. Then, there exists $x \in X$ such that $F(x) \subset -\text{int } C$. By (i) of Proposition 2.2, $\psi_C^F(x; k) < 0$ for all $k \in \text{int } C$, which shows that system (II) does not hold.

Next, we assume that system (II) does not hold. Then, for all $k \in \text{int } C$, there exists $x \in X$ such that $\psi_C^F(x; k) < 0$. By (ii) of Proposition 2.2, system (I) holds. ■

Corollary 3.1 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. Assume that if $F(x) \subset -\text{int } C$, then there exists a compact subset $B \subset -\text{int } C$ such that $F(x) \subset B - C$. Then, exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \subset -\text{int } C$;*
 (II) *There exists $k \in \text{int } C$ such that $\psi_C^F(x; k) \geq 0$ for all $x \in X$.*

Proof. First, we assume that system (I) holds. Then, there exists $x \in X$ such that $F(x) \subset -\text{int } C$. By Corollary 2.1, $\psi_C^F(x; k) < 0$ for all $k \in \text{int } C$, which shows that system (II) does not hold.

Next, we assume that system (II) does not hold. Then, for all $k \in \text{int } C$, there exists $x \in X$ such that $\psi_C^F(x; k) < 0$. By (ii) of Proposition 2.2, system (I) holds. ■

Theorem 3.3 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. Then, exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \cap \text{int } C \neq \emptyset$;*
 (II) *There exists $k \in \text{int } C$ such that $-\varphi_C^{-F}(x; k) \leq 0$ for all $x \in X$.*

Proof. The proof is completed simply by replacing F by $-F$ in the proof of Theorem 3.1. ■

Theorem 3.4 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. If F is compact-valued on X , then exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \subset \text{int } C$;*

(II) *There exists $k \in \text{int } C$ such that $-\psi_C^{-F}(x; k) \leq 0$ for all $x \in X$.*

Proof. The proof is completed simply by replacing F by $-F$ in the proof of Theorem 3.2. ■

Corollary 3.2 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. Assume that if $F(x) \subset \text{int } C$, then there exists a compact subset $B \subset \text{int } C$ such that $F(x) \subset B + C$. Then, exactly one of the following two systems holds:*

(I) *There exists $x \in X$ such that $F(x) \subset \text{int } C$;*

(II) *There exists $k \in \text{int } C$ such that $-\psi_C^{-F}(x; k) \leq 0$ for all $x \in X$.*

Proof. The proof is completed simply by replacing F by $-F$ in the proof of Corollary 3.1. ■

Theorem 3.5 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. Then, exactly one of the following two systems holds:*

(I) *There exists $x \in X$ such that $F(x) \cap (-\text{int } C) \neq \emptyset$ or $F(x) \cap \text{int } C \neq \emptyset$;*

(II) *There exists $k \in \text{int } C$ such that $\varphi_C^F(x; k) \geq 0$ and $-\varphi_C^{-F}(x; k) \leq 0$ for all $x \in X$.*

Proof. The proof is straightforward from the same way as the proofs of Theorems 3.1 and 3.3. ■

Next, we present five types of alternative theorems for set-valued maps when we compare each image of set-valued map with the zero vector with respect to the closure of a convex cone.

Theorem 3.6 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. If F is compact-valued on X , then exactly one of the following two systems holds:*

(I) *There exists $x \in X$ such that $F(x) \cap (-\text{cl } C) \neq \emptyset$;*

(II) *There exists $k \in \text{int } C$ such that $\varphi_C^F(x; k) > 0$ for all $x \in X$.*

Proof. First, we assume that system (I) holds. Then, there exists $x \in X$ such that $F(x) \cap (-\text{cl } C) \neq \emptyset$. By (i) of Proposition 2.3, $\varphi_C^F(x; k) \leq 0$ for all $k \in \text{int } C$, which shows that system (II) does not hold.

Next, we assume that system (II) does not hold. Then, for all $k \in \text{int } C$, there exists $x \in X$ such that $\varphi_C^F(x; k) \leq 0$. By (ii) of Proposition 2.3, system (I) holds. ■

Theorem 3.7 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. Then, exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \subset -\text{cl } C$;*
- (II) *There exists $k \in \text{int } C$ such that $\psi_C^F(x; k) > 0$ for all $x \in X$.*

Proof. First, we assume that system (I) holds. Then, there exists $x \in X$ such that $F(x) \subset -\text{cl } C$. By (i) of Proposition 2.4, $\psi_C^F(x; k) \leq 0$ for all $k \in \text{int } C$, which shows that system (II) does not hold.

Next, we assume that system (II) does not hold. Then, for all $k \in \text{int } C$, there exists $x \in X$ such that $\psi_C^F(x; k) \leq 0$. By (ii) of Proposition 2.4, system (I) holds. ■

Theorem 3.8 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. If F is compact-valued on X , then exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \cap \text{cl } C \neq \emptyset$;*
- (II) *There exists $k \in \text{int } C$ such that $-\varphi_C^{-F}(x; k) < 0$ for all $x \in X$.*

Proof. The proof is completed simply by replacing F by $-F$ in the proof of Theorem 3.6. ■

Theorem 3.9 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. Then, exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \subset \text{cl } C$;*
- (II) *There exists $k \in \text{int } C$ such that $-\psi_C^{-F}(x; k) < 0$ for all $x \in X$.*

Proof. The proof is completed simply by replacing F by $-F$ in the proof of Theorem 3.7. ■

Theorem 3.10 *Let X and Y be a nonempty set and a topological vector space, C a convex cone in Y with its nonempty interior, and $F : X \rightarrow 2^Y$ a set-valued map, respectively. If F is compact-valued on X , then exactly one of the following two systems holds:*

- (I) *There exists $x \in X$ such that $F(x) \cap (-\text{cl } C) \neq \emptyset$ or $F(x) \cap \text{cl } C \neq \emptyset$;*
- (II) *There exists $k \in \text{int } C$ such that $\varphi_C^F(x; k) > 0$ and $-\varphi_C^{-F}(x; k) < 0$ for all $x \in X$.*

Proof. The proof is straightforward from the same way as the proofs of Theorems 3.6 and 3.8. ■

4 Optimality Conditions

Throughout this section, let X be a nonempty set, and let Y and Z be ordered topological vector spaces with convex cones C and D , respectively. We assume that $C \neq Y$ and $\text{int } C \neq \emptyset$. Let $F : X \rightarrow 2^Y$ and $G : X \rightarrow 2^Z$ be set-valued maps. A constrained set-valued optimization problem is written as

$$\begin{aligned} \text{(MP)} \quad & \min_K F(x) \\ & \text{subject to } G(x) \cap (-D) \neq \emptyset, \end{aligned}$$

where K is a convex cone in Y . The feasible set of problem (MP) is defined by $V = \{x \in X : G(x) \cap (-D) \neq \emptyset\}$. Problem (MP) is to find all solutions $x_0 \in V$ such that there exists $y_0 \in F(x_0)$ and for each $x \in V$, there exists no $y \in F(x)$ satisfying $y_0 \in y + K \setminus \{0_Y\}$. Such solution x_0 is called an efficient solution of problem (MP) with respect to K , and in case of $\text{int } K$ instead of K , its solution is called a weakly efficient solution (traditionally in vector optimization). Since the constraint in problem (MP) is reduced to $G(x) \leq 0$ when G is a real-valued function and D is the cone of nonnegative reals, it is a generalization of the inequality constraints of a standard nonlinear programming problem. Thus, we consider the following optimization problems:

$$\begin{aligned} \text{(MP1)} \quad & \min_{\text{int } C} F(x) \quad \text{subject to } G(x) \cap (-D) \neq \emptyset; \\ \text{(MP2)} \quad & \min_C F(x) \quad \text{subject to } G(x) \cap (-D) \neq \emptyset. \end{aligned}$$

Definition 4.1 A point $x_0 \in V$ is said to be a *weakly efficient solution* of (MP1) if there exists $y_0 \in F(x_0)$ and for each $x \in V$, there exists no $y \in F(x)$ satisfying $y_0 \in y + \text{int } C$, that is,

$$F(V) \cap (y_0 - \text{int } C) = \emptyset; \quad (4.1)$$

A pair (x_0, y_0) is said to be a *weakly efficient element* for (MP1) if $x_0 \in V$ and $y_0 \in F(x_0)$ satisfies (4.1).

Definition 4.2 A point $x_0 \in V$ is said to be an *efficient solution* of (MP2) if there exists $y_0 \in F(x_0)$ and for each $x \in V$, there exists no $y \in F(x)$ satisfying $y_0 \in y + C \setminus \{0_Y\}$, that is,

$$F(V) \cap (y_0 - C \setminus \{0_Y\}) = \emptyset; \quad (4.2)$$

A pair (x_0, y_0) is said to be an *efficient element* for (MP2) if $x_0 \in V$ and $y_0 \in F(x_0)$ satisfies (4.2).

Definition 4.3 Let $k \in \text{int } C$. Consider the following scalar minimization problem

$$\min_{x \in V} \varphi_C^F(x; k). \quad (4.3)$$

Let $x_0 \in V$ be given. Then, a pair (x_0, y_0) is said to be an *optimal element* for the problem if the following conditions hold:

- (i) $\varphi_C^F(x; k) \geq \varphi_C^F(x_0; k)$ for all $x \in V$;
- (ii) $\varphi_C^F(x_0; k) = h_C(y_0; k)$ and $y_0 \in F(x_0)$.

Remark 4.1 Under $k \in \text{int } C$, we have the following: a pair (x_0, y_0) is an optimal element for (4.3) if and only if $x_0 \in V$ and $y_0 \in F(x_0)$ satisfies

$$h_C(y; k) \geq h_C(y_0; k) \text{ for all } y \in F(V).$$

Definition 4.4 Let $k \in \text{int } C$. Consider problem (4.3). Let $x_0 \in V$ be given. Then, a pair (x_0, y_0) is said to be a *strict optimal element* if the following conditions hold:

- (i) $\varphi_C^F(x; k) > \varphi_C^F(x_0; k)$ for all $x \in V \setminus \{x_0\}$;
- (ii) $\varphi_C^F(x_0; k) = h_C(y_0; k)$ and $y_0 \in F(x_0)$;
- (iii) $h_C(y; k) > h_C(y_0; k)$ for all $y \in F(x_0) \setminus \{y_0\}$.

Remark 4.2 Under $k \in \text{int } C$, we have the following: a pair (x_0, y_0) is a *strict optimal element* for (4.3) if and only if $x_0 \in V$ and $y_0 \in F(x_0)$ satisfies

$$h_C(y; k) > h_C(y_0; k), \text{ for all } y \in F(V) \setminus \{y_0\}.$$

Theorem 4.1 (Sufficient condition for (MP1).) Let $\bar{x} \in V$ and $\bar{y} \in F(\bar{x})$. If there exists $k \in \text{int } C$ such that $(\bar{x}, \bar{y})$ is an optimal element for (4.3), then $(\bar{x}, \bar{y})$ is a weakly efficient element for (MP1).

Proof. Assume that $(\bar{x}, \bar{y})$ is not a weakly efficient element for (MP1). Then, there exist $x \in V$ and $y \in F(x)$ such that $\bar{y} \in y + \text{int } C$. Since $k \in \text{int } C$, it follows from (i) of Lemma 2.4 that $h_C(\bar{y}; k) > h_C(y; k)$. By Remark 4.1, it contradicts the assumption that $(\bar{x}, \bar{y})$ is an optimal element for (4.3). ■

Theorem 4.2 (Necessary and sufficient condition for (MP1).) Let $\bar{x} \in V$ and $\bar{y} \in F(\bar{x})$. Then $(\bar{x}, \bar{y})$ is a weakly efficient element for (MP1) if and only if there exists $k \in \text{int } C$ such that $h_C(y - \bar{y}; k) \geq 0$ for all $y \in F(V)$.

Proof. Suppose first that $(\bar{x}, \bar{y})$ is a weakly efficient element for (MP1). By definition, we have $(F(V) - \bar{y}) \cap (-\text{int } C) = \emptyset$. By applying Theorem 3.1 to $F(V) - \bar{y}$ instead of $F(x)$, there exists $k \in \text{int } C$ such that $h_C(y - \bar{y}; k) \geq 0$ for all $y \in F(V)$.

Conversely, suppose that there exists $k \in \text{int } C$ such that $h_C(y - \bar{y}; k) \geq 0$ for all $y \in F(V)$. Assume that $(\bar{x}, \bar{y})$ is not a weakly efficient element for (MP1). Then, there exist $x \in V$ and $y \in F(x)$ such that $y - \bar{y} \in -\text{int } C$. Since $k \in \text{int } C$, it follows from (i) of Lemma 2.1 that $h_C(y - \bar{y}; k) < 0$, which contradicts the assumption. ■

Theorem 4.3 (Sufficient condition for (MP2).) *Let $\bar{x} \in V$ and $\bar{y} \in F(\bar{x})$. If there exists $k \in \text{int } C$ such that $(\bar{x}, \bar{y})$ is a strict optimal element for (4.3), then $(\bar{x}, \bar{y})$ is an efficient element for (MP2).*

Proof. By applying the same argument as the proof of Theorem 4.1 to problem (MP2), the proof is straightforward from (ii) of Lemma 2.4 and Remark 4.2. ■

Theorem 4.4 (Necessary and sufficient condition for (MP2).) *Let $\bar{x} \in V$ and $\bar{y} \in F(\bar{x})$. If F is compact-valued on V and C is closed, then $(\bar{x}, \bar{y})$ is an efficient element for (MP2) if and only if there exists $k \in \text{int } C$ such that $h_C(y - \bar{y}; k) > 0$ for all $y \in F(V) \setminus \{\bar{y}\}$.*

Proof. For problem (MP2), by using the same argument as the proof of Theorem 4.2, it follows from Theorem 3.6 that the necessity is shown. By (i) of Lemma 2.2, we can also show the sufficiency. ■

5 Conclusions

Based on a nonlinear scalarization technique for sets, we establish five types of alternative theorems for set-valued maps without any convexity assumption. Moreover, we obtain optimality conditions for set-valued optimization problems.

References

- [1] G. Y. Chen and X. Q. Yang, Characterizations of Variable Domination Structures via Nonlinear Scalarization, *J. Optim. Theory Appl.* 112 (2002), 97–110.
- [2] O. Ferrero, Theorems of the Alternative for Set-Valued Functions in Infinite-Dimensional Spaces, *Optimization* 20 (1989), 167–175.
- [3] P. G. Georgiev and T. Tanaka, Vector-Valued Set-Valued Variants of Ky Fan's Inequality, *J. Nonlinear Convex Anal.* 1 (2000), 245–254.
- [4] P. Gr. Georgiev and T. Tanaka, Fan's Inequality for Set-Valued Maps, *Nonlinear Anal.* 47 (2001), 607–618.
- [5] C. Gerth and P. Weidner, Nonconvex Separation Theorems and Some Applications in Vector Optimization, *J. Optim. Theory Appl.* 67 (1990), 297–320.
- [6] F. Giannessi, Theorems of the Alternative for Multifunctions with Applications to Optimization: General Results, *J. Optim. Theory Appl.* 55 (1987), 233–256.
- [7] R. B. Holmes, *Geometric Functional Analysis and its Applications*, Springer-Verlag, New York, 1975.

- [8] V. Jeyakumar, A Generalization of a Minimax Theorem of Fan via a Theorem of the Alternative, *J. Optim. Theory Appl.* 48 (1986), 525–533.
- [9] D. Kuroiwa, T. Tanaka, and T.X.D. Ha, On cone convexity of set-valued maps, *Nonlinear Anal.* 30 (1997), 1487–1496.
- [10] Z. Li, A Theorem of the Alternative and Its Application to the Optimization of Set-Valued Maps, *J. Optim. Theory Appl.* 100 (1999), 365–375.
- [11] D. T. Luc, *Theory of Vector Optimization*, Lecture Note in Economics and Mathematical Systems, 319, Springer, Berlin, 1989.
- [12] S. Nishizawa and T. Tanaka, On Inherited Properties for Set-Valued Maps and Existence Theorems for Generalized Vector Equilibrium Problems, to appear in *J. Nonlinear Convex Anal.* (2004).
- [13] S. Nishizawa, T. Tanaka and P. Gr. Georgiev, On Inherited Properties of Set-Valued Maps, in *Nonlinear Analysis and Convex Analysis*, W. Takahashi and T. Tanaka (eds.), Yokohama Publishers, Yokohama, 2003, pp.341–350.
- [14] S. Nishizawa, T. Tanaka and P. Gr. Georgiev, On Inherited Properties for Vector-Valued Multifunctions, in *Multi-Objective Programming and Goal-Programming*, T. Tanino, T. Tanaka and M. Inuiguchi (eds.), Springer, Berlin, 2003, pp.215–220.
- [15] C. Tammer, A Generalization of Ekeland's Variational Principle, *Optimization* 25 (1992), 129–141.
- [16] C. Tammer, Stability Results for Approximately Efficient Solutions, *OR Spectrum* 16 (1994), 47–52.
- [17] X. M. Yang, X. Q. Yang and G. Y. Chen, Theorems of the Alternative and Optimization with Set-Valued Maps, *J. Optim. Theory Appl.* 107 (2000), 627–640.