

NOTES ON BI-AD_{ω_1}

DAISUKE IKEGAMI

ABSTRACT. In this note, we show that the axiom BI-AD_{ω_1} is inconsistent under $\text{ZF} + \text{AC}_{\omega}(\mathbb{R})$. This answers the question of Löwe in [3, Question 52].

1. INTRODUCTION AND BASIC DEFINITIONS

In this note, we will prove the following:

Theorem 1.1 ($\text{ZF} + \text{AC}_{\omega}(\mathbb{R})$). The axiom BI-AD_{ω_1} is inconsistent.

The axiom BI-AD_{ω_1} is a natural strengthening of the Axiom of Blackwell determinacy (BI-AD). For the background of this research topic, one can refer to the survey paper on Blackwell determinacy [3]. We mostly follow the standard notations and we use basic notions from the Jech's textbook on set theory [1].

First, let us define what BI-AD_{ω_1} is.

Let X be a set with more than one element and assume $\text{AC}_{\omega}({}^{\omega}X)$. By $\text{Prob}(X)$, we denote the set of all Borel probability measures on X with a countable support, i.e., the set of all Borel probability measures p such that there is a countable set $C \subseteq X$ with $p(C) = 1$.¹ From now on, we regard X as a discrete topological space and topologize ${}^{\omega}X$ as the product space. For any finite sequence s of elements in X , let $[s]$ be the basic open set generated by s , i.e., $[s] = \{x \in {}^{\omega}X; s \subseteq x\}$.

Let X^{Even} (X^{Odd}) be the set of finite sequences in X with even (odd) length. We call a function $\sigma: X^{\text{Even}} \rightarrow \text{Prob}(X)$ a *mixed strategy for player I* and a function $\tau: X^{\text{Odd}} \rightarrow \text{Prob}(X)$ a *mixed strategy for player II*. Given mixed strategies σ and τ for players I and II respectively, let $\nu(\sigma, \tau): {}^{<\omega}X \rightarrow \text{Prob}(X)$ as follows: For each finite sequence s of elements in X ,

$$\nu(\sigma, \tau)(s) = \begin{cases} \sigma(s) & \text{if } \text{lh}(s) \text{ is even,} \\ \tau(s) & \text{if } \text{lh}(s) \text{ is odd,} \end{cases}$$

where $\text{lh}(s)$ is the length of s . Since some of the calculations in this paper require a lot of parentheses, let us reduce their number by convention. If $(x_0, \dots, x_n)$ is a finite sequence, we

2010 *Mathematics Subject Classification*. Primary 03E60; Secondary 03E15, 91A05, 91A44.

Key words and phrases. Infinite games, Blackwell determinacy, descriptive set theory.

The author would like to thank Teruyuki Yorioka for organizing the RIMS workshop in set theory 2016. He is also grateful to JSPS for support through the grant with JSPS KAKENHI Grant Number 15K17586.

¹We are going to amalgamate a sequence of such measures to produce a product measure on ${}^{\omega}X$ as we construct the Lebesgue measure on ${}^{\omega}\omega$. For this purpose, the condition of having a countable support is essential.

write $[x_0, \dots, x_n]$ for the basic open set $[(x_0, \dots, x_n)]$. Similarly, if $x \in X$ and $\mu \in \text{Prob}(X)$, we write $\mu(x)$ for $\mu(\{x\})$. Now, for each finite sequence s of elements in X , define

$$\mu_{\sigma, \tau}([s]) = \prod_{i=0}^{\text{lh}(s)-1} \nu(\sigma, \tau)(s|_i)(s(i)).$$

By using $\text{AC}_\omega(\mathbb{R} \times {}^\omega X)$ (which follows from $\text{AC}_\omega({}^\omega X)$), we can uniquely extend $\mu_{\sigma, \tau}$ to a Borel probability measure on ${}^\omega X$, i.e., the probability measure whose domain is the set of all Borel sets in ${}^\omega X$. Let us also use $\mu_{\sigma, \tau}$ for denoting this Borel probability measure.

Let A be a subset of ${}^\omega X$. A mixed strategy σ for player I is *optimal in A* if for any mixed strategy τ for player II, A is $\mu_{\sigma, \tau}$ -measurable and $\mu_{\sigma, \tau}(A) = 1$. Similarly, a mixed strategy τ for player II is *optimal in A* if for any mixed strategy σ for player I, A is $\mu_{\sigma, \tau}$ -measurable and $\mu_{\sigma, \tau}(A) = 0$. We say that A is *Blackwell determined* if either player I or II has an optimal strategy in A . Finally, BI-AD_X is the statement “for any subset A of ${}^\omega X$, A is Blackwell determined.”²

Note that we only need $\text{AC}_\omega(\mathbb{R})$ to define BI-AD_{ω_1} because $\text{AC}_\omega({}^\omega \omega_1)$ follows from $\text{AC}_\omega(\mathbb{R})$.

2. PROOF OF THEOREM

Proof of Theorem 1.1. To derive a contradiction, let us assume BI-AD_{ω_1} .

We will prove the following two claims, which will be inconsistent to each other:

Claim 1. There is no injection from ω_1 to $\mathcal{P}(\omega)$.

Claim 2. There is an injection from ω_1 to $\mathcal{P}(\omega)$.

Proof of Claim 1. We will use the following fact:

Fact 2.1. Assume BI-AD . Then ω_1 is measurable.

Proof. See e.g., [2, Corollary 4.19]. □

Let μ be a normal measure on ω_1 and towards a contradiction, suppose there is an injection $f: \omega_1 \rightarrow \mathcal{P}(\omega)$.

We will derive a contradiction by the following arguments: for each natural number n , let $X_n = \{\alpha < \omega_1 \mid n \in f(\alpha)\}$. Since μ is a measure on ω_1 , for each n , either $X_n \in \mu$ or $\omega_1 \setminus X_n \in \mu$. Let $X = \bigcap \{X_n \mid X_n \in \mu\} \cap \bigcap \{\omega_1 \setminus X_n \mid X_n \notin \mu\}$. Then since μ is σ -complete, X is also in μ . Hence one can pick $\alpha, \beta \in X$ with $\alpha \neq \beta$.

We argue that $f(\alpha) = f(\beta)$, which would contract the assumption that f is injective. In fact, for each natural number n ,

$$n \in f(\alpha) \iff \alpha \in X_n \iff \beta \in X_n \iff n \in f(\beta),$$

²This formulation of Blackwell determinacy axioms does not involve imperfect information games; the original formulation due to Blackwell did, but these axioms turned out to be equivalent to the version we defined here which could be described as “perfect information determinacy with mixed strategies”. The imperfect information axiom would allow the players to move simultaneously, but at each move at least one of the players would have only finitely many choices. The proof of [4] adapts to show that the perfect information axiom implies the imperfect information one. Vervoort’s proof in [5] shows that optimal strategies exist for the perfect information games. For more details, cf. [3, §5].

where the second equivalence holds because both α and β are in X . Therefore, $f(\alpha) = f(\beta)$. $\square$

Proof of Claim 2. We will construct an injection $g: \omega_1 \rightarrow \mathcal{P}(\omega)$ using determinacy of Blackwell games with choosing countable ordinals.

Let us consider the following game $\mathcal{G}$: Player I chooses a countable ordinal α at first and then Player II chooses either 0 or 1 ω -many times and produces a real $y \subseteq \omega$. Player II wins if the real y codes α . Otherwise Player I wins.

This game can be formulated as a game in the definition of Bl-AD_{ω_1} and therefore, one of the players has an optimal strategy in the game $\mathcal{G}$.

Notice that Player I cannot have an optimal strategy in the game $\mathcal{G}$ because of the following argument: Suppose σ is an optimal strategy in the game $\mathcal{G}$ for Player I and then by the definition of a mixed strategy, there is an ordinal $\alpha < \omega_1$ such that $\sigma(\emptyset)(\{\alpha\}) > 0$. Then let τ_0 be the mixed strategy for Player II in the game $\mathcal{G}$ such that τ_0 let II play a real y coding α with measure 1. Then the probability of the payoff set in the $\mathcal{G}$ via μ_{σ, τ_0} is less than 1 and hence the mixed strategy σ is not optimal.

Hence Player II has an optimal strategy τ in the game $\mathcal{G}$ instead. We will produce an ω_1 -sequence $(x_\alpha \mid \alpha < \omega_1)$ of distinct reals using this strategy τ in the following way:

Given a countable ordinal α , let τ_α be the mixed strategy of Player II in the game $\mathcal{G}$ after Player I plays α with probability 1. Then τ_α can be regarded as a Borel probability on the Baire space and hence it can be seen as a real x_α .

We claim that the function $g(\alpha) = x_\alpha$ is injective. Let $\alpha < \beta$ be countable ordinals. Then the strategy τ_α concentrates on reals coding α with probability 1 while τ_β concentrates on reals coding β with probability 1. Therefore, τ_α and τ_β are distinct and so are x_α and x_β , as desired. $\square$

$\square$

REFERENCES

- [1] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.
- [2] Benedikt Löwe. The simulation technique and its applications to infinitary combinatorics under the axiom of Blackwell determinacy. *Pacific J. Math.*, 214(2):335–358, 2004.
- [3] Benedikt Löwe. Set theory of infinite imperfect information games. In Alessandro Andretta, editor, *Set theory: recent trends and applications*, volume 17 of *Quad. Mat.*, pages 137–181. Dept. Math., Seconda Univ. Napoli, Caserta, 2006.
- [4] Donald A. Martin. The determinacy of Blackwell games. *J. Symbolic Logic*, 63(4):1565–1581, 1998.
- [5] Donald A. Martin, Itay Neeman, and Marco Vervoort. The strength of Blackwell determinacy. *J. Symbolic Logic*, 68(2):615–636, 2003.

(D. Ikegami) DEPARTMENT OF MATHEMATICS, SCHOOL OF ENGINEERING, TOKYO DENKI UNIVERSITY,
5 ASAHI-CHO, SENJU, ADACHI-WARD, TOKYO 120-8551, JAPAN
E-mail address, D. Ikegami: ikegami@mail.dendai.ac.jp