

Optical Nano-science Research Section

K. Matsuda, Professor

K. Shinokita, Assistant Professor

1. Introduction

We are engaged in fundamental and applied research of nano-materials from a viewpoint of optics and material science. Our research aims to explore new physical and chemical phenomena leading to the applications of novel nano-materials including carbon nanotubes, layered transition metal dichalcogenides, perovskites for the efficient utilization of light energy and the development of future optoelectronic devices with ultra-low energy consumption. The followings are main the research achievements in the year of 2022.

1. Valley Relaxation of the Moiré Excitons in a WSe₂/MoSe₂ Heterobilayer

Due to their unusual physical properties and ability to function as extreme two-dimensional (2D) systems, atomically thin materials such as graphene and semiconducting transition-metal dichalcogenides (TMDs) and their van der Waals (vdW) heterostructures have gained increasing attention. The physical properties of atomically thin materials originating from a particular atomic arrangement are significantly altered by emerging moiré superlattices made up of lattice- or angular-mismatched vdW heterostructures. The moiré superlattice is gaining attention for its ability to engineer optically excited, tightly bound electron-hole pairs (excitons) and can result in exciton trapping through in-plane periodic moiré potentials in twisted semiconducting TMDs heterobilayers. The quantum confined zero-dimensional (0D) as a two-level system created by the moiré exciton trapped in the moiré potentials is encouraging for the development of quantum optics and quantum information processes.

The confinement of moiré potential of exciton alters the optical characteristics of the system. Under low-excitation power density conditions, the moiré exciton results in the appearance of sharp peaks in the low-temperature photoluminescence (PL) spectrum. Additionally, the moiré pattern (atomic registry) has a significant impact on the optical selection rule of the exciton. The intrinsic valley degrees of freedom control the optical properties of monolayer TMDs. The K and -K valleys at the edge of the Brillouin zone are inequivalent but energetically degenerate due to the strong spin-orbit interaction and lack of inversion symmetry. The optical selection rule and corresponding circular polarized light emission are both valley-dependent due to the inherent coupling of the valley




Fig. 1 (a) Schematic of moiré exciton in WSe₂/MoSe₂ heterobilayer. (b) Degree of circular polarization as a function of excitation power density. (c) Low-temperature circularly polarized PL spectrum under σ⁺ circularly polarized excitation (1.676 eV) with different excitation power densities. The σ⁺ and σ⁻ circularly polarized components are shown as the blue and red lines, respectively. (d) Excitation power density dependence of the PL intensity monitored at 1.321, 1.332, and 1.343 eV in the polarization-resolved PL spectra.

and spin degrees of freedom.

Contrary to monolayer TMDs, the optical selection rule of moiré superlattice also influences the circularly polarized light emission due to the atomic registry's C₃ symmetry and the valley degrees of freedom. The fascinating optical and electrical properties requires an understanding of the intra- and intervalley relaxation processes. The valley relaxation process in monolayer TMDs has thus been successfully explained and controlled through extensive experimental and theoretical studies. Electro-static carrier doping techniques were used to continuously control the valley relaxation process by altering the electron-hole exchange interaction. It has been discovered that momentum-dependent long-range electron-hole exchange interactions dominate the valley relaxation of neutral excitons in monolayer TMDs. On the other hand, it is still unclear how

the valley relaxation process in the moiré superlattice works, where the reduced dimensionality might also have an impact.

Here, we used circularly polarized PL and photoluminescence excitation (PLE) spectroscopy to investigate the intra- and intervalley relaxation of the moiré excitons in a twisted WSe₂/MoSe₂ hetero-bilayer. Contrary to 2D monolayer TMDs, the experimentally observed circularly polarized emission is strongly dependent on the excitation power density. The excitation power density dependence of circularly polarized emission leads to the inter-valley relaxation of the moiré exciton, which are originated from Pauli blocking due to the low density of states typical of 0D systems. The intravalley relaxation of the moiré exciton states from the triplet to singlet state via Γ_5 phonon emission is also revealed by the resonant PLE measurement. Circularly polarized quantum light emitter may be used in quantum optics and information processing as a result of our discoveries regarding the valley relaxation of moiré excitons.

2. Dynamics of Moiré Exciton in a Twisted MoSe₂/WSe₂ Heterobilayers

Studies in artificial vdW heterostructures made of two atomically thin 2D materials have received a lot of attention. A number of distinct physical phenomena, such as superconductivity, ferromagnetism near 3/4 filling, correlated insulator phases, and Hofstadter butterflies in graphene moiré superlattices, have new pathways thanks to the recent discovery of moiré superlattices at the interface of vdW hetero-structures with small lattice mismatches or twist angles between two monolayers. When optically generated electrons and holes in a spatially separated layer interact via Coulomb interactions, an interlayer exciton is created. This exciton is then contained within the moiré trap potential and spatially organized as a moiré exciton ensemble. The dense array of coherent quantum emitters is one application of the moiré trapped exciton that has great potential for many-body physics. However, little is known about the discrete excitonic states and dynamics of the confined exciton within the moiré potential.

The excitons can be divided into optically active bright excitons and inactive dark excitons in semiconductors. The dipole-allowed bright exciton exhibits radiative recombination during the emission process and directly couples to light via strong light-matter interaction. Optical absorption and PL spectroscopy make it possible to observe and track the dynamic behaviors of bright excitons. However, due to the weak light-matter interaction and reduced oscillator strength, optical access to the dark exciton is challenging. Dark excitons can be used in research on spin storage and long-lived qubits. The dark exciton state in monolayer 2D materials exhibits nonequilibrium dynamics as well as optical responses, and studies have shown that




Fig. 2 (a) Schematic of moiré exciton states confined in the potential. The splitting of bright and dark moiré exciton states is about 8 meV. (b) PL decay profiles at various temperatures. The contribution of bright and dark moiré exciton states is changed from low to high temperature.

it is energetically inferior to the bright exciton state.

The energy structures and characteristics of the bright and dark moiré exciton states in the moiré potential, however, may raise important and crucial questions. Therefore, it is essential to conduct experiments to demonstrate the existence of dark exciton within moiré potential. The dynamics of bright and dark moiré excitonic states in twisted semiconducting vdW heterobilayers requires further research in order to fully understand their revolutionary significance for fundamental moiré physics and potential quantum optics applications in the novel platform of moiré excitonic systems.

Here, we used PL spectroscopy and rate-equation analysis to investigate the moiré exciton states and their dynamics in twisted MoSe₂/WSe₂ heterobilayers. The experimental results demonstrate the existence of a dark moiré exciton state above the lowest emissive bright singlet exciton state, and the dynamics of bright excitons are determined by radical recombination and transition to the dark exciton state with the aid of phonons. In-depth information is provided regarding the dynamics and nature of moiré exciton states.

Our findings lay the groundwork for further investigation into quantum phenomena in moiré physics for use in quantum optics while illuminating novel aspects of quantum states of moiré excitons and fascinating moiré exciton dynamics.

Collaboration Works

松田一成, University of Bordeaux (フランス), 単一ナノ物質における先端分光

Financial Support

1. Grant-in-Aid for Scientific Research

松田一成, 基盤研究(S), 原子層人工ヘテロ構造におけるバレースピン量子光学の開拓と応用

松田一成, 学術変革領域研究(A), 2.5次元構造の分析技術開発

松田一成, 挑戦的研究(萌芽), バレースピン制御とデバイス応用: バレートロニクスに向けた課題と挑戦

松田一成, 学術変革領域研究(A), 2.5次元物質科学の総括(分担金)

篠北啓介, 基盤研究(B), モアレ超構造における協力的量子光学現象の開拓

田中絢也, 特別研究員奨励費, 室温動作超広帯域光検出器の実現に向けた量子物質赤外応答の解明

2. Others

篠北啓介, 創発的研究支援事業, 半導体モアレ超構造を用いた量子電磁力学の創生

Publications

M. Shimasaki, T. Nishihara, K. Matsuda, T. Endo, Y. Takaguchi, Z. Liu, Y. Miyata, Y. Miyauchi, Directional Exciton-Energy Transport in a Lateral Heteromonolayer of WSe_2 - MoSe_2 , ACS Nano, 16, 5, 8205-8212, 2022

H. Ago, S. Okada, Y. Miyata, K. Matsuda, M. Koshino, K. Ueno, K. Nagashio, Science of 2.5 Dimensional Materials: Paradigm Shift of Materials Science Toward Future Social Innovation, Science and Technology of Advanced Materials, 23, 1, 275-299, 2022

M. Li, J.Y. Wu, K. Sambe, Y. Yakiyama, T. Akutagawa, T. Kajitani, T. Fukushima, K. Matsuda, H. Sakurai, Dielectric Response of 1,1-difluorosumanene Caused by an In-plane Motion, Materials Chemistry Frontiers, 2022, 6, 1752-1758, 2022

N. Wada, J. Pu, Y. Takaguchi, W. Zhang, A. Liu, T. Endo, T. Irisawa, K. Matsuda, Y. Miyauchi, T. Takenobu, Y. Miyata, Efficient and Chiral Electroluminescence from

In-Plane Heterostructure of Transition Metal Dichalcogenide Monolayers, Advanced Functional Materials, 2203602, 2022

J. Pu, H. Ou, T. Yamada, N. Wada, H. Naito, H. Ogura, T. Endo, Z. Liu, T. Irisawa, K. Yanagi, Y. Nakanishi, Y. Gao, M. Maruyama, S. Okada, K. Shinokita, K. Matsuda, Y. Miyata, T. Takenobu, Continuous Color-Tunable Light-emitting Devices Based on Compositionally Graded Monolayer Transition Metal Dichalcogenide Alloys, Advanced Materials, 34, 2203250, 2022

K. Shinokita, K. Watanabe, T. Taniguchi, K. Matsuda, Valley Relaxation of the Moiré Excitons in a $\text{WSe}_2/\text{MoSe}_2$ Heterobilayer, ACS Nano, 16, 10, 16862-16868, 2022

K. Tanaka, T. Nishihara, A. Takakura, Y. Segawa, K. Matsuda, Y. Miyauchi, Polarized Raman spectroscopy on topological semimetal $\text{Co}_3\text{Sn}_2\text{S}_2$, Journal of Raman Spectroscopy, 2022

M. Shimasaki, T. Nishihara, N. Wada, Z. Liu, K. Matsuda, Y. Miyata, Y. Miyauchi, Anisotropic Exciton Drift-diffusion in a Monolayer $\text{WS}_{2-x}\text{Se}_{(2-2x)}$ Alloy with a Gradually Changing Composition, Applied Physics Express, 16, 1, 012010, 2023

Presentations

K. Matsuda, Novel optical science and its application in monolayer transition metal dichalcogenide and its heterostructure, Brookhaven National Laboratory Users Meeting, ハイブリッド開催, 2022.5.26

松田一成, 2次元層状物質および人工ヘテロ構造の光科学とその応用, 日本学術振興会第125光電変換委員会, オンライン開催, 2022.7.22

H. Kim, K. Aino, K. Shinokita, W. Zhang, K. Watanabe, T. Taniguchi, K. Matsuda, Observation of moiré exciton dynamics in twisted MoSe_2 - WSe_2 heterostrobilayer, Conference on the Lasers and Electro-Optics Pacific Rim 2022, Sapporo Convention Center, 2022.7.31-8.5

D. Dong, W. Zhang, K. Watanabe, T. Taniguchi, K. Shinokita, K. Matsuda, Gate tunable moiré excitonic states in twisted $\text{WSe}_2/\text{MoSe}_2$ heterobilayers, Conference on the Lasers and Electro-Optics Pacific Rim 2022, Sapporo Convention Center, 2022.7.31-8.5

S. Asada, K. Shinokita, K. Matsuda, Laser position and polarization angle dependence of spontaneous photocurrent in $\text{WSe}_2/\text{CrPS}_4$ hetero-interface., The 63th Fullerenes-Nanotube-Graphene General Symposium, Tokyo

Metropolitan University, 2022.8.31-9.2

H. Wang, K. Matsuda, Ab initio simulation study on linear and nonlinear optical properties of thin layer SnS, The 63th Fullerenes-Nanotube-Graphene General Symposium, Tokyo Metropolitan University, 2022.8.31-9.2

Y. Xiang, K. Matsuda, Simulation studies of nonlinear optical properties in two-dimensional Weyl semimetal WTe₂, The 63th Fullerenes-Nanotube-Graphene General Symposium, Tokyo Metropolitan University, 2022.8.31-9.2

篠北啓介, 相野薫平, 渡邊賢司, 谷口尚, 松田一成, WSe₂/MoSe₂ ヘテロ二層におけるモアレ励起子の円偏光発光励起スペクトル, 日本物理学会 2022 年秋季大会, ハイブリッド開催, 2022.9.12-15

H. Kim, K. Anio, K. Shinokita, W. Zhang, K. Watanabe, T. Taniguchi, K. Matsuda, Higher moiré excitonic states and its dynamics in twisted MoSe₂/WSe₂ heterostructure, JSAP-Optica Symposium, Tohoku University, 2022.9.20-9.23

K. Matsuda, Optical science of moiré excitonic systems in artificial van der Waals heterostructures, The 12th A3 Symposium on Emerging Materials: Nanomaterials for Electronics, Energy, and Environment, Waseda University, 2022.11.7-10

K. Matsuda, Optical science and application of moiré excitonic systems in two-dimensional hetero-structures, The 13th Recent Progress in Graphene and Two-dimensional Materials Research Conference, ハイブリッド開催, 2022.11.14-17

松田一成, 2.5次元物質における光科学と物性開拓, 第16回物性科学領域横断研究会, ハイブリッド開催, 2022.11.25-26

K. Matsuda, Optical science in two-dimensional material and its heterostructure, 11th International Workshop on 2D Materials, Hotel President, Seoul, Hybrid, 2023.2.17-18

D. Murase, K. Shinokita, K. Watanabe, T. Taniguchi, K. Matsuda, Fabrication of artificially designed two-dimensional material and van der Waals heterostructure by fully dry transfer method, The 64th Fullerenes-Nanotube-Graphene General Symposium, Nagoya University, 2023.3.1-3

H. Wang, H. Kim, K. Shinokita, K. Watanabe, T. Taniguchi, K. Matsuda, Observation of a single moiré exciton in nano-fabricated twisted MoSe₂/WSe₂ heterobilayers, The 64th Fullerenes-Nanotube-Graphene

General Symposium, Nagoya University, 2023.3.1-3

K. Aino, K. Shinokita, K. Matsuda, Bayesian optimization for the mechanical exfoliation of monolayer WSe₂, The 64th Fullerenes-Nanotube-Graphene General Symposium, Nagoya University, 2023.3.1-3

Y. Taniguchi, M. Hasegawa, T. Kitao, T. Uemura, K. Suzuki, H. Kaji, K. Matsuda, Optical properties of armchair graphene nanoribbon synthesized by metal-organic framework, The 64th Fullerenes-Nanotube-Graphene General Symposium, Nagoya University, 2023.3.1-3

松田一成, ナノサイエンスに立脚した光科学: 未来へ向けて, 1次元、2次元物質科学の展望と課題, 東北大学, 2023.3.10