

ORIGINAL ARTICLE

Protective effects of Alda-1, an ALDH2 activator, on alcohol-derived DNA damage in the esophagus of human ALDH2*2 (Glu504Lys) knock-in mice

Kenshiro Hirohashi^{1,2,*}, Shinya Ohashi², Yusuke Amanuma², Yukie Nakai², Tomomi Ida², Kiichiro Baba², Yosuke Mitani², Ayaka Mizumoto², Yoshihiro Yamamoto², Osamu Kikuchi³, Junichi Matsubara², Atsushi Yamada², Shin'ichi Miyamoto¹, Hiroshi Seno¹, Tomonari Matsuda⁴ and Manabu Muto^{2,*}

¹Department of Gastroenterology and Hepatology and ²Department of Therapeutic Oncology, Graduate School of Medicine, Kyoto University, 54 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan, ³Department of Medical Oncology, Dana-Farber Cancer Institute, 450 Brookline Avenue, Boston, MA 02215, USA and ⁴Research Center for Environmental Quality Management, Kyoto University, 1-2 Yumihama, Otsu 520-0811, Japan

*To whom correspondence should be addressed. Tel: +81 75 751 4592; Fax: +81 75 751 4594; Email: mmuto@kuhp.kyoto-u.ac.jp

Abstract

Alcohol consumption is the key risk factor for the development of esophageal squamous cell carcinoma (ESCC), and acetaldehyde, a metabolite of alcohol, is an alcohol-derived major carcinogen that causes DNA damage. Aldehyde dehydrogenase2 (ALDH2) is an enzyme that detoxifies acetaldehyde, and its activity is reduced by ALDH2 gene polymorphism. Reduction in ALDH2 activity increases blood, salivary and breath acetaldehyde levels after alcohol intake, and it is deeply associated with the development of ESCC. Heavy alcohol consumption in individuals with ALDH2 gene polymorphism significantly elevates the risk of ESCC; however, effective prevention has not been established yet. In this study, we investigated the protective effects of Alda-1, a small molecule ALDH2 activator, on alcohol-mediated esophageal DNA damage. Here, we generated novel genetically engineered knock-in mice that express the human ALDH2*1 (wild-type allele) or ALDH2*2 gene (mutant allele). Those mice were crossed, and human ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 knock-in mice were established. They were given 10% ethanol for 7 days in the presence or absence of Alda-1, and we measured the levels of esophageal DNA damage, represented by DNA adduct (N²-ethylidene-2'-deoxyguanosine). Alda-1 significantly increased hepatic ALDH2 activity both in human ALDH2*1/*2 and/or ALDH2*2/*2 knock-in mice and reduced esophageal DNA damage levels after alcohol drinking. Conversely, cyanamide, an ALDH2-inhibitor, significantly exacerbated esophageal DNA adduct level in C57BL/6N mice induced by alcohol drinking. These results indicate the protective effects of ALDH2 activation by Alda-1 on esophageal DNA damage levels in individuals with ALDH2 gene polymorphism, providing a new insight into acetaldehyde-mediated esophageal carcinogenesis and prevention.

Introduction

Esophageal squamous cell carcinoma (ESCC) is the disease with high morbidity and mortality (1,2). Alcohol consumption is the major risk factor for ESCC, and acetaldehyde, a metabolite of ethanol, is considered to play a central role in alcohol-related

esophageal carcinogenesis (3–5). Acetaldehyde causes various DNA damages, such as DNA adducts including N²-ethylidene-2'-deoxyguanosine (N²-ethylidene-dG) (6) and N²-ethyl-2'-deoxyguanosine (N²-Et-dG) (7), single- and/or double-strand

Received: December 2, 2018; Revised: April 15, 2019; Accepted: May 9, 2019

© The Author(s) 2019. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

For commercial re-use, please contact journals.permissions@oup.com

Abbreviation

ALDH2	aldehyde dehydrogenase 2
cDNA	complementary DNA
ESCC	esophageal squamous cell carcinoma
N ² -Et-dG	N ² -ethyl-2'-deoxyguanosine
UTR	untranslated region

breaks, point mutations, sister chromatid exchanges and DNA-DNA cross-links (8–13). 'Acetaldehyde associated with alcohol consumption' is designated as a definite carcinogen for the esophagus (14).

Acetaldehyde is detoxified to acetic acid by aldehyde dehydrogenase 2 (ALDH2) mainly in the liver (15). ALDH2 gene has two major alleles, ALDH2*1 (active ALDH2) and ALDH2*2 (inactive ALDH2) (16). ALDH2*2 (rs671) allele, also known as Glu504Lys, encodes the ALDH2 protein, which is defective at metabolizing acetaldehyde (17). Combination of ALDH2*1 and ALDH2*2 alleles makes up three genotypes with various enzymatic activity; wild-type (ALDH2*1/*1), heterozygous (ALDH2*1/*2) and homozygous (ALDH2*2/*2) (18,19). As ALDH2*2 acts in a dominant negative manner (20), hepatic activity of heterozygous (ALDH2*1/*2) and homozygous (ALDH2*2/*2) mutant ALDH2 is less than 10% and 0%, respectively, compared with that of wild-type (ALDH2*1/*1) ALDH2 (21–23). ALDH2*2/*2 carriers do not tolerate alcohol and few of them habitually drink alcoholic beverages, meanwhile, ALDH2*1/*2 carriers may drink various amount of alcohol (24,25). Previous epidemiological analysis showed that ALDH2*1/*2 carriers with heavy alcohol consumption are at high risk for ESCC (26–28). According to a meta-analysis, ALDH2*1/*2 carriers have a 7.12-fold increased risk of ESCC, compared with ALDH2*1/*1 carriers (29). Moreover, alcoholics with the ALDH2*1/*2 carriers have a 13.5-fold increased risk of ESCC, compared with ALDH2*1/*1 carriers (30). Thus, the risk of ESCC in ALDH2*1/*2 carriers rises according to the amount of alcohol consumption (31). Of note, about 70% of ESCC patients in East Asian countries such as Japan or Taiwan are revealed to have ALDH2*2 allele carriers (32,33).

Alcohol intake increases acetaldehyde concentrations in the blood, saliva and the breath, and ALDH2 gene polymorphisms affect those levels (34–36). Importantly, salivary acetaldehyde concentration reaches a high level in individuals with ALDH2*1/*2 after alcohol consumption (37). Consequently, high acetaldehyde-containing saliva can be directly exposed to the pharynx and esophagus and may induce various acetaldehyde-mediated DNA damages in these individuals. Indeed, esophageal DNA damage levels in ALDH2 knockout mice were much higher than those in control ALDH2 wild-type mice after drinking 10% ethanol for 2 months (38,39).

Thus, impaired ALDH2 activity due to ALDH2 gene polymorphism is considered to be deeply involved in esophageal carcinogenesis. Therefore, restoration of ALDH2 activity might be beneficial to ALDH2*1/*2 carriers in ESCC prevention. At present, a small molecule Alda-1 has been identified as an ALDH2 activator (40,41). Here, we hypothesized that Alda-1 may reduce esophageal DNA injury in the ALDH2*2 allele carriers with alcohol consumption. To verify that hypothesis, we generated novel genetically engineered knock-in mice that express the human ALDH2*1 and/or ALDH2*2 and investigated the effects of Alda-1 on these knock-in mice.

The aim of this study was to clarify whether Alda-1 has protective effects against alcohol-derived DNA damage in the esophagus of ALDH2*2 allele carriers.

Materials and methods

Mouse preparation

Age-matched control C57BL/6N male mice, which carry the wild-type *Aldh2* gene, were purchased from Charles River Laboratories Japan (Yokohama, Japan). All experiments conformed to the relevant regulatory standards and were approved by the institutional animal care and use committee of Kyoto University (Med Kyo 16196). To obtain tissues from mice, they were euthanized painlessly under anesthesia with diethyl ether inhalation followed by cervical dislocation. For the measurement of DNA adduct levels, esophageal tissues were frozen in liquid nitrogen and stored at -80°C until use. Liver tissues were collected in liquid nitrogen and stored at -80°C for analysis of ALDH activity.

Generation of genetically engineered knock-in mice that express the human ALDH2*1 (wild-type allele) or ALDH2*2 gene (mutant allele)

Human ALDH2*1 or ALDH2*2 knock-in mice in C57BL/6N background were generated by homologous recombination in TransGenic (Fukuoka, Japan) in accordance with the institutional guidelines. We designed targeting vector to replace mouse *Aldh2* gene with human ALDH2*1 or ALDH2*2 complementary DNA (cDNA) (Figure 1A). The design of the vector construction is summarized in Figure 1B. As 5' homologous arms, we used a 2.6k base pair (bp) mouse genomic fragment containing 5' untranslated region (UTR) of exon 1 of *Aldh2* gene. As 3' homologous arms, we used a 5.1k bp fragment containing exon 3–7 of *Aldh2* gene. These genomic fragments were amplified by PCR from RENKA ES cell genomic DNA (42).

Mouse *Aldh2* 5' UTR sequence was amplified by PCR from RENKA ES cell genomic DNA and cloned into a vector, named pUC118 generated by TransGenic, followed by human ALDH2 (ALDH2*1 or ALDH2*2) cDNA fragment (1554 bp) and bovine growth hormone polyA signal (5' UTR + cDNA + polyA).

3' homologous arm was amplified by PCR and cloned into a vector, named pPND2.1 generated by TransGenic, which contains lox-P flanked PGK_neo cassette (phosphoglycerate kinase 1 promoter-driven neomycin resistant gene) as a positive selection marker and MC1.DTA cassette (polyoma enhancer/herpes simplex virus thymidine kinase promoter driven diphtheria toxin A gene) as a negative selection marker.

Then, PCR-amplified 5' homologous arm and 5' UTR + cDNA + polyA were subcloned into the plasmid with 3' homologous arm, PGK_neo cassette and MC1.DTA cassette. Resulting targeting vector contains MC1.DTA cassette, 5' homologous arm, human ALDH2 (ALDH2*1 or ALDH2*2) cDNA fragment, polyA, lox-P flanked PGK_neo cassette and 5.1 kb 3' homologous arm. Thus, ALDH2*1 or ALDH2*2 targeting vector was generated. To amplify the fragment from genomic DNA, primer sets as shown in Table 1 were used.

This targeting vector was linearized and introduced into RENKA ES cells (C57BL/6N) by electroporation. After selection using neomycin, the resistant clones were isolated, and their DNAs were screened for homologous recombination by PCR using following primer set: sc_5AF2: 5'-ACC ATC CAT TCA AGG TAA AGT TCC -3' and neo_108r: 5'-CCT CAG AAG AAC TCG TCA AGA AG-3'. PCR positive ES clones were expanded, and isolated DNAs were further analyzed by PCR amplification using following primer sets: sc_5AF2 and neo_108r for 5' amplification, sc_3AR3: 5'-CAG GCA CAG GTT ACT ACT CTT CC-3' and neo_MS: 5'-ATT CGC AGC GCA TCG CCT TCT ATC GCC TTC -3' for 3' amplification. Homologous recombination of these clones was also confirmed by genomic Southern hybridization probed with neomycin-resistant gene.

Homologous recombinant ES cell clones were aggregated with ICR 8 cell embryos to generate chimeric mice. Germline transmitted heterozygous ALDH2 knock-in mice (F1 heterozygous ALDH2*1 or ALDH2*2 knock-in mice) were obtained by crossing chimeric mice with a high contribution of the RENKA background with C57BL/6N mice. Targeted allele was identified by PCR with the following primer sets: sc_5AF2 and neo_108r.

F1 heterozygous ALDH2*1 or ALDH2*2 knock-in mice were crossed each other to obtain F2 homozygous ALDH2*1 or ALDH2*2 knock-in mice. The F2 homozygous mice were crossed with B6.Cg-Tg (CAG-Cre) CZ-MO2Osb




Figure 1. Scheme of human *ALDH2**1 or *ALDH2**2 targeting vector to generate genetically engineered knock-in mice that express the human *ALDH2**1 or *ALDH2**2 gene. (A) Human *ALDH2* (*ALDH2**1 or *ALDH2*) cDNA knock-in strategy. bp: base pair. (B) Construction for human *ALDH2**1 or *ALDH2**2 knock-in targeting vector. 5' homologous arm (2.6k bp), 5' UTR of exon 1 of *Aldh2* gene + human *ALDH2* (*ALDH2**1 or *ALDH2*) cDNA (1554 bp) + polyA containing vector, and 3' homologous arm (5.1k bp) are included in human *ALDH2**1 or *ALDH2**2 knock-in targeting vector.

mice (CAG-Cre mice), which was provided by the RIKEN BRC (Tsukuba, Japan), to eliminate neomycin-resistant gene. Consequently, human *ALDH2**1/*1 or *ALDH2**2/*2 knock-in mice were generated. And then, the human *ALDH2**1/*1 and *ALDH2**2/*2 knock-in mice were crossed to generate human *ALDH2**1/*2 knock-in mice.

The genotype of *Aldh2* mice and human *ALDH2* mice was confirmed by PCR amplification using DNA extracted from mouse tails. The primers used to identify mouse *Aldh2* were as follows: forward, 5'-GAGGACTGTGTTGGGAGGTC-3'; reverse, 5'-GTAGTCCGGTCCCGTTC-3' (264 bp fragment). The PCR conditions used were 94°C for 3 min, followed by 30 cycles of 94°C for 30 s, 62°C for 30 s and 72°C for 30 s. A final

extension was conducted at 72°C for 3 min. The primers used for the detection of human *ALDH2* were as follows: forward, 5'-AGATGTGCAGGATG GCATGACC-3'; reverse, 5'-ACCCGGTAGAATTCGACGACC-3' (790 bp fragment). The PCR conditions used were 94°C for 3 min, followed by 30 cycles of 98°C for 10 s, 62°C for 30 s and 68°C for 1 min. A final extension was conducted at 68°C for 3 min. To confirm the elimination of the neomycin-resistance gene, we used the following primers: forward, 5'-AGATGTGC AGGATGGCATGACC-3'; reverse, 5'-GGATGCTATGAGCTTCATTC-3' (748 bp fragment). The PCR conditions used were 94°C for 3 min, followed by 30 cycles of 98°C for 10 s, 62°C for 30 s and 68°C for 2 min. A final extension was conducted at 68°C for 3 min.

Table 1. Primer sets for amplification from genomic DNA

	Primer name	Sequence(5'-3')	Size (mer)
PCR for 5' homologous arm			
Forward	5AF1	CAC TAT CTT CTT CTT CCC TGT GC	23
Reverse	5AR1	GAA GTG AGT CTC ACC TGG TTG C	22
PCR for 3' homologous arm			
Forward	3AF1	TGA ATG AAG CTC ATA GCA TCC	21
Reverse	3AR1	GGT CCT TTG TAG GAG GTT ACA GC	23
5' PCR for screening			
Forward	sc_5AF2	CC ATC CAT TCA AGG TAA AGT TCC	23
Reverse	neo_108r	CCT CAG AAG AAC TCG TCA AGA AG	23
3' PCR for screening			
Forward	neo_MS	ATT CGC AGC GCA TCG CCT TCT ATC GCC TTC	30
Reverse	sc_3AR3	CAG GCA CAG GTT ACT ACT CTT CC	23

In addition, the genotype of human ALDH2*1 and/or ALDH2*2 was confirmed by the PCR-restriction fragment length polymorphism (RFLP) methods (43). We used the specific restriction enzyme, *AcuI* (R0641S, New England Biolabs Tokyo, Japan), to distinguish between human ALDH2*1 and ALDH2*2. The PCR conditions for *AcuI* digestion were 37°C for 90 min, followed by 65°C for 20 min.

Measurement of enzymatic hepatic ALDH2 activity in mice

Hepatic ALDH2 activity in mice was determined as described previously (40). Briefly, it was determined spectrophotometrically by using extracted protein from tissue homogenate by monitoring the reductive reaction of NAD⁺ to NADH at λ 340 nm. All the assays were carried out at 25°C in 0.1 M sodium pyrophosphate buffer, pH = 9.5, 2.4 mM NAD⁺ and 10 mM acetaldehyde as the substrate.

Measurement of esophageal N²-ethylidene-dG levels in mice

Esophageal N²-ethylidene-dG level was quantified as described previously (44). Briefly, DNA was isolated from esophageal tissue specimens. NaBH₃CN (100 mM; 156159, Sigma-Aldrich, St. Louis, MO), a reducing reagent, was added to DNA samples. This converts N²-ethylidene-dG to stable N²-Et-dG. As the endogenous N²-Et-dG level in tissues is extremely low, the N²-Et-dG level that is converted from N²-ethylidene-dG indicates the endogenous N²-ethylidene-dG level (45). The DNA adduct standard, N²-Et-dG and its stable isotope, [U-15N5]-labeled N²-Et-dG, were synthesized as described previously (46). DNA samples were digested as described previously (46) and subjected to liquid chromatography-tandem mass spectrometry (LC/MS/MS). LC/MS/MS analyses were performed using a Shimadzu LC system (Shimadzu Corp., Kyoto, Japan) interfaced with a Quattro Ultimo triple-stage quadrupole mass spectrometer or an ACQUITY UPLC H-Class system interfaced with a XEVO-TSQ triple-stage quadrupole mass spectrometer (Waters Corp., Milford, MA), as reported previously (46). Shim-pack XR-ODS columns (3.0 × 75 mm, 2.2 μ m; Shimadzu Corp.) or ACQUITY UPLC BEH C18 columns (2.1 × 100 mm, 1.7 μ m; Waters Corp.) were used to separate the samples.

Single injection of Alda-1, an ALDH2 activator, on ALDH2 knock-in mice

To examine the effects of Alda-1 on ALDH2 activation, we injected Alda-1 (20 mg/kg body weight; SML-0462, Sigma-Aldrich) or vehicle control (dimethyl sulfoxide, 472301, Sigma-Aldrich) intraperitoneally on ALDH2*1/*1, ALDH2*1/*2 and/or ALDH2*2/*2 knock-in mice. Liver of those mice was collected 3 h after the injection and used for measuring the hepatic ALDH activity.

Alda-1 treatment in ALDH2 knock-in mice with alcohol drinking

ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 knock-in mice were given 10% ethanol instead of water and allowed to drink freely and injected

intraperitoneally with either Alda-1 (20 mg/kg body weight twice a day, n = 8) or vehicle control (dimethyl sulfoxide, n = 8) every 12 h for 7 days. And the esophageal tissues were collected for measuring N²-ethylidene-dG levels.

Single injection of cyanamide, an ALDH2-inhibitor, on C57BL/6N mice

Cyanamide (187364, Sigma-Aldrich), a well-established ALDH2-inhibitor (47), was dissolved in distilled water. We injected cyanamide (1.5 mg/kg body weight) or distilled water intraperitoneally in male C57BL/6N mice, and then hepatic ALDH activity was measured at 2 h after the injection.

Cyanamide treatment on alcohol drinking in C57BL6 mice

Male C57BL/6N mice were allowed to drink 10% ethanol freely and were injected intraperitoneally with either cyanamide (1.5 mg/kg body weight, twice a day, n = 6) or vehicle control (distilled water, n = 6) every 12 h for 7 days, and the esophageal tissues were collected to measure N²-ethylidene-dG level.

TaqMan gene expression assays

RNA was isolated using RNeasy Plus Mini Kits (QIAGEN, Hilden, Germany) and cDNA was synthesized using PrimeScript RT reagent kits (Takara Bio, Kusatsu, Japan) according to the manufacturer's instructions. Real-time reverse-transcription PCR was conducted with TaqMan Gene Expression Assays (Life Technologies Corp., Tokyo, Japan) for ALDH2 (Assay ID; Hs01007998_m1) and for β -actin (Assay ID; Hs99999903_m1) using a LightCycler 480 Instrument II (Hoffmann-La Roche Ltd., Basel, Switzerland) as described previously (48). All PCRs were performed in triplicate. The relative ALDH2 messenger RNA expression level was normalized to that of β -actin as an internal control.

Western blot analysis

Hepatic whole-cell lysates were prepared as described previously (49). The denatured protein samples (20 μ g) were fractionated on Any kD Mini-PROTEAN TGX Precast Gels (Bio-Rad Laboratories, Hercules, CA). Primary antibodies and the titers used for western blotting were as follows: goat polyclonal anti-ALDH2 (N-14) (sc-48838; Santa Cruz Biotechnology, Santa Cruz, CA; 1:1000); rabbit monoclonal anti- β -actin (13E5; Cell Signaling Technology; 1:2000). These were then reacted with the appropriate horseradish peroxidase-conjugated secondary antibody (GE Healthcare, Little Chalfont, UK). The signal was visualized using an Immobilon Western Chemiluminescent Horseradish Peroxidase Substrate (Merck Millipore, Darmstadt, Germany) and was exposed using a ChemiDoc XRS system equipped with Image Lab software (Bio-Rad Laboratories).

Statistical analysis

Data are presented as the mean $\pm$ standard deviation. The data were first tested for normality of distribution. The differences between two groups were analyzed using two-tailed paired Student's *t*-tests for equal variance data. *P*-values of less than 0.05 were considered significant.

Results

Generation of human *ALDH2**1/*1, *ALDH2**1/*2, or *ALDH2**2/*2 knock-in mice

To confirm the generation of human *ALDH2* (*ALDH2**1 or *ALDH2**2) gene knock-in mice, we first checked the deletion of mice *Aldh2* gene and insertion of human *ALDH2* gene by PCR using the primers specifically recognizing mice *Aldh2* or human *ALDH2*. As shown in Figure 2A, control C57BL/6N mice (mice/mice) were positive for mice *Aldh2* and negative for human *ALDH2*. F1 heterozygous human *ALDH2* knock-in mice (mice *Aldh2*/human *ALDH2**1 or *ALDH2**2) showed positive for both mice *Aldh2* and human *ALDH2*. F2 homozygous human *ALDH2* knock-in mice (human *ALDH2**1/*1 or *ALDH2**2/*2) showed negative for mice *Aldh2* and positive for human *ALDH2*.

Next, we eliminated neomycin-resistant gene in F2 homozygous human *ALDH2**1/*1 or *ALDH2**2/*2 knock-in mice by crossed with CAG-Cre mice, and its absence was confirmed by PCR. When the mice possess the neomycin-resistant gene, 2498 bp bands are seen. On the other hand, 748 bp bands are observed when neomycin-resistant gene is deleted (Figure 2B).

Next, in F2 homozygous human *ALDH2* knock-in mice, we checked the genotype (i.e. either *ALDH2**1/*1, *ALDH2**1/*2 or *ALDH2**2/*2) by PCR-RFLP method. After digestion by restriction enzyme *AclI*, PCR products derived from human *ALDH2**1/*1 knock-in mice were recognized by showing three bands at 430 bp, 235 bp and 83 bp, whereas those from human *ALDH2**2/*2

knock-in mice presented with two bands at 665 bp and 83 bp. After cleavage by *AclI*, PCR products derived from human *ALDH2**1/*2 knock-in mice showed four bands at 665 bp, 430 bp, 235 bp and 83 bp, and thus, human *ALDH2**1/*1, *ALDH2**1/*2 and *ALDH2**2/*2 knock-in mice were generated, and hereafter, those mice are described as *ALDH2**1/*1, *ALDH2**1/*2 and *ALDH2**2/*2 mice, respectively.

Characterization of *ALDH2**1/*1, *ALDH2**1/*2 and *ALDH2**2/*2 mice

We then characterized *ALDH2**1/*1, *ALDH2**1/*2 and *ALDH2**2/*2 mice, in terms of expression and enzymatic activity of *ALDH2*. When measured using liver tissue homogenate, the *ALDH2* messenger RNA (Figure 3A) and *ALDH2* protein (Figure 3B) expression levels did not differ among three groups. In contrast, *ALDH* activity in *ALDH2**1/*2 and *ALDH2**2/*2 mice was significantly lower than that in *ALDH2**1/*1 mice (Figure 3C).

Next, we investigated the effect of *ALDH2* genotype in mice on alcohol consumption as well as alcohol-derived DNA damage in esophageal tissues. When mice were given 10% ethanol as substitute for water for 7 days, the average amount of ethanol consumption in *ALDH2**1/*2 or *ALDH2**2/*2 mice was significantly less than that in *ALDH2**1/*1 mice (Figure 4A). Alcohol drinking increased esophageal DNA damage represented by N²-ethylidene-dG levels in all groups. Of note, *ALDH2**1/*2 and *ALDH2**2/*2 mice showed significantly higher DNA damage than *ALDH2**1/*1 mice (Figure 4B).




Figure 2. Genotyping of human *ALDH2**1/*1, *ALDH2**1/*2 and *ALDH2**2/*2 knock-in mice. (A) PCR images for mice *Aldh2* (left panel), and human *ALDH2* (right panel). Left lane in left and right panel: molecular weight (MW) marker, 100 base pair (bp) DNA ladder. PCR products of mice *Aldh2* and human *ALDH2* were 264 bp and 790 bp, respectively. (B) Elimination of neomycin resistant gene. Left lane: MW marker (100 bp DNA ladder). PCR products of human *ALDH2* with or without neomycin-resistant gene were 2498 bp and 748 bp, respectively. Neo (+): before elimination of neomycin resistant gene, Neo (-): after elimination of neomycin resistant gene. (C) The PCR images (PCR-RFLP methods) for *ALDH2**1/*1, *ALDH2**1/*2 and *ALDH2**2/*2 mice. Left lane: MW marker (100 bp DNA ladder). Digestion of PCR products with *AclI* harbored three bands (430, 235 and 83 bp) in *ALDH2**1/*1 mice, four bands (665, 430, 235 and 83 bp) in *ALDH2**1/*2 mice and two bands (665 and 83 bp) in *ALDH2**2/*2 mice.

Alda-1 treatment on ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 mice

To investigate the effects of Alda-1, an ALDH2 activator, we intraperitoneally injected Alda-1 in ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 mice. As shown in Figure 5A, Alda-1 significantly elevated hepatic ALDH activity in all the human ALDH2 knock-in mice. The amount of alcohol drinking in ALDH2*1/*2 and ALDH2*2/*2 mice was significantly increased by Alda-1 treatment, but not in ALDH2*1/*1 mice (Figure 5B). Importantly, esophageal N²-ethylidene-dG level in both ALDH2*1/*2 and ALDH2*2/*2 mice

with alcohol drinking were significantly reduced by Alda-1 treatment despite the increased alcohol consumption, indicating the protective role of Alda-1 from alcohol-derived DNA damage in these mice (Figure 5C).

Effects of cyanamide, an ALDH2 inhibitor, on esophageal DNA damage in mice with alcohol drinking

Finally, we investigated whether cyanamide, an ALDH2 inhibitor, enhances alcohol-derived DNA damage in the esophagus of mice with alcohol drinking. As shown in Figure 6A, single intraperitoneal injection of cyanamide in C57BL/6N mice significantly reduced hepatic ALDH activity. Thereafter, we injected cyanamide to the C57BL/6N mice and let them drink 10% ethanol for 7 days. Cyanamide treatment resulted in a significant reduction of alcohol consumption (Figure 6B); however, esophageal N²-ethylidene-dG level was conversely significantly exacerbated by cyanamide treatment (Figure 6C).

Discussion

In this study, we established novel genetic engineering mouse, human ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 knock-in mice. So far, Aldh2 knockout mice have been widely used as an experimental mouse model for analyzing Aldh2 function; however, because ALDH2 dysfunction in human being mainly occurs as a result of having ALDH2*2 allele, Aldh2 knockout mice do not serve as bona fide model of human diseases related to acetaldehyde. In addition, because Aldh2 knockout mice lack Aldh2 protein, they cannot be used for examining the effect of ALDH2 enzymatic modulator such as ALDH2 activator and/or ALDH2 inhibitors. Indeed, treatment with ALDH2 activator (Alda-1) or ALDH2 inhibitor (cyanamide) did not affect Aldh2 activity in Aldh2 knockout mice (data not shown). To overcome disadvantages of Aldh2 knockout mice, we generated human ALDH2 knock-in mice. In this study, we newly established mice model, in which mouse Aldh2 allele was replaced to human ALDH2*1 (wild-type allele) or ALDH2*2 (mutant allele) by homologous recombination, and neomycin-resistant genes are also removed using CAG promoter-mediated Cre-loxP system. Here, we showed that ALDH2 activity in ALDH2*1/*2 and ALDH2*2/*2




Figure 3. Levels of ALDH2 expression and activity in human ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 knock-in mice. (A) Hepatic ALDH2 messenger RNA (mRNA) expression levels in ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 mice: ALDH2 mRNA expression level was measured by real-time RT-PCR. Relative ALDH2 mRNA levels in the liver of ALDH2*1/*2 and ALDH2*2/*2 mice relative to that of ALDH2*1/*1 mice is indicated. β-actin served as an internal control. (NS, not significant versus ALDH2*1/*1; n = 3). (B) Hepatic ALDH2 protein expression in human ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 knock-in mice: ALDH2 protein expression was demonstrated by western blotting. β-actin served as a loading control. (C) Hepatic ALDH activity in ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 mice. ALDH activity in mice with ALDH2*2 allele (ALDH2*1/*2 and/or ALDH2*2/*2 mice) were significantly lower than that in ALDH2*1/*1 mice. n = 6 *P < 0.05 versus ALDH2*1/*1 mice.




Figure 4. Influences of alcohol drinking on human ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 knock-in mice. (A) The amount of alcohol consumption per day in ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 mice. They were allowed to drink 10% ethanol freely for 7 days. The average amount of alcohol consumption per day in each mice is shown. n = 6, *P < 0.05, versus ALDH2*1/*1 mice. (B) Esophageal N²-ethylidene-dG levels in ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 mice that were allowed to drink 10% ethanol or water for 7 days. Alcohol drinking significantly increased esophageal N²-ethylidene-dG levels in all groups. Esophageal N²-ethylidene-dG levels in ALDH2*1/*2 and/or ALDH2*2/*2 mice with alcohol drinking were significantly higher than those in ALDH2*1/*1 mice with alcohol drinking. n = 6, *P < 0.05 versus each group of mice that drink water. *P < 0.05 versus ALDH2*1/*1 mice with alcohol drinking.

mice was significantly lower than that in $ALDH2^{*1/*1}$ mice, although the messenger RNA as well as protein expression levels of $ALDH2$ did not differ among them. Therefore, it is conceivable that reduced $ALDH2$ activity in $ALDH2^{*1/*2}$ and/or $ALDH2^{*2/*2}$ mice is caused by genetic polymorphism, but not by different expression levels of $ALDH2$. Notably, even with lesser

amount of ethanol intake, esophageal DNA damage levels with alcohol drinking were significantly higher in $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ mice than those in $ALDH2^{*1/*1}$ mice. This result suggests that esophagus of $ALDH2^{*1/*2}$ and/or $ALDH2^{*2/*2}$ mice is more susceptible to ethanol, probably due to decreased activity of $ALDH2$. Taken together, we established experimental mice model that possess identical polymorphism seen in human $ALDH2$ gene and would serve as an ideal experimental model for studying human diseases associated with impaired $ALDH2$ activity.




Figure 5. Effect of Alda-1 treatment on human $ALDH2^{*1/*1}$, $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ knock-in mice. (A) Hepatic $ALDH$ activity in $ALDH2^{*1/*1}$, $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ mice treated with Alda-1 or vehicle control. Alda-1 or vehicle control (20 mg/kg body weight) was intraperitoneally injected to $ALDH2^{*1/*1}$, $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ mice, and then liver of those mice was collected 3 h after the injection and hepatic $ALDH$ activity was measured. Alda-1 significantly increased hepatic $ALDH$ activity in $ALDH2^{*1/*1}$, $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ mice. $n = 8$, $P < 0.05$ versus each group of mice treated with vehicle control. (B) Amount of ethanol intake in $ALDH2^{*1/*1}$, $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ mice treated with Alda-1 or vehicle control. Alda-1 significantly increased the amount of alcohol drinking per day in both $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ mice. $n = 8$, $P < 0.05$ versus each group of mice treated with vehicle control. N.S.: no significant. (C) Esophageal N^2 -ethylidene-dG levels in $ALDH2^{*1/*1}$, $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ mice with alcohol drinking in the presence or absence of Alda-1 treatment. Alda-1 treatment resulted in a significant reduction of the N^2 -ethylidene-dG level in the esophagus of both $ALDH2^{*1/*2}$ and $ALDH2^{*2/*2}$ mice with alcohol drinking. $n = 8$, $P < 0.05$ versus each group of mice treated with vehicle control. N.S.: no significant.




Figure 6. Effect of cyanamide treatment on C57BL/6N mice. (A) Hepatic $ALDH$ activity in C57BL/6N mice treated with cyanamide or vehicle control. Cyanamide or vehicle control (1.5 mg/kg body weight) was intraperitoneally injected to C57BL/6N mice, and then liver of those mice was collected 3 h after the injection and hepatic $ALDH$ activity was measured. $n = 6$, $P < 0.05$ versus control. (B) The amount of alcohol consumption per day in C57BL/6N mice treated with cyanamide or vehicle control. They were allowed to drink 10% ethanol freely for 7 days. The average amount of alcohol consumption per day in each mice is shown. $n = 6$, $P < 0.05$ versus control. (C) Esophageal N^2 -ethylidene-dG levels in C57BL/6N mice with alcohol drinking in the presence or absence of cyanamide treatment. Cyanamide treatment significantly increased esophageal N^2 -ethylidene-dG level in C57BL/6N mice with alcohol drinking, compared with control. $n = 6$, $P < 0.05$ versus control.

In this study, we used Alda-1, a small molecule compound that has been identified as ALDH2 activators, and examined its effects on human ALDH2 knock-in mice. Alda-1 has been firstly reported by Chen *et al.*, identified by high-throughput screening using a fluorescent ALDH2 enzymatic assay based on the emission of resorufin (40). It was shown to be effective for an ischemic damage to the heart (40) and an acute inflammatory pain (50) in human ALDH2*1/*2 knock-in mice established in their group. In our present study, Alda-1 treatment significantly restored ALDH2 activity in ALDH2*1/*2 and ALDH2*2/*2 mice. It also increased the amount of alcohol drinking in those mice, nevertheless, esophageal DNA damage levels were significantly reduced by Alda-1 treatment, presumably due to the increased ALDH2 activity by Alda-1 treatment. Conversely, treatment with cyanamide, an ALDH2 inhibitor, reduced ALDH2 activity in C57BL/6N mice. Despite that cyanamide decreased the alcohol consumption in those mice, esophageal DNA damage levels were significantly enhanced by cyanamide treatment. These results indicate that modulation of ALDH2 activity using ALDH2 activator (Alda-1) or ALDH2 inhibitor (cyanamide) deeply affects the levels of esophageal DNA damage associated with alcohol drinking. Moreover, because esophageal DNA damage has been linked to esophageal carcinogenesis (51), Alda-1 might be effective in the prevention of alcohol-mediated ESCC via restoration of ALDH2 activity.

Molecular mechanisms underlying the activation of ALDH2 by Alda-1 is previously reported by Perez-Miller *et al.*, and they noted that Alda-1 increases ALDH2 enzymatic activity of ALDH2*2 because it acts as a structural chaperon and it restores the abnormal structure of ALDH2*2 (41). Furthermore, Alda-1 activates both ALDH2*1 and ALDH2*2 by binding at the entrance of the catalytic tunnel in close proximity to Cys302 and Glu286, which are critical to its substrate catalysis (41). In line with their report, we showed that Alda-1 treatment increased ALDH2 activity not only in ALDH2*1/*2 and ALDH2*2/*2 but also in ALDH2*1/*1 mice. Our results were consistent with a previous report that Alda-1 increased ALDH2 activity in human ALDH2*1/*1, ALDH2*1/*2 and ALDH2*2/*2 knock-in mice that were established by other groups independently (50). Although Alda-1 did not affect alcohol consumption as well as DNA adduct level in ALDH2*1/*1 mice, we presume that basal ALDH2 activity in wild-type ALDH2 might be inherently sufficient, and the increase of ALDH2 activity by Alda-1 in ALDH2*1/*1 may not have substantial benefit for protecting esophageal DNA damage.

In this study, we did not find histological abnormalities in the esophagus of ALDH2*1/*2 and ALDH2*2/*2 mice after drinking alcohol for 7 days. Although we made ALDH2*1/*2 mice drink 10% alcohol for as long as 12 months, even those mice did not develop esophageal cancer (data not shown). We suspect that more long-time drinking may be necessary for a development of ESCC, or other factors besides alcohol drinking alcohol may also be involved in the development of esophageal cancer, as there has been a report that esophageal dysplasia was caused by 4-nitroquinoline-1-oxide and ethanol intake for 10 weeks (52). Furthermore, we may be able to test the preventive effect of Alda-1 on alcohol-mediated ESCC development with those models.

The limitation of our study is that we could not evaluate the influences of Alda-1 on alcohol-abuse. As cyanamide is used as an alcohol deterrent drug for alcoholics, ALDH2 activators may have the risk of alcohol abuse due to enabling more alcohol drinking and therefore alcohol abuse. We should further define the indication for treatment of ALDH2 activators as a preventive agent for ESCC.

In conclusion, we established a novel mice model representing human ALDH2 gene polymorphism which causes reduced ALDH2 activity and is associated with various diseases including ESCC. We further showed that ALDH2 activation by Alda-1 in mice with ALDH2 dysfunction alleviates esophageal DNA damages associated with alcohol drinking. Our findings provide a new insight into alcohol-mediated esophageal carcinogenesis and prevention.

Funding

This work was supported by a Grant-in-Aid for Scientific Research 16K09281 (Shinya Ohashi), the Takeda Science Foundation (Shinya Ohashi), and Grant-in Aid from the Japanese Society of Gastroenterology (Shinya Ohashi), and for practical research for innovative cancer control from the Japan Agency for Medical Research and Development, AMED (Manabu Muto). None of the funding sources contributed to the writing of the manuscript.

Acknowledgements

The authors are grateful for the technical assistance from the Center for Anatomical, Pathological and Forensic Medical Researches and the Medical Research Support Center, Kyoto University Graduate School of Medicine.

Conflict of Interest Statement: The authors declare no competing interests.

References

- Ohashi, S. *et al.* (2015) Recent advances from basic and clinical studies of esophageal squamous cell carcinoma. *Gastroenterology*, 149, 1700–1715.
- Pennathur, A. *et al.* (2013) Oesophageal carcinoma. *Lancet*, 381, 400–412.
- Seitz, H.K. *et al.* (2007) Molecular mechanisms of alcohol-mediated carcinogenesis. *Nat. Rev. Cancer*, 7, 599–612.
- Brooks, P.J. *et al.* (2014) Acetaldehyde and the genome: beyond nuclear DNA adducts and carcinogenesis. *Environ. Mol. Mutagen.*, 55, 77–91.
- Mizumoto, A. *et al.* (2017) Molecular mechanisms of acetaldehyde-mediated carcinogenesis in squamous epithelium. *Int J Mol Sci.*, 18, 1943.
- Wang, M. *et al.* (2000) Identification of DNA adducts of acetaldehyde. *Chem. Res. Toxicol.*, 13, 1149–1157.
- Fang, J.L. *et al.* (1995) Development of a 32P-postlabelling method for the analysis of adducts arising through the reaction of acetaldehyde with 2'-deoxyguanosine-3'-monophosphate and DNA. *Carcinogenesis*, 16, 2177–2185.
- Lambert, B. *et al.* (1985) DNA cross-links in human leucocytes treated with vinyl acetate and acetaldehyde *in vitro*. *Mutat. Res.*, 146, 301–303.
- Obe, G. *et al.* (1986) Metabolism of ethanol *in vitro* produces a compound which induces sister-chromatid exchanges in human peripheral lymphocytes *in vitro*: acetaldehyde not ethanol is mutagenic. *Mutat. Res.*, 174, 47–51.
- Noori, P. *et al.* (2001) Mutational spectrum induced by acetaldehyde in the HPRT gene of human T lymphocytes resembles that in the p53 gene of esophageal cancers. *Carcinogenesis*, 22, 1825–1830.
- Singh, N.P. *et al.* (1995) Acetaldehyde: genotoxicity and cytotoxicity in human lymphocytes. *Mutat. Res.*, 337, 9–17.
- Dellarco, V.L. (1988) A mutagenicity assessment of acetaldehyde. *Mutat. Res.*, 195, 1–20.
- Helander, A. *et al.* (1991) Increased frequency of acetaldehyde-induced sister-chromatid exchanges in human lymphocytes treated with an aldehyde dehydrogenase inhibitor. *Mutat. Res.*, 264, 103–107.
- Secretan, B. *et al.*; WHO International Agency for Research on Cancer Monograph Working Group. (2009) A review of human carcinogens—Part E: tobacco, areca nut, alcohol, coal smoke, and salted fish. *Lancet Oncol.*, 10, 1033–1034.
- Crabb, D.W. *et al.* (2004) Overview of the role of alcohol dehydrogenase and aldehyde dehydrogenase and their variants in the genesis of alcohol-related pathology. *Proc. Nutr. Soc.*, 63, 49–63.

16. Higuchi, S. et al. (1995) Alcohol and aldehyde dehydrogenase polymorphisms and the risk for alcoholism. *Am. J. Psychiatry*, 152, 1219–1221.
17. Wang, M.F. et al. (2009) Substrate specificity of human and yeast aldehyde dehydrogenases. *Chem. Biol. Interact.*, 178, 36–39.
18. Yoshida, A. et al. (1984) Molecular abnormality of an inactive aldehyde dehydrogenase variant commonly found in Orientals. *Proc. Natl. Acad. Sci. U. S. A.*, 81, 258–261.
19. Li, H. et al. (2009) Refined geographic distribution of the oriental ALDH2*504Lys (nee 487Lys) variant. *Ann. Hum. Genet.*, 73(Pt 3), 335–345.
20. Larson, H.N. et al. (2005) Disruption of the coenzyme binding site and dimer interface revealed in the crystal structure of mitochondrial aldehyde dehydrogenase “Asian” variant. *J. Biol. Chem.*, 280, 30550–30556.
21. Chiang, C.P. et al. (2016) Ethanol-metabolizing activities and isozyme protein contents of alcohol and aldehyde dehydrogenases in human liver: phenotypic traits of the ADH1B*2 and ALDH2*2 variant gene alleles. *Pharmacogenet. Genomics*, 26, 184–195.
22. Farrés, J. et al. (1994) Effects of changing glutamate 487 to lysine in rat and human liver mitochondrial aldehyde dehydrogenase. A model to study human (Oriental type) class 2 aldehyde dehydrogenase. *J. Biol. Chem.*, 269, 13854–13860.
23. Zhou, J. et al. (2000) Basis for half-of-the-site reactivity and the dominance of the K487 oriental subunit over the E487 subunit in heterotetrameric human liver mitochondrial aldehyde dehydrogenase. *Biochemistry*, 39, 12019–12024.
24. Matsuo, K. et al. (2006) Alcohol dehydrogenase 2 His47Arg polymorphism influences drinking habit independently of aldehyde dehydrogenase 2 Glu487Lys polymorphism: analysis of 2,299 Japanese subjects. *Cancer Epidemiol. Biomarkers Prev.*, 15, 1009–1013.
25. Nakamura, Y. et al. (2002) Genetic variation in aldehyde dehydrogenase 2 and the effect of alcohol consumption on cholesterol levels. *Atherosclerosis*, 164, 171–177.
26. Yokoyama, A. et al. (1996) Cancer screening of upper aerodigestive tract in Japanese alcoholics with reference to drinking and smoking habits and aldehyde dehydrogenase-2 genotype. *Int. J. Cancer*, 68, 313–316.
27. Matsuo, K. et al. (2001) Gene-environment interaction between an aldehyde dehydrogenase-2 (ALDH2) polymorphism and alcohol consumption for the risk of esophageal cancer. *Carcinogenesis*, 22, 913–916.
28. Muto, M. et al. (2005) Risk of multiple squamous cell carcinomas both in the esophagus and the head and neck region. *Carcinogenesis*, 26, 1008–1012.
29. Yang, S.J. et al. (2010) Relationship between genetic polymorphisms of ALDH2 and ADH1B and esophageal cancer risk: a meta-analysis. *World J. Gastroenterol.*, 16, 4210–4220.
30. Yokoyama, A. et al. (2001) Alcohol and aldehyde dehydrogenase gene polymorphisms and oropharyngolaryngeal, esophageal and stomach cancers in Japanese alcoholics. *Carcinogenesis*, 22, 433–439.
31. Yokoyama, A. et al. (2002) Genetic polymorphisms of alcohol and aldehyde dehydrogenases and glutathione S-transferase M1 and drinking, smoking, and diet in Japanese men with esophageal squamous cell carcinoma. *Carcinogenesis*, 23, 1851–1859.
32. Katada, C. et al. (2014) Factors associated with the presence of multiple Lugol-voiding lesions in patients with esophageal squamous-cell carcinoma. *Dis. Esophagus*, 27, 457–462.
33. Lee, C.H. et al. (2009) Genetic modulation of ADH1B and ALDH2 polymorphisms with regard to alcohol and tobacco consumption for younger aged esophageal squamous cell carcinoma diagnosis. *Int. J. Cancer*, 125, 1134–1142.
34. Muto, M. et al. (2002) Association between aldehyde dehydrogenase gene polymorphisms and the phenomenon of field cancerization in patients with head and neck cancer. *Carcinogenesis*, 23, 1759–1765.
35. Yokoyama, A. et al. (2010) Polymorphisms of alcohol dehydrogenase-1B and aldehyde dehydrogenase-2 and the blood and salivary ethanol and acetaldehyde concentrations of Japanese alcoholic men. *Alcohol. Clin. Exp. Res.*, 34, 1246–1256.
36. Aoyama, I. et al. (2017) Establishment of a quick and highly accurate breath test for ALDH2 genotyping. *Clin. Transl. Gastroenterol.*, 8, e96.
37. Yokoyama, A. et al. (2008) Salivary acetaldehyde concentration according to alcoholic beverage consumed and aldehyde dehydrogenase-2 genotype. *Alcohol. Clin. Exp. Res.*, 32, 1607–1614.
38. Hori, H. et al. (1997) Genetic polymorphisms of tobacco- and alcohol-related metabolizing enzymes and human esophageal squamous cell carcinoma susceptibility. *J. Clin. Gastroenterol.*, 25, 568–575.
39. Amanuma, Y. et al. (2015) Protective role of ALDH2 against acetaldehyde-derived DNA damage in oesophageal squamous epithelium. *Sci. Rep.*, 5, 14142.
40. Chen, C.H. et al. (2008) Activation of aldehyde dehydrogenase-2 reduces ischemic damage to the heart. *Science*, 321, 1493–1495.
41. Perez-Miller, S. et al. (2010) Alda-1 is an agonist and chemical chaperone for the common human aldehyde dehydrogenase 2 variant. *Nat. Struct. Mol. Biol.*, 17, 159–164.
42. Mishina, M. et al. (2007) Conditional gene targeting on the pure C57BL/6 genetic background. *Neurosci. Res.*, 58, 105–112.
43. Hayashida, M. et al. (2010) Genotyping of polymorphisms in alcohol and aldehyde dehydrogenase genes by direct application of PCR-RFLP on dried blood without DNA extraction. *Anal. Sci.*, 26, 503–505.
44. Yukawa, Y. et al. (2012) Combination of ADH1B*2/ALDH2*2 polymorphisms alters acetaldehyde-derived DNA damage in the blood of Japanese alcoholics. *Cancer Sci.*, 103, 1651–1655.
45. Wang, M. et al. (2006) Identification of an acetaldehyde adduct in human liver DNA and quantitation as N2-ethyldeoxyguanosine. *Chem. Res. Toxicol.*, 19, 319–324.
46. Matsuda, T. et al. (2007) Increased formation of hepatic N2-ethylidene-2'-deoxyguanosine DNA adducts in aldehyde dehydrogenase 2-knockout mice treated with ethanol. *Carcinogenesis*, 28, 2363–2366.
47. Nagasawa HT, et al. (1990) Evidence for nitroxyl in the catalase-mediated bioactivation of the alcohol deterrent agent cyanamide. *J. Med. Chem.*, 33, 3120–3102.
48. Takaoka, M. et al. (2004) Epidermal growth factor receptor regulates aberrant expression of insulin-like growth factor-binding protein 3. *Cancer Res.*, 64, 7711–7723.
49. Yoshioka, M. et al. (2017) Distinct effects of EGFR inhibitors on epithelial- and mesenchymal-like esophageal squamous cell carcinoma cells. *J. Exp. Clin. Cancer Res.*, 36, 101.
50. Zambelli, V.O. et al. (2014) Aldehyde dehydrogenase-2 regulates nociception in rodent models of acute inflammatory pain. *Sci. Transl. Med.*, 6, 251ra118.
51. Garaycochea, J.I. et al. (2018) Alcohol and endogenous aldehydes damage chromosomes and mutate stem cells. *Nature*, 553, 171–177.
52. Osei-Sarfo, K. et al. (2015) Initiation of esophageal squamous cell carcinoma (ESCC) in a murine 4-nitroquinoline-1-oxide and alcohol carcinogenesis model. *Oncotarget*, 6, 6040–6052.