

The classification of log del Pezzo surfaces and their quasi-universal coverings

By D. Q. ZHANG

(Department of Mathematics, Osaka University)

We Work over an algebraically closed field k of characteristic zero. Let V be a nonsingular projective surface over k and D a reduced effective divisor on V with only simple normal crossings.

Write $D = \sum_{i=1}^n D_i$, where D_i 's are irreducible components.

Definition 1 (cf. [61]). $Bk(D) = \sum_{i=1}^n \alpha_i D_i$ is defined as the effective $\mathbb{Q}$ -divisor satisfying :

- (i) $\text{Supp } Bk(D)$ consists of all connected components of D which are contractible to quotient singularities and which contains no (-1) curves, and all admissible rational twigs T_i 's in D , i.e., by definition, $T := T_i$ is written as $T = G_1 + \cdots + G_s$ such that: $G_1 \cong \mathbb{P}^1$, $(G_1^2) \leq -2$, $(G_i, G_j) \leq 1$ and $(G_i, G_j) = 1$ if and only if $j-i = \pm 1$, $D-T$ does not meet $G_1 + \cdots + G_{s-1}$ when $s \geq 2$.
- (ii) $(D - Bk(D) + K_V, D_i) = 0$, for every $D_i \subseteq \text{Supp } Bk(D)$.

Then $0 \leq \alpha_i \leq 1$, where $i=1, 2, \dots, n$.

Denote by $D^\#$ the effective $\mathbb{Q}$ -divisor $D - Bk(D) = \sum_{i=1}^n \beta_i D_i$

with $\beta_i = 1 - \alpha_i$ ($1 \leq i \leq n$).

If $\bar{\kappa}(V-D) \geq 0$, $D + K_V = (D^\# + K_V) + Bk(D)$ is the Zariski decomposition, where $D^\# + K_V$ is the numerically effective part and $Bk(D)$ is the negative definite part.

Lemma 2 (cf. [6]). *We have $0 \leq \beta_i \leq 1$. More precisely,*

- (i) $\beta_i = 0$ if and only if every connected component F of D containing D_i is contractible to a rational double point and there are no (-1) curves in F .
- (ii) $\beta_i < 1$ if and only if $D_i \subseteq \text{Supp } Bk(D)$.

In [6] M. Miyanishi and S. Tsunoda constructed an almost minimal model $\sigma: (V, D) \rightarrow (\tilde{V}, \tilde{D})$ for each pair (V, D) . $(\tilde{V}, \tilde{D})$ is almost minimal. Namely, for every irreducible curve $\tilde{E}$ on $\tilde{V}$, either $(\tilde{E}, \tilde{D}^\# + K_{\tilde{V}}) \geq 0$ or the intersection matrix of $\tilde{E} + Bk(\tilde{D})$ is not negative definite. $(\tilde{V}, \tilde{D})$ has properties : (i) $\sigma_*(D) = \tilde{D}$, (ii) $\bar{\kappa}(\tilde{V} - \tilde{D}) = \bar{\kappa}(V - D)$, etc.

In virtue of this result, we shall assume in the following arguments that (V, D) is almost minimal.

Y. Kawamata proved (cf. [5]) :

Suppose (V, D) is almost minimal. Then $D^\# + K_V$ is numerically effective if and only if $\bar{\kappa}(V - D) \geq 0$.

By the study done by M. Miyanishi and S. Tsunoda (cf. [6, 7]),

we know well the structure of a pair (V, D) with $\bar{\kappa}(V-D)=-\infty$, unless (V, D) is the following log del Pezzo surface of rank one with contractible boundaries; in this case V is rational.

Definition 3. A pair (V, D) is called a log del Pezzo surface with contractible boundaries if

(i) D is contractible to quotient singular points on a projective normal surface by the contraction morphism

$g: V \rightarrow \bar{V}$; there are no (-1) curves in D ,

i.e., g is a minimal resolution of $\text{Sing}(\bar{V})$;

(ii) the anticanonical divisor $-K_{\bar{V}}$ is ample.

Moreover, if the Picard number $\rho(\bar{V}) := \text{rk}(\text{Pic}(\bar{V}) \otimes_{\mathbb{Z}} \mathbb{Q}) = 1$, (V, D) is said to be of rank one.

We often confuse (V, D) with $\bar{V}$.

Let (V, D) and $g: V \rightarrow \bar{V}$ be the same as in Definition 3. Then $-(D^{\#} + K_V) (\equiv g^*(-K_{\bar{V}}))$ is numerically effective as a $\mathbb{Q}$ -divisor and for every curve E on V , $-(E, D^{\#} + K_V) = 0$ if and only if $E \leq D$. Fix a natural number P such that $PD^{\#}$ is integral. Then $-P(G, D^{\#} + K_V) \in \mathbb{Z}_+$ for every curve G . Thus, we can fix an irreducible curve C such that $-(C, D^{\#} + K_V)$ attains the smallest positive value.

Lemma 4 (M. Miyanishi). (I) If $|C + D + K_V| \neq \emptyset$ (called the case (α)), then $(V, C + D)$ is a quasi-Iitaka surface (see Definition 5 below) by replacing C by a member of $|C|$ if necessary.

(II) If $|C + D + K_V| = \emptyset$ and $(V, D) \neq (\Sigma_n, M_n)$ where Σ_n is a

Hirzebruch surface of deg n and M_n is its minimal section (called the case (β)), then we may assume that C is a (-1) curve by replacing C by a curve C' with $|C'+D+K_V| = \emptyset$ and $-(C', D^\# + K_V) = -(C, D^\# + K_V)$.

Definition 5. A pair (W, B) with W rational is said to be a **quasi-Iitaka surface** if there exists a decomposition of B into reduced effective integral divisors $B = A + N$ such that

$$(i) \quad A + K_W \sim 0,$$

(ii) N is contractible to rational double singular points; there are no (-1) curves in N , hence $(\text{Supp } N) \cap (\text{Supp } A) = \emptyset$.

Moreover, if B has only simple normal crossings, (W, B) is called an **Iitaka surface**.

We sketch how Iitaka surfaces are classified (cf. [10]).

Let $h: W \rightarrow \bar{W}$ be the contraction of N to rational double singular points. Then, since $K_{\bar{W}} \sim -h_* A$, $K_{\bar{W}}$ is not numerically effective. Applying the Mori theory, we can find an extremal rational curve $\bar{L}$ on $\bar{W}$. Namely, $-3 \leq (\bar{L}, K_{\bar{W}}) < 0$, and if Z_1 and Z_2 are in the closure of the cone $\overline{NE}(\bar{W})$ of effective 1-cycles and $Z_1 + Z_2 \in \mathbb{R}_+[\bar{L}]$, then $Z_1, Z_2 \in \mathbb{R}_+[\bar{L}]$. There exists a numerically effective divisor $\bar{H}$ on $\bar{W}$ such that $\bar{H}^\perp \cap \overline{NE}(\bar{W}) = \mathbb{R}_+[\bar{L}]$.

Case(1) $\bar{H} \equiv 0$. Then $\rho(\bar{W}) = 1$ and $-K_{\bar{W}}$ is ample.

Lemma 6. *If $\rho(\bar{W})=1$ (hence $\rho(W) = \#(N)+1$) then $\bar{W}$ is a rational Gorenstein log del Pezzo surface of rank 1. So, N has one of the following 27 Dynkin types (cf. [41], [9], [3], [10]).*

$A_1, A_1+A_2, A_4, 2A_1+A_3, D_5, A_1+A_5, 3A_2, E_6, 3A_1+D_4, A_7,$
 $A_1+D_6, E_7, A_1+2A_3, A_2+A_5, D_8, 2A_1+D_6, E_8, A_1+E_7, A_1+A_7, 2A_4,$
 $A_8, A_1+A_2+A_5, A_2+E_6, A_3+D_5, 4A_2, 2A_1+2A_3, 2D_4.$

Case(2) $\bar{H} \neq 0$ and $(\bar{H}^2)=0$. Then $\bar{H} \in \mathbb{R}_+[\bar{\ell}]$ and $(\bar{\ell}^2)=0$. Suppose $W \not\cong \mathbb{P}^2$ or Σ_n . Then, for $N \gg 0$ such that $N\bar{\ell}$ is a Cartier divisor, $\Phi|_{h*N\bar{\ell}}$ is composed with a $\mathbb{P}^1$ -fibration $\Phi: W \rightarrow \mathbb{P}^1$. Every singular fiber of Φ is written as $f_1 = 2(E+B_1+\dots+B_{s-2})+B_{s-1}+B_s$, where E is a (-1) curve and B_i 's are irreducible components of N .

Case(3) $(\bar{H}^2) > 0$. Then the proper transform $\ell = h'\bar{\ell}$ is a (-1) curve. Furthermore, if ℓ is not an irreducible component of A then the connected component R of $\ell+N$ containing ℓ is contractible to a smooth point. Indeed, R is a rod and ℓ is a tip. Let $\sigma: W \rightarrow \bar{W}$ be the contraction of ℓ if ℓ is an irreducible component of A and of R otherwise. Consider a quasi-litaka surface (W, σ_*B) and apply the Mori theory again. After a finitely many steps, we are reduced to the case (1) or (2) above.

This completes the classification of litaka surfaces. For quasi-litaka surfaces, this process works, too.


We now consider the case(β) of Lemma 4. We study this case according as how many irreducible components of D intersect with the (-1) curve C . We recall that a surface $V-D$ is called affine uniruled (resp. affine-ruled) if there exists a dominant morphism (resp. open immersion) $\mathbb{A}^1 \times G \rightarrow V-D$ with an affine curve G .

Theorem 7. *Suppose that the case(β) occurs, i.e., $|C+D+K_V| = \emptyset$ and $(V,D) \neq (\Sigma_n, M_n)$, and that $V-D$ is not affine-ruled. Suppose the following case($\beta-1$) does not occur.*

Case($\beta-1$) The (-1) curve C meets exactly two irreducible components D_1 and D_2 of D with $(D_1^2) = -2$ and $(D_2^2) \leq -3$.

Then there exists a $\mathbb{P}^1$ -fibration $\Phi: V \rightarrow \mathbb{P}^1$ such that all singular fibers of Φ and the configuration of $C+D$ are precisely described as follows :

Group(I). It consists of the following 8 Figures.



$$b = -(D_7^2) - (D_8^2) - 4$$

Figure (1)





Figure (2)



Figure(3)




Figure 17)




Figure (8)




Figure (4)




Figure (5)





Figure (6)

Group(I). (V, D) is obtained by blowing up a quasi-litaka surface (W, B) in the way shown below, where $B=A+N$, A is a cuspidal curve in $|-K_W|$ and $\rho(W) = \#(N)+1$ (hence all possible Dynkin types of N are given in Lemma 6).



$\xleftarrow{\sigma}$

$$\begin{array}{ccc} -2 & -3 & -\alpha \\ \left| \begin{array}{c} D_1 \\ - \\ - \\ - \end{array} \right| & \left| \begin{array}{c} D_2 \\ - \\ - \\ - \end{array} \right| & \left| \begin{array}{c} D_3 = \sigma^*(A) \\ - \\ - \\ -1 \end{array} \right| C \end{array}$$

$$(A^2) = 1, 2 \text{ or } 3;$$

$$\alpha = 6 - (A^2) = 5, 4 \text{ or } 3.$$

We set $D = D_1 + D_2 + D_3 + \sigma^*(N)$.

Group(III). C meets only one component D_1 of D and the connected component Δ_1 of D containing D_1 is a fork. Moreover, either D_1 is the central component of Δ_1 with $(D_1^2) = -2$ or D_1 is contained in a twig T_1 of Δ_1 such that $C+T_1$ is negative definite. Let $\sigma: V \rightarrow W$ be the contraction of C in the first case and all contractible curves of $C+T_1$ in the second case. Then $(W, \sigma_*(D))$ is a log del Pezzo surface of rank one with non-contractible boundaries, M. Miyanishi and S. Tsunoda proved :

There exists a $\mathbb{P}^1$ -fibration $\Psi: W \rightarrow \mathbb{P}^1$ where all singular fibers and the configuration of $\sigma_*(D)$ are described precisely in [7; Lemma 2.6 and Theorems 4 and 6].

So, the configuration of $C+D$ and all singular fibers of $\Phi := \Psi \circ \sigma: V \rightarrow \mathbb{P}^1$ can be obtained.

(*) The surfaces given in Figures 2 ~ 8 and *Group(I)* are deduced from our unpublished notes.

Lemma 8 (M. Miyanishi). Suppose that the case $(\beta-1)$ occurs, i.e., $|C+D+K_V| = \emptyset$, $(V,D) \neq (\Sigma_n, M_n)$ and C meets exactly two irreducible components D_1 and D_2 of D with $(D_1^2) = -2$ and $(D_2^2) \leq -3$. Let $\eta: V \rightarrow W$ be the contraction of the (-1) curve C . Then $(W, \eta_*(D-D_1))$ is a log del Pezzo surface of rank one with contractible boundaries.

Combining the above results, we get a method to classify log del Pezzo surfaces of rank one with contractible boundaries. Actually, in some easy cases, we can obtain all possibilities of pairs (V,D) fitting the case $(\beta-1)$ by Lemma 8.

Definition 9. A log del Pezzo surface (V,D) of rank one with contractible boundaries is called a **dP3** surface if $\bar{V}$, with the contraction morphism $g: V \rightarrow \bar{V}$, has exactly one rational triple and several rational double singular points.

Let (V,D) be a dP3 surface fitting the case $(\beta-1)$. Then $\eta_*(D-D_1)$ consists of (-2) curves and (-2) forks, where $\eta: V \rightarrow W$ is the contraction of C . By Lemma 2, (i), $-K_W = -((\eta_*(D-D_1))^{\#} + K_W)$, which is numerically effective. Note that $0 < (K_W^2) = 10 - \rho(W) = 9 - \#(\eta_*(D-D_1)) \leq 7$. So, W is obtained by blowing up $9 - (K_W^2) (\leq 8)$ points on $\mathbb{P}^2$, which might include infinitely near points. Demazure proved that there is a nonsingular elliptic curve A in $|-K_W|$ (cf. [2; Theorem 1, p.39]). So, $(W, A + \eta_*(D-D_1))$ is an Iitaka surface with $\rho(W) = \#(\eta_*(D-D_1)) + 1$. Thus, $\eta_*(D-D_1)$

has one of 27 Dynkin types given in Lemma 6. By considering suitable $\mathbb{P}^1$ -fibrations, we can work out all possibilities of pairs (V, D) fitting the case $(\beta-1)$. At the same time we find a $\mathbb{P}^1$ -fibration $\Psi: V \rightarrow \mathbb{P}^1$ for each pair (V, D) , and write out explicitly the configuration of $C+D$ and singular fibers of Ψ .

Now we let the field $k = \mathbb{C}$.

Theorem 10 (cf. [12]). *Let (V, D) be a $dP3$ surface. Then we have:*

- (I) *The configuration of D is No. n of Table 1 ($1 \leq n \leq 97$), which is given at the end of the paper.*
- (II) *We find a $\mathbb{P}^1$ -fibration $\Psi: V \rightarrow \mathbb{P}^1$ and write out explicitly the configuration *Picture*(n) of $C+D$ and all singular fibers of Ψ (cf. *Picture*(20) below). By reason of spaces, we omit them.*
- (III) *$\pi_1(V^0)$ is a finite group. The quasi-universal covering $\bar{U}$ of $\bar{V}$ (cf. Definition 11 below) is a log del Pezzo surface with contractible boundaries and $\bar{U}$ is rational.*
- (IV) *Suppose $\pi_1(V^0) = (0)$. Then $V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$ where $\mathbb{C}^* := \mathbb{C} - \{0\}$.*
- (V) *Suppose $\pi_1(V^0) \neq (0)$. Then $\bar{V}$ is the quotient of $\mathbb{P}^2$ by a finite subgroup H of $\text{PGL}(2, \mathbb{C})$ iff $\rho(\bar{U}) = 1$. If this is the case, there is a cyclic normal subgroup H_1 of H such that $H/H_1 \cong \pi_1(V^0)$ and $\mathbb{P}^2/H_1 \cong \bar{U}$.*

Definition 11. *Let U^0 be the topologically universal covering of $V-D$. If the fundamental group $\pi_1(V-D)$ is finite, then U^0 is algebraic and we let $\bar{U}$ be the normalization of*

$\bar{V}$ in the function field $k(U^0)$. $\bar{U}$ is said to be the quasi universal covering of $\bar{V}$, which is unique by Zariski Main Theorem.

In the joint work with M.Miyanishi we prove the following Theorem 12 (cf. [8; Theorems 1 and 2]).

Theorem 12. *Let S be a singular rational Gorenstein log del Pezzo surface with $\rho(S)=1$. Then the quasi-universal covering $\tilde{S}$ of S exists and $\tilde{S}$ is given in Table 2 at the end of the present paper.*

Moreover, we have :

- (1) $S^0 := \text{Reg}(S)$ is simply connected, i.e., $\pi_1(S^0)=0$ if and only if S is a Gorenstein algebraic compactification of the affine plane $\mathbb{C}^2$ by an irreducible curve at infinity; for another proof of the "if" part, see Brenton[1; Theorem 6].
- (2) Suppose $\pi_1(S^0) \neq 0$. Then $S \cong \mathbb{P}^2/G$ for a finite group $G \subseteq \text{PGL}(2;k)$ if and only if the Picard number $\rho(\tilde{S}) = 1$.

Remark. Actually, we proved that S^0 is simply connected if and only if S is a Gorenstein algebraic compactification of the affine plane $\mathbb{C}^2$ (the boundary might be *reducible*) under the assumption that S is a singular rational Gorenstein log del Pezzo surface (not necessarily $\rho(S) = 1$).

We want to construct the quasi-universal covering of a dP3 surface. We need the following Proposition 13 which is

proved by the purity of the branch locus.

Proposition 13. *Let Y be a normal surface with only quotient singularities and let $\pi: X \rightarrow Y$ be a finite morphism which is étale outside $\text{Sing}(Y)$. Then X has only quotient singularities. In particular, if Y is a log del Pezzo surface so is X , because $-K_X = \pi^*(-K_Y)$.*

Corollary 14. *Let (V, D) be a dP3 surface and $g: V \rightarrow \bar{V}$ the contraction of D . Suppose that there exist integral divisors F and Δ on V and an integer $l \geq 2$ such that $\Delta > 0$, $\Delta \sim lF$ and every irreducible components of Δ is contained in D and has coefficient less than l . Denote by $\tau: X \rightarrow V$ the normalization of the covering surface which is defined by the relation $\Delta \sim lF$ and which hence branches exactly over Δ . Then $\tau^{-1}(D)$ is contractible to points on a projective normal surface $\bar{X}$ and τ induces a finite morphism $\bar{\tau}: \bar{X} \rightarrow \bar{V}$ such that $\bar{\tau} = \tau$ over $\text{Reg}(\bar{V})$ (hence $\bar{\tau}$ is étale outside $\text{Sing}(\bar{V})$). So, $\bar{X}$ is a log del Pezzo surface.*

For the computation of $H_1(V-D; \mathbb{Z})$ we use

Lemma 15 (cf. [8; Lemma 11]). *Let (V, D) be a dP3 surface. Then we have :*

$$H_1(V-D; \mathbb{Z}) \cong (H^2(D; \mathbb{Z})/H_2(D; \mathbb{Z})) / (Cl(\bar{V})/Pic(\bar{V})).$$

We explain our method of finding the quasi-universal covering of a dP3 surface $\bar{V}$ by the following :

Example 16. Consider the No.20 surface (V, D) of Theorem 10 (cf. Picture(20) below). Let $v: V \rightarrow \Sigma_2$ be the contraction of curves in the singular fibers of the vertical fibration such that $(v_* D_3)^2 = -2$. Then we have :

$M := v_* D_3$, $L := v_* D_2$, $v_* D_4 \sim M + 3L$, $v_* D_1 \sim 2M + 4L$, $L \sim v_* E \sim v_* E_1$. These implies $5v^*(M+L) \sim v^* v_* D_4 + 2v^* v_* D_2 + 4D_3$. Hence if we let

$\Delta = 2D_2 + 4D_3 + D_4 + 3D_5 + D_6 + 2D_7 + 3D_8 + 4D_9$ and $F = v^* L + D_3 - E_2 - E_3$, then $\Delta \sim 5F$. By Corollary 14, there exists a finite morphism $\bar{\tau}$:

$\bar{X} \rightarrow \bar{V}$ which is étale outside $\text{Sing}(\bar{V})$ and which has $\deg \bar{\tau} = 5$

(where $g: V \rightarrow \bar{V}$ is the contraction of D). Let $h: U \rightarrow \bar{X}$

be the minimal resolution of $\text{Sing}(\bar{X})$. Then $h^{-1}(\text{Sing } \bar{X}) =$

$\sum_{i=1}^5 H_i$ (cf. Picture(20.1) below). We see $\bar{X} - \text{Sing}(\bar{X}) = U -$


$h^{-1}(\text{Sing } \bar{X}) \supseteq \mathbb{C}^2$. Indeed, let $u: U \rightarrow \Sigma_0$ be the contraction

of $C_1, C_4, H_4, H_2, C_3, F_2, H_5$ and H_3 . Then $\Sigma_0 - u_* \left(\sum_{i=1}^5 H_i \right)$


$\supseteq \mathbb{C}^2$. So, $\bar{X} - \text{Sing}(\bar{X})$ is simply connected. This implies

that $\bar{\tau}^{-1}(\bar{V} - \text{Sing}(\bar{V}))$ is simply connected. Hence $\bar{X}$ is


the quasi-universal covering of $\bar{V}$ and $\pi_1(V-D) \cong \mathbb{Z}/5\mathbb{Z}$.




Picture (20)



Picture (20.1)



Picture (20)



Picture (20.1)

Table 1

[No]	Sing. type of $\bar{V}$	$ H_1(V^0; \mathbb{Z}) $	$\pi_1(V^0)$	$ \rho(\bar{U}) $	Sing. type of $\bar{U}$	Ruledness of V^0, U^0
[1]	*	(0)	(0)	1	$\bar{U} = \bar{V} = \bar{\Sigma}_3$	$V^0 \supseteq \mathbb{C}^2$
[2]	$A_4 + (*-o^4)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[3]	$*-o^8$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[4]	$A_1 + (*-o^7)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[5]	$D_6 + (o-*o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$A_7 + (-4)$	$U^0 \supseteq \mathbb{C}^2$
[6]	$2A_1 + D_4 + (o-*o)$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$ \pi_1 = 16$	2	$\bar{U} = \Sigma_0$	$U^0 = \Sigma_0$
[7]	$2A_3 + (o-*o)$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	D_4 or Q_3	4	$2A_1$	$U^0 \supseteq \mathbb{C}^2$
[8]	$A_1 + A_2 + (*-o^5)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[9]	$D_5 + (*-o^3)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[10]	$2A_1 + A_3 + (*-o^3)$	$\mathbb{Z}/2\mathbb{Z}$	S_3	3	$3A_1 + 2*$	$U^0 \supseteq \mathbb{C}^2 - P$
[11]	$A_1 + A_5 + (*-o^2)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$A_2 + 2(*-o-o)$	$U^0 \supseteq \mathbb{C}^2$
[12]	$E_6 + (*-o^2)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[13]	$A_1 + D_6 + (*-o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$D_4 + 2(*-o)$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[14]	$A_7 + (*-o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	3	$A_3 + 2(*-o)$	$U^0 \supseteq \mathbb{C}^2$
[15a]	$E_7 + (*-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[15b]	$E_7 + (*-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[16]	$D_8 + *$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	3	$D_5 + 2*$	$U^0 \supseteq \mathbb{C}^2 - P$
[17]	$A_1 + E_7 + *$	$\mathbb{Z}/2\mathbb{Z}$	S_3	4	D_4	$U^0 \supseteq \mathbb{C}^2$
[18a]	$E_8 + *$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[18b]	$E_8 + *$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
[19]	$A_1 + A_7 + *$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	5	$A_1 + 4*$	$U^0 \supseteq \mathbb{C}^2 - P$
[20]	$2A_4 + *$	$\mathbb{Z}/5\mathbb{Z}$	$\mathbb{Z}/5\mathbb{Z}$	5	$5*$	$U^0 \supseteq \mathbb{C}^2$
[21]	$A_8 + *$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	5	$A_2 + 3*$	$U^0 \supseteq \mathbb{C}^2$

22	$A_1 + A_2 + A_5 + *$	$\mathbb{Z}/6\mathbb{Z}$	$ \pi_1 = 18$	4	smooth del Pezzo surface of deg 6	$U^0 = \bar{U}$
23	$3A_2 + (*-o^2)$	$\mathbb{Z}/3\mathbb{Z}$	$ \pi_1 = 21$	1	$\bar{U} = \mathbb{P}^2$	$U^0 = \mathbb{P}^2$
24	$A_2 + A_5 + (*-o)$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	3	$A_1 + 3(*-o)$	$U^0 \supseteq \mathbb{C}^2$
25	$A_2 + E_6 + *$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	3	$D_4 + 3*$	$U^0 \supseteq \mathbb{C}^2$
26	$A_3 + D_5 + *$	$\mathbb{Z}/4\mathbb{Z}$	$ \pi_1 = 12$	6	smooth del Pezzo surface of deg 4	$U^0 = \bar{U}$
27	$A_1 + 2A_3 + (*-o)$	$ H_1 = 4$	$ \pi_1 = 20$	2	$\bar{U} = \Sigma_0$	$U^0 = \Sigma_0$
28	$2A_1 + D_5 + *$	$\mathbb{Z}/2\mathbb{Z}$	S_3	1	$\bar{U} = \bar{\Sigma}_2$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^{**}$
29	$2A_1 + (*-o^3 - \overset{\circ}{o}-o)$	$\mathbb{Z}/2\mathbb{Z}$	S_3	5	$*-o-*$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^{**}$
30	$A_1 + (*-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C}^2$
31	$2A_1 + (o-* - o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	1	$\bar{U} = \bar{\Sigma}_4$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
32	$A_1 + (o-* - o - o - o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$(-4) - o$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
33	$A_1 + (o-* - o - \overset{\circ}{o} - o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$(-4) - \overset{\circ}{o} - o$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
34	$3A_1 + (o - \overset{\circ}{*} - o)$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	1	$\bar{U} = \bar{\Sigma}_6$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^{**}$
35	$2A_1 + (o - o - o - \overset{\circ}{*} - o)$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	3	$o - (-6) - o$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^{**}$
36	$*-o-o-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C}^2$
37	$o-o-o-* - o-o-o$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	3	$o - (-4) - o$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
38	$*-o-o-o-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C}^2$
39	$o-* - o-o-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C}^2$
40	$A_1 + (*-o^5)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
41	$A_1 + (o-o-* - o^3)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
42	$*-o^7$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
43	$o-o-o-* - o^4$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
44	$A_2 + (o-* - o^4)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
45	$o-* - o^7$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$
46	$o-o-o-o-* - o^4$	(0)	(0)	1	$\bar{U} = \bar{V}$	$V^0 \supseteq \mathbb{C} \times \mathbb{C}^*$

47	$A_1 + (o-o-*o^5)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
48	$o-o-o-*o-\overset{\circ}{o}-o$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	3	$o-(-4)-\overset{\circ}{o}-o$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
49	$o-o-o-*o-o-\overset{\circ}{o}-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
50	$*-o-o-o-o-o-\overset{\circ}{o}-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
51	$2A_1 + (o-\overset{\circ}{o}-*o^3)$	$\mathbb{Z}/2\mathbb{Z}$	S_3	5	$*-(-4)-*$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^{**} $
52	$*-o-\overset{\circ}{o}-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
53	$*-o-\overset{\circ}{o}-o-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
54	$*-\overset{\circ}{o}-o-o-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
55	$o-*-\overset{\circ}{o}-o-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
56	$A_1 + (*-\overset{\circ}{o}-o^4)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
57	$o-o-*-\overset{\circ}{o}-o$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
58	$A_1 + (o-o-\overset{\circ}{o}-o^3)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$o-o-(-4)-\overset{\circ}{o}-o$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
59	$A_3 + (o-o-o-\overset{\circ}{o}-o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$ A_1 + (o-o-o-*o^3) $	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
60	$D_4 + (*-o-\overset{\circ}{o}-o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$A_3 + (*-o^3-*)$	$ U^0 \supseteq \mathbb{C}^2 $
61	$A_3 + (*-o-o-\overset{\circ}{o}-o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$A_1 + (*-o^5-*)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
62	$D_5 + (*-\overset{\circ}{o}-o)$	$\mathbb{Z}/2\mathbb{Z}$	S_3	6	$A_1 + (-4)$	$ U^0 \supseteq \mathbb{C}^2 $
63	$A_2 + (*-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
64	$A_1 + A_2 + (*-o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
65	$A_2 + (*-o-o-o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
66	$A_3 + *$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
67	$A_1 + A_4 + *$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
68	$A_6 + *$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
69	$A_1 + A_4 + (*-o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $

70	$A_2 + A_3 + (*-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
71	$A_6 + (*-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
72	$A_3 + (*-o-o-o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
73	$A_1 + (o-* -o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
74	$2A_1 + (o-* -o-o-o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$2A_1 + ((-4)-o)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
75	$A_1 + (o-* -o^5)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	3	$2A_1 + ((-4)-o-o)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
76	$A_1 + A_2 + (o^2-* -o^2)$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	3	$3A_1 + (-5)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
77	$A_1 + (o-o-* -o^5)$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	5	$3A_1 + ((-5)-o)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
78	$2A_1 + (o^3-* -o^3)$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	5	$4A_1 + (-6)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
79	$A_2 + A_3 + (o-* -o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$A_1 + 2A_2 + (-4)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
80	$A_2 + (o-* -o^5)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	3	$2A_2 + ((-4)-o-o)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
81	$2A_2 + (o-o-* -o-o)$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	3	$3A_2 + (-5)$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
82	$A_1 + A_3 + (o-* -o^3)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$2A_3 + (o-(-4))$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
83	$A_5 + (o-* -o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
84	$A_3 + D_4 + *$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$A_1 + A_3 + 2*$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
85	$A_1 + (*-\overset{\circ}{o}-o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
86	$A_1 + (*-o-\overset{\circ}{o}-o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C}^2 $
87	$2A_1 + (*-\overset{\circ}{o}-o-o-o)$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	$2A_1 + ((-3)-\overset{\circ}{o}-(-3))$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
88	$2A_1 + A_2 + (*-\overset{\circ}{o}-o)$	$\mathbb{Z}/2\mathbb{Z}$	S_3	1	$\bar{U} = \bar{\Sigma}_4$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^{**} $
89	$A_2 + (*-o-o-\overset{\circ}{o}-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
90	$A_3 + (*-o-\overset{\circ}{o}-o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
91	$A_4 + (*-\overset{\circ}{o}-o-o)$	(0)	(0)	1	$\bar{U} = \bar{V}$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
92	$A_1 + A_2 + (*-\overset{\circ}{o}-o^3)$	$\mathbb{Z}/2\mathbb{Z}$	S_3	4	$o-(-\overset{\circ}{4})-o$	$ v^0 \supseteq \mathbb{C} \times \mathbb{C}^{**} $
93	$A_1 + D_4 + (o-\overset{\circ}{*}-o)$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$ D_4 \text{ or } Q_3 $	4	$*$	$ U^0 \supseteq \mathbb{C}^2 $

94	$A + (o - * - o^3 - \overset{\circ}{o} - o)$		$\mathbb{Z}/2\mathbb{Z}$		$\mathbb{Z}/2\mathbb{Z}$		3		$2A_1 + ((-4) - o - \overset{\circ}{o} - o)$	$V^0 \supset \mathbb{C} \times \mathbb{C}^*$
95	$3A_1 + (o - * - o - \overset{\circ}{o} - o)$		$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$		D_4 or Q_3		3		$o - (-4) - o$	$ V^0 \supseteq \mathbb{C} \times \mathbb{C}^{**}$
96	$A_1 + A_3 + (o - * - \overset{\circ}{o} - o)$		$\mathbb{Z}/4\mathbb{Z}$		$\mathbb{Z}/4\mathbb{Z}$		4		$4A_1 + ((-4) - (-4))$	$ V^0 \supseteq \mathbb{C} \times \mathbb{C}^* $
97	$D_7^+ *$		(0)		(0)		1		$\bar{U} = \bar{V}$	$ V^0 \supset \mathbb{C} \times \mathbb{C}^* $

Let $f: U \rightarrow \bar{U}$ be the minimal resolution of singularities on the quasi-universal covering $\bar{U}$ of a dP3 surface $\bar{V}$. The singularity of $\bar{V}$ (resp. $\bar{U}$) is described in terms of the Dynkin graph of $D := g^{-1}(\text{Sing } \bar{V}) \subseteq V$ (resp. $B := f^{-1}(\text{Sing } \bar{U}) \subseteq U$).

V^0, U^0 : stand for $V-D$ and $U-B$, respectively;

$\mathbb{C}^*, \mathbb{C}^{**}, \mathbb{C}^2-P$: stand for $\mathbb{C} - \{0\}$, $\mathbb{C} - \text{two distinct points}$, and $\mathbb{C}^2 - \text{one point } P$, respectively;

$\bar{\Sigma}_n (n \geq 2)$: the surface obtained by contracting the minimal section on a Hirzebruch surface Σ_n of degree n ;

$*, o, (-n)$: stands for the unique (-3) curve on V , a (-2) curve on V , and a $(-n)$ curve on U , respectively;

$*-o^3-*, \text{etc.}$: stands for $*-o-o-o-*$, etc., respectively;

$(*-o) + A_m + D_n$: disjoint union of $*-o$ and Dynkin types A_m and D_n ;

We employ the following notations for finite groups.

D_4 : the dihedral group of order 8;

Q_3 : the quaternion group of order 8;

S_3 : the third symmetric group of order 6.

In No.22, $\pi_1 = \langle x, y, z \mid x^3=y^3=z^2=1, xy=yx, yz=zy, xz=zx^2 \rangle$.

In No.26, $\pi_1 = \langle a, b \mid a^3=b^4=1, ab=ba^2 \rangle$.

Though the surfaces $\bar{V}^{(a)}$ and $\bar{V}^{(b)}$ of No.18a and No.18b (resp. No.15a and No.15b) have singularities of the same type, they are not isomorphic to each other.

Table 2

Dynkin type of S	$\pi_1(S^\circ)$	$Cl(S)/Pic(S)$	$\rho(\tilde{S})$	Sing. of $\tilde{S}$
A_1	0	$\mathbb{Z}/2\mathbb{Z}$	1	$\tilde{S} = S$
$A_1 + A_2$	0	$\mathbb{Z}/6\mathbb{Z}$	1	$\tilde{S} = S$
A_4	0	$\mathbb{Z}/5\mathbb{Z}$	1	$\tilde{S} = S$
$2A_1 + A_3$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$	1	$\tilde{S} = S(A_1)$
D_5	0	$\mathbb{Z}/4\mathbb{Z}$	1	$\tilde{S} = S$
$A_1 + A_5$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/6\mathbb{Z}$	2	A_2
$3A_2$	$\mathbb{Z}/3\mathbb{Z}$	$(\mathbb{Z}/3\mathbb{Z})^{\oplus 2}$	1	$\tilde{S} = \mathbb{P}^2$
E_6	0	$\mathbb{Z}/3\mathbb{Z}$	1	$\tilde{S} = S$
$3A_1 + D_4$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 3}$	1	$\tilde{S} = S(A_1)$
A_7	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/4\mathbb{Z}$	3	A_3
$A_1 + D_6$	$\mathbb{Z}/2\mathbb{Z}$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	2	D_4

Dynkin type of S	$\pi_1(S^\circ)$	$\text{Cl}(S)/\text{Pic}(S)$	$\rho(\tilde{S})$	Sing. of $\tilde{S}$
E_7	0	$\mathbb{Z}/2\mathbb{Z}$	1	$\tilde{S} = S$
$A_1 + 2A_3$	$\mathbb{Z}/4\mathbb{Z}$	$\mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$	2	$\tilde{S} = \mathbb{P}^1 \times \mathbb{P}^1$
$A_2 + A_5$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/6\mathbb{Z}$	3	A_1
D_8	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	3	D_5
$2A_1 + D_6$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	3	$\tilde{S} = \widetilde{S(A_7)}$
E_8	0	0	1	$\tilde{S} = S$
$A_1 + E_7$	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	2	E_6
$A_1 + A_7$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$\mathbb{Z}/4\mathbb{Z}$	5	A_1
$2A_4$	$\mathbb{Z}/5\mathbb{Z}$	$\mathbb{Z}/5\mathbb{Z}$	5	a smooth del Pezzo surface of degree 5
A_8	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	5	A_2
$A_1 + A_2 + A_5$	$\mathbb{Z}/6\mathbb{Z}$	$\mathbb{Z}/6\mathbb{Z}$	4	a smooth del Pezzo surface of degree 6

Dynkin type of S	$\pi_1(S^\circ)$	$\text{Cl}(S)/\text{Pic}(S)$	$\rho(\tilde{S})$	Sing. of $\tilde{S}$
$A_2 + E_6$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	3	D_4
$A_3 + D_5$	$\mathbb{Z}/4\mathbb{Z}$	$\mathbb{Z}/4\mathbb{Z}$	4	A_2
$4A_2$	$(\mathbb{Z}/3\mathbb{Z})^{\oplus 2}$	$(\mathbb{Z}/3\mathbb{Z})^{\oplus 2}$	1	$\tilde{S} = \mathbb{P}^2$
$2A_1 + 2A_3$	$\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$	2	$\tilde{S} = \widetilde{S(A_1 + 2A_3)}$
$2D_4$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	$(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$	4	$2A_1$

References

- [11] L.Brenton et al, Graph theoretic techniques in algebraic geometry I : The extended Dynkin diagram $\bar{E}_8$ and minimal singular compactifications of $\mathbb{C}^2$, Recent Developments in Several Complex Variables, 47 - 63, Princeton University Press, 1981.
- [12] M. Demazure, Surfaces de del Pezzo, Lecture Notes in Math. No.777, Berlin-Heidelberg-New York: Springer, 1980,
- [13] M. Furushima, Singular del Pezzo surfaces and analytic compactifications of 3-dimensional complex affine space $\mathbb{C}^3$, Nagoya Math. J., 104(1986), 1 - 28.
- [14] F. Hidaka and K. Watanabe, Normal Gorenstein surfaces with ample anti-canonical divisor, Tokyo J. Math. 4(1981), 319 - 330.
- [15] Y. Kawamata, On the classification of non-complete algebraic surfaces, Proc. Copenhagen Summer Meeting in Algebraic Geometry, Lecture Notes in Math. No.732, 215-232, Berlin - Heidelberg - New York: Springer, 1978.
- [16] M. Miyanishi - S. Tsunoda, Non-complete algebraic surfaces with logarithmic Kodaira dimension $-\infty$ and with non-connected boundaries at infinity, Japan. J. Math. 10(1984), 195-242.
- [17] ——— - ———, Logarithmic del Pezzo surfaces of rank one with non-contractible boundaries, Japan. J. Math., 10(1984), 271-319.
- [18] M. Miyanishi - D.Q. Zhang, Gorenstein log del Pezzo

surfaces of rank one, to appear in J. Algebra.

- [9] T. Urabe, On singularities on degenerate
del Pezzo surfaces of degree 1,2,
Proc. Symp. Pure Math. 40(1983), 587 - 591.
- [10] D.Q.Zhang, On Iitaka surfaces, Osaka. J. Math.,
24(1987), 417-460.
- [11] D.Q.Zhang, Logarithmic del Pezzo surfaces of rank one
with contractible boundaries, Osaka. J. Math. 25(1988).
- [12] D. Q. Zhang, The classification and the quasi-universal
covering of log del Pezzo surfaces with only rational
double or rational triple singular points. Preprint.