

A GENERALIZED EXTENSION OF AVERAGED MAPPINGS DEFINED IN A HILBERT SPACE TO BANACH SPACES

HIROKO MANAKA

ABSTRACT. It is well known that there is an averaged mapping as one of useful nonlinear mappings for many iterative algorithms. In a Hilbert space we can obtain a theorem that an averaged mapping is determined by some inequality, since we can use the important tool of relation between a norm and an inner product. However, this determinism can not have a power in Banach spaces. In this paper, we introduce a generalized extension of an averaged mapping determined in a Hilbert space to Banach spaces.

1. AN AVERAGED MAPPING AND ITS PROPERTIES

At first we give an introduction of an averaged mapping defined in a Hilbert space.

Definition 1.1. Let H be a Hilbert space, and let $\langle \cdot, \cdot \rangle$ be an inner product, and let $\|\cdot\|$ be a norm on H . Let T be a mapping from H to H . T is called an averaged mapping if there exists $\alpha \in [0, 1]$ and a nonexpansive mapping $N : H \rightarrow H$ such that

$$T = (1 - \alpha)I + \alpha N,$$

where I is an identity mapping. More explicitly, T is called an α -averaged mapping, or an averaged mapping with α .

We obtained some properties of averaged mappings in a Hilbert space as follows([1]).

Proposition 1.1. *In a Hilbert space H , the following hold:*

- (a) *A mapping $T : H \rightarrow H$ is firmly nonexpansive if and only if T is a $(\frac{1}{2})$ -averaged mapping.*
- (b) *Let T_1 and T_2 be firmly nonexpansive on H . Then, T_2T_1 is $(\frac{2}{3})$ -averaged.*
- (c) *Let T_i be α_i -averaged for every $i \in I = \{1, 2, 3, \cdot, \cdot, m\}$. Then, $T_mT_{m-1} \cdot \cdot T_2T_1$ is α -averaged with*

$$\alpha = \frac{\sum_{i \in I} (\frac{\alpha_i}{1 - \alpha_i})}{1 + \sum_{i \in I} (\frac{\alpha_i}{1 - \alpha_i})}.$$

In a Hilbert space, we obtain the following theorem by using a good relation between a norm and an inner product (cf. [1]).

Date: 2025.4.12.

2000 Mathematics Subject Classification. Primary 49J40, Secondary 47J20.

Key words and phrases. averaged mapping, V-strongly nonexpansive mapping, strong convergence theorem, Bregman distance, Hilbert space, Banach space.

Theorem 1.1. *Let $T : H \rightarrow H$, and suppose $\alpha \in (0, 1)$. Then (a) and (b) are equivalence.*

(a) *T is an α -averaged mapping.*

(b) *T satisfies the following inequality (1) for all $x, y \in E$*

$$(1) \quad \|Tx - Ty\|^2 \leq \|x - y\|^2 - \left(\frac{1 - \alpha}{\alpha}\right) \|(I - T)x - (I - T)y\|^2.$$

2. BREGMAN DISTANCE IN BANACH SPACES

We also introduce Bregman distance defined in Banach spaces (cf. [6]).

Definition 2.1. Let E be a Banach space. Let $f : E \rightarrow \mathbb{R}$ a *Gâteaux* differentiable and convex function and $f'(x)$ stands for the derivative of f at $x \in E$.

Then, Bregman distance D_f is defined and denoted as follows:

$$D_f(x, y) = f(x) - f(y) - \langle x - y, f'(y) \rangle,$$

for all $x, y \in E$, where $\langle \cdot, \cdot \rangle$ stands for the duality pair

Moreover, we give a definition of $V(\cdot, \cdot)$, as a special version of Bregman distance with $f = \|\cdot\|^2$.

Definition 2.2. Let E be a smooth Banach space with a norm $\|\cdot\|$. We denote a function V on a $E \times E$ as follows:

$$V(x, y) = \|x\|^2 + \|y\|^2 - 2 \langle x, Jy \rangle$$

for all $x, y \in E$, where $\langle \cdot, \cdot \rangle$ stands for the duality pair and J is the normalized duality mapping, i.e.

$$Jx = \{x^* \in E^* : \langle x, x^* \rangle = \|x\|^2, \quad \|x^*\| = \|x\|\}.$$

It is well known that the value of the normalized duality mapping J is singleton in a smooth Banach space. It is trivial $V(x, y) \geq 0$ for all $x, y \in E$. In a Hilbert space H , as a special case of a Banach space, it is shown that $V(x, y) = \|x - y\|^2$ for all $x, y \in H$.

Lemma 2.1. (cf.[10]) *The V has the following properties.*

(i) *For all $x, y, z \in E$,*

$$2 \langle x - y, Jy - Jz \rangle \leq V(x, y) + V(x, z).$$

(ii) *If a sequence $\{x_n\} \subset E$ satisfies $\lim_{n \rightarrow \infty} V(x_n, p) < \infty$ for some $p \in E$, then $\{x_n\}$ is bounded.*

(iii) *Let E be a smooth and uniformly convex Banach space. There exists a continuous, strictly increasing and convex function $g : [0, \infty) \rightarrow [0, \infty)$ such that $g(0) = 0$ and for each real number $r > 0$ and all $x, y \in \{z \in E : \|z\| \leq r\}$*

$$0 \leq g(\|x - y\|) \leq V(x, y).$$

Remark 1. This $V(\cdot, \cdot)$ does not mean a distance, since the so-called triangle inequality is not satisfied. In a smooth and uniformly convex Banach spaces, $V(x, y) = 0$ implies $x = y$ by the above lemma 2.1 (iii).

3. A NONLINEAR MAPPING DETERMINED BY BREGMAN DISTANCE V

We introduce a definition of a special mapping in Banach spaces and named V -strongly nonexpansive mapping as follows ([10]):

Definition 3.1. Let E be a smooth Banach space. A mapping $T : E \rightarrow E$ is said to be V -strongly nonexpansive if there exists a constant $\lambda > 0$ such that

$$(2) \quad V(Tx, Ty) \leq V(x, y) - \lambda V((I - T)x, (I - T)y),$$

for all $x, y \in E$. More explicitly, T is called a V -strongly nonexpansive mapping with λ .

For the sake of showing the relation among other nonlinear mappings, we give definitions of them in a Banach space E and a Hilbert space H .

Definition 3.2. (1) $T : E \rightarrow E$ is said to be firmly nonexpansive if

$$\|Tx - Ty\|^2 \leq \langle x - y, J(Tx - Ty) \rangle$$

for all $x, y \in E$.

(2) $T : H \rightarrow H$ is said to be β -inverse strongly monotone if

$$\beta \|Tx - Ty\|^2 \leq \langle x - y, Tx - Ty \rangle$$

for all $x, y \in H$.

(3) $T : H \rightarrow H$ is said to be strongly nonexpansive if T is nonexpansive with some fixed point and if it holds that

$$(x_n - y_n) - (Tx - Ty) \rightarrow 0$$

for sequences $\{x_n\}$ and $\{y_n\}$ such that $\{x_n - y_n\}$ is bounded and $\lim_{n \rightarrow \infty} (\|x_n - y_n\| - \|Tx_n - Ty_n\|) = 0$.

We give a proposition with respect to the relation among the above nonlinear mappings.

Proposition 3.1. ([10]) *In a Hilbert space,*

- (a) *A firmly nonexpansive mapping is V -strongly nonexpansive with $\lambda = 1$.*
- (b) *If T is a β -inverse strongly monotone mapping for some $\beta > \frac{1}{2}$, then $(I - T)$ is V -strongly nonexpansive with $\lambda = (2\beta - 1)$.*
- (c) *A V -strongly nonexpansive mapping with a nonempty subset of asymptotically fixed points is strongly nonexpansive.*

Proposition 3.2. ([10], [12]) *In smooth Banach spaces,*

- (a) *For any $c \in (-1, 1)$, cI is V -strongly nonexpansive with any $\lambda \in (0, \frac{1+c}{1-c})$. Especially, for $c = 1$, I is V -strongly nonexpansive with any $\lambda > 0$.*
- (b) *If T is V -strongly nonexpansive with $\lambda > 0$, then for any $\alpha \in [-1, 1]$ with $\alpha \neq 0$, αT is V -strongly nonexpansive with $\alpha^2 \lambda$.*
- (c) *If T is V -strongly nonexpansive with $\lambda \geq 1$, then $(I - T)$ is also V -strongly nonexpansive with λ^{-1} .*
- (d) *Suppose that T is V -strongly nonexpansive with $\lambda > 0$, and that $\alpha \in [-1, 1]$ satisfies $\alpha^2 \lambda \geq 1$. Let $T_\alpha = (I - \alpha T)$. Then T_α is V -strongly nonexpansive with $(\alpha^2 \lambda)^{-1}$. Moreover,*

$$V(T_\alpha x, T_\alpha y) \leq V(x, y) - \lambda^{-1} V(Tx, Ty).$$

Remark 2. The V -strongly nonexpansive mapping is not necessarily nonexpansive. We have an example of a V -strongly nonexpansive mapping which has a nonempty subset of fixed points and which is not even quasi-nonexpansive in some Banach space ([12]).

4. COMPARISON BETWEEN THE AVERAGED MAPPING AND THE V -STRONGLY NONEXPANSIVE MAPPING IN BANACH SPACES

We recall Theorem 1.1 that a mapping T is α -averaged mapping if and only if T satisfies the following inequality for all $x, y \in E$

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 - \left(\frac{1-\alpha}{\alpha}\right) \|(I-T)x - (I-T)y\|^2,$$

where $\alpha \in (0, 1)$. Exchanging $\frac{1-\alpha}{\alpha}$ for λ , we get

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 - \lambda \|(I-T)x - (I-T)y\|^2,$$

as $\lambda > 0$. In a Hilbert space, this means that T is an averaged mapping if and only if T is a V -strongly nonexpansive mapping, since the following correspondance holds:

$$\lambda = \frac{1-\alpha}{\alpha} \iff \alpha = \frac{1}{\lambda+1}.$$

In a Banach space, we could also give a definition of an averaged mapping as the same as one in a Hilbert space;

Definition 4.1. Let E be a Banach space with a norm $\|\cdot\|$. Let T be a mapping from E to E . T is called an averaged mapping if there exists $\alpha \in (0, 1)$ and a nonexpansive mapping $N : E \rightarrow E$ such that

$$T = (1-\alpha)I + \alpha N.$$

Then T is a nonexpansive mapping since

$$\begin{aligned} \|Tx - Ty\| &= \|(1-\alpha)x + \alpha Nx - (1-\alpha)y - \alpha Ny\| \\ &= \|(1-\alpha)(x-y) + \alpha(Nx - Ny)\| \\ &\leq (1-\alpha) \|x-y\| + \alpha \|Nx - Ny\| \\ &\leq (1-\alpha) \|x-y\| + \alpha \|x-y\| = \|x-y\|. \end{aligned}$$

This definition means that an averaged mapping is a nonexpansive mapping in an arbitrary Banach space. However, with respect to a V -strongly nonexpansive mapping, we have obtained an example of V -strongly nonexpansive mappings which has a nonempty set of fixed points and which is not even quasi nonexpansive in a Banach space l^p with $p = \frac{3}{2}$ (see [12]).

The properties of averaged mappings are useful for the convergence theorems for fixed points with an iterative scheme, especially in a Hilbert space. However in Banach spaces, instead of those V -strongly nonexpansive mappings ought to play important role.

At last we should introduce a theorem with respect to this nonlinear mapping in a Banach space.

Theorem 4.1. ([10]) *Let E be a reflexive, smooth and strictly convex Banach space. Let C be a nonempty, closed, and convex subset of E , and $R_C : E \rightarrow C$ sunny and generalized nonexpansive retraction. Let $B : E^* \rightarrow 2^E$ be a maximal monotone operator and let $J_{r_n} = (I + r_n B J)^{-1}$ be a generalized resolvent of B for a sequence $\{r_n\}$, where the duality mapping J of E is weakly sequential continuous, and $\{r_n\} \subset (0, \infty)$. Suppose $A : C \rightarrow E$ is a V strongly nonexpansive mapping with $\lambda \geq 1$ such that $C_0 = A^{-1} \cap (B J)^{-1}(0) \neq \emptyset$. For an $\alpha \in [-1, 1]$ such that $\alpha^2 \lambda \geq 1$, let a sequence $\{x_n\} \subset C$ be defined as follows: For any $x = x_1 \in C$ and every $n \in \mathbb{N}$,*

$$\begin{aligned} y_n &= R_C(I - \alpha A)x_n, \\ x_{n+1} &= R_C(\beta_n x + (1 - \beta_n)J_{r_n}y_n), \end{aligned}$$

where $\{\beta_n\} \subset [0, 1]$ and $\{r_n\} \subset (0, \infty)$ satisfy that

$$\sum_{n \geq 1} \beta_n < \infty, \quad \liminf_{n \rightarrow \infty} r_n > 0.$$

Then there exists an element $u \in C_0$ such that

$$x_n \rightharpoonup u, \quad R_{C_0}(x_n) \rightarrow u.$$

Acknowledgement 1. *This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.*

REFERENCES

- [1] Heinz H. Bauschke, and Walaa M. Mouri, *An Introduction to Convexity, Optimization, and Algorithms*, MOS-SIAM Series on Optimization, SIAM, (2024).
- [2] Ya, Alber and S. Guerre-Delabriere, *Principles of weakly contractive maps in Hilbert space*, Operator Theory, Advances and Applications **98** (1997), 7-21.
- [3] H. Bauschke, *Fenchel duality, Fitzpatrick functions and the extension of firmly nonexpansive mapping*, Proceeding of the American Mathematical Society, **135** (2007), 135-139.
- [4] V. Barbu, *Nonlinear semigroups and differential equations in Banach spaces*, Noordhoff International Publishing, (1976).
- [5] R. E. Bruck and S. Reich, *Nonexpansive projections and resolvents of accretive operators in Banach spaces*, Houston, J. Math. **3** (1977), 459-470.
- [6] D. Butnariu and E. Resmerita, *Bregman distances, totally convex function and a method for solving operator equations in Banach spaces*, Abstract and Applied Analysis, (2006), Art. ID 84919, 39pp.
- [7] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, Cambridge Studies in Advanced Mathematics, **28**, Cambridge University Press, Cambridge, 1990.
- [8] T. Ibaraki and W. Takahashi, *A new projection and convergence theorems for the projections in Banach spaces*, J. Approximation Theory, **149**(2007), 1-14.
- [9] S. Kamiruma and W. Takahashi, *Strong Convergence of a Proximal-Type Algorithm in a Banach space*, SIAM Journal on Optimization, **13**(2002), 938-945.
- [10] H. Manaka, *Convergence theorems for a maximal monotone operator and a V -strongly nonexpansive mapping in a Banach space*, Abstract and Applied Analysis, **2010**, Article ID 189814, (20 pages).
- [11] H. Manaka, *Convergence Theorems for Fixed Points with Iterative Methods in Banach spaces*, Thesis (Yokohama National University), 2011.
- [12] Hiroko Manaka, *Fixed point theorems for an elastic nonlinear mapping in Banach spaces*, Abstract and Applied Analysis, Article ID 760671,(2015).

(Hiroko Manaka) COLLEGE OF SCIENCE AND TECHNOLOGY, NIHON UNIVERSITY, 7-24-1, NARASHINO-DAI, FUNABASHI-SHI, CHIBA, 274-8501, JAPAN

Email address: manaka.hiroko@nihon-u.ac.jp