

The eigenvalues of the principal minor of a Hermitian random matrix

By

Takahiro HASEBE*

Abstract

This is a summary of the paper [8]. We study the difference between the empirical eigenvalue distribution of a unitarily invariant Hermitian random matrix and that of its principal minor. Under some technical assumptions, we prove that the difference can be described by the Markov–Krein correspondence as the size of the matrix goes to infinity.

§ 1. Introduction

Let X_N be a hermitian random matrix of size N whose distribution on the matrix space $M_N(\mathbb{C})$ is invariant under conjugacy by any unitary matrix. Let $\Lambda_N = (\lambda_1^{(N)} \leq \dots \leq \lambda_N^{(N)})$ be its eigenvalues. Then a diagonalization $X_N = U_N D_N U_N^*$ exists, where $D_N = \text{diag}(\lambda_1^{(N)}, \lambda_2^{(N)}, \dots, \lambda_N^{(N)})$ and U_N is a Haar unitary random matrix of size N and independent of D_N (see [6, Proposition 6.1]). We denote by $\mathfrak{m}_N$ the *empirical eigenvalue distribution* of X_N :

$$\mathfrak{m}_N := \frac{1}{N} \sum_{i=1}^N \delta_{\lambda_i^{(N)}}.$$

Given a Schwartz distribution ζ on $\mathbb{R}$, let $M_k(\zeta)$ denote the evaluation of ζ by the test function x^k and call it the k -th moment of ζ , as long as it is well defined. If ζ is a probability measure, then $M_k(\zeta)$ is the usual k -th moment.

Received December 28, 2023. Revised March 5, 2024.

2020 Mathematics Subject Classification(s): 46L54, 60B20

This work is supported by JSPS kakenhi 19K14546 and 18H01115 and JSPS Open Partnership Joint Research Projects grant no. JPJSBP120209921 and Bilateral Joint Research Projects (JSPS-MEAE-MESRI, grant no. JPJSBP120203202).

*Department of Mathematics, Hokkaido University, North 10 West 8, Kita-Ku, Sapporo 060-0810, Japan.

e-mail: `thasebe@math.sci.hokudai.ac.jp`

Assumption 1.1. The following are assumed throughout this paper.

$$(A1) \quad \sup_{N \geq 1} \mathbb{E}[M_k(\mathbf{m}_N)] < \infty, \quad k \in 2\mathbb{N}.$$

(A2) There exists a (non-random) probability distribution $\mathbf{m}$ on $\mathbb{R}$ having finite moments of all orders such that $\mathbf{m}_N$ converges in moments in probability to $\mathbf{m}$:

$$\lim_{N \rightarrow \infty} \mathbb{P}[|M_k(\mathbf{m}_N) - M_k(\mathbf{m})| \geq \epsilon] = 0, \quad k \in \mathbb{N}, \quad \epsilon > 0.$$

Let $\tilde{X}_N$ be the principal minor of X_N made by removing the last row and column of X_N . Cauchy's interlacing law says that the eigenvalues $\tilde{\Lambda}_N = (\tilde{\lambda}_1^{(N)} \leq \dots \leq \tilde{\lambda}_{N-1}^{(N)})$ of $\tilde{X}_N$ interlace with Λ_N (see [16, Exercise 1.3.14]):

$$\lambda_1^{(N)} \leq \tilde{\lambda}_1^{(N)} \leq \lambda_2^{(N)} \leq \tilde{\lambda}_2^{(N)} \leq \dots \leq \lambda_{N-1}^{(N)} \leq \tilde{\lambda}_{N-1}^{(N)} \leq \lambda_N^{(N)}.$$

It is not hard to see under Assumption 1.1 that the empirical eigenvalue distribution $\tilde{\mathbf{m}}_N$ of $\tilde{X}_N$ also converges in moments in probability to the same limit $\mathbf{m}$. Our main purpose is to investigate the difference of $\mathbf{m}_N$ and $\tilde{\mathbf{m}}_N$. Let us investigate the rescaled difference $N(\mathbf{m}_N - \tilde{\mathbf{m}}_N)$, which can be calculated as

$$N(\mathbf{m}_N - \tilde{\mathbf{m}}_N) = \sum_{i=1}^N \delta_{\lambda_i^{(N)}} - \frac{N}{N-1} \sum_{j=1}^{N-1} \delta_{\tilde{\lambda}_j^{(N)}} = \underbrace{\sum_{i=1}^N \delta_{\lambda_i^{(N)}} - \sum_{j=1}^{N-1} \delta_{\tilde{\lambda}_j^{(N)}}}_{=: \hat{\tau}_N} - \tilde{\mathbf{m}}_N.$$

Because $\tilde{\mathbf{m}}_N$ converges to $\mathbf{m}$, the problem is the convergence of the signed measure $\hat{\tau}_N$.

To describe the convergence of $\hat{\tau}_N$, recall the notions of *free cumulants* and *Markov-Krein correspondence*. The free cumulants $R_n(\mathbf{m}), n \in \mathbb{N}$, of $\mathbf{m}$ are the real numbers defined by the recursive formula

$$(1.1) \quad M_k(\mathbf{m}) = \sum_{\rho \in \text{NC}(k)} R_\rho(\mathbf{m}),$$

where $\text{NC}(k)$ is the set of non-crossing partitions on the set $\{1, 2, \dots, k\}$ and $R_\rho(\mathbf{m}) := \prod_{B \in \rho} R_{|B|}(\mathbf{m})$. Free cumulants are helpful for computing free convolutions and are well studied in free probability. For further details, the reader is referred to [15]. On the other hand, let τ be the Schwartz distribution determined by formula

$$(1.2) \quad \int_{\mathbb{R}} \frac{1}{1 - zx} d\mathbf{m}(x) = \exp \left[\int_{\mathbb{R}} \log \frac{1}{1 - zx} d\tau(x) \right], \quad z \in \mathbb{C} \setminus \mathbb{R}.$$

The correspondence between $\mathbf{m}$ and τ gives a bijection between the set of probability measures on $\mathbb{R}$ and the set of certain Schwartz distributions, and is called the *Markov-Krein correspondence*; see [12] for further details.

Here are three remarks on the Markov–Krein correspondence. Firstly, in many examples, τ is a signed measure, and in such a case it is called the Rayleigh measure of $\mathbf{m}$. In general, τ is the Schwartz-distributional derivative of a so-called Rayleigh function.

Secondly, because we assumed that $\mathbf{m}$ has finite moments of all orders, so is τ (see [1, Theorem A (d)] and [12, Section 3.4]). The relationship between the two kinds of moments $M_k(\mathbf{m})$ and $M_k(\tau)$, $k \in \mathbb{N}$, is exactly the one between the complete symmetric functions and Newton’s power sums, i.e.,

$$(1.3) \quad M_k(\tau) = k M_k(\mathbf{m}) - \sum_{r=1}^{k-1} M_r(\tau) M_{k-r}(\mathbf{m}), \quad k \in \mathbb{N}.$$

Thirdly, the Markov–Krein correspondence appears in different contexts to describe interlacing sequences: limit shapes of large random Young diagrams [13, 17, 3, 4]; roots of two consecutive orthogonal polynomials of large degrees [11]; eigenvalues of large random matrices and of their principal minor [11, 5] (the present paper deals with this category). There are also situations where the distribution τ above appears as a probability measure: Poisson–Dirichlet processes (see [12, Section 4.1] and references therein); self-decomposable distributions for monotone convolution [7]; Harish-Chandra–Izykson–Zuber integral of rank one at a high temperature regime [14]. The reason why the same correspondence appears in different contexts seems still unclear.

§ 2. The main result

The main result of [8] is as follows.

Theorem 2.1. *Let $\mathbf{m}_N, \mathbf{m}, \hat{\tau}_N, \tau$ be as in Section 1. Then $M_k(\hat{\tau}_N)$ converges to $M_k(\tau)$ in L^2 as $N \rightarrow \infty$ for every $k \in \mathbb{N}$. In particular, $\hat{\tau}_N$ converges to τ in moments in probability.*

This result is known as a folklore theorem and is announced in [10] as a conjecture. Similar results are previously obtained in [11] for randomly rotated real Wigner matrices and then in [5] for Wigner and Wishart matrices (without random rotation).

The proof is based on a free probability technique and is sketched below; for further details see [8]. We first compute the expectation of $M_k(\hat{\tau}_N)$, which is given by

$$(2.1) \quad \mathbb{E}[M_k(\hat{\tau}_N)] = \mathbb{E} \circ \text{Tr}[D_N^k] - \mathbb{E} \circ \text{Tr}[(D_N U_N P_N U_N^*)^k],$$

where $P_N = \text{diag}(1, 1, \dots, 1, 0)$. Weingarten calculus fits this problem and yields the asymptotic expansion of $\mathbb{E} \circ \text{Tr}[(D_N U_N P_N U_N^*)^k]$ in variable N^{-1} . The first order term

of the expansion is $\mathbb{E} \circ \text{Tr}[D_N^k]$ that is of order $O(N)$ and exactly cancels the first term of (2.1). The second order term, which is the crucial part, turns out to be

$$(2.2) \quad - \sum_{\rho \in \text{NC}(k)} (k + 1 - |\rho|) R_\rho(\mathbf{m}),$$

where $|\rho|$ is the number of blocks of ρ . To prove the main theorem, expression (2.2) should equal $-M_k(\tau)$, and indeed it is. This equality can be easily proved by combining combinatorial formulas for complete symmetric functions. The higher order terms will vanish in the limit and are unnecessary for the proof. Considering the above, we obtain $\mathbb{E}[M_k(\widehat{\tau}_N)] = M_k(\tau) + o(1)$ as $N \rightarrow \infty$. To complete the proof of L^2 convergence $M_k(\widehat{\tau}_N) \rightarrow M_k(\tau)$, we also need to compute and estimate $\mathbb{E}[M_k(\widehat{\tau}_N)^2]$. Although the computation is more involved, Weingarten calculus also works for this case.

As a final remark, the recent paper [9] investigated equation (2.1) from the viewpoint of noncommutative independence. As a result, the equality of (2.2) and $-M_k(\tau)$ was generalized to a multivariate situation, see [9, Theorem 6.1].

Acknowledgement

This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

References

- [1] Aronszajn, N. and Donoghue, W. F., On exponential representations of analytic functions in the upper half-plane with positive imaginary part, *J. Analyse Math.* Vol. V (1956-57), 321–388.
- [2] Baryshnikov, Y., GUEs and queues, *Probab. Theory Relat. Fields* 119 (2001), 256–274.
- [3] Biane, P., Representations of symmetric groups and free probability, *Advances in Mathematics* 138 (1998), 126–181.
- [4] Biane, P., Approximate factorization and concentration for characters of symmetric groups, *International Mathematics Research Notices* 2001, No. 4 (2001), 179–192.
- [5] Bufetov, A., Kerov’s interlacing sequences and random matrices, *J. Math. Phys.* 54, (2013), 113302, 10pp.
- [6] Collins, B. and Male, C., The strong asymptotic freeness of Haar and deterministic matrices, *Annales Scientifiques de l’École Normale Supérieure* 47 (2014), 147–163.
- [7] Franz, U., Hasebe, T. and Schleissinger, S., Monotone increment processes, classical Markov processes and Loewner chains, *Dissertationes Mathematicae* 552 (2020), 1–119.
- [8] Fujie, K. and Hasebe, T., The spectra of principal submatrices in rotationally invariant hermitian random matrices and the Markov–Krein correspondence, *ALEA Lat. Am. J. Probab. Math. Stat.* 19 (2022), 109–123.
- [9] Fujie, K. and Hasebe, T., Free probability of type B prime. *ArXiv:2310.14582*
- [10] Goel, G. and Yao, A., A quantized analogue of the Markov–Krein correspondence, *International Mathematics Research Notices* 2023, Issue 6 (2023), 4805–4838.
- [11] Kerov, S., Asymptotics of the separation of roots of orthogonal polynomials, *St. Petersburg Mathematical Journal* 5 (1994), 925–941.
- [12] Kerov, S., Interlacing measures, in: *Kirillov’s seminar on representation theory*, 35–83, *Amer. Math. Soc. Transl. Ser. 2, Volume 181* (G.I. Olshanski ed.), Amer. Math. Soc., Providence, RI, 1998.
- [13] Logan, F. and Shepp, L. A., A variational problem for random Young tableaux, *Advances in Mathematics* 26 (1977), 206–222.
- [14] Mergny, P. and Potters, M., Rank one HCIZ at high temperature: interpolating between classical and free convolutions, *SciPost Phys.* 12 (2022), 022.
- [15] Nica, A. and Speicher, R., *Lectures on the combinatorics of free probability*. London Mathematical Society Lecture Note Series, 335. Cambridge University Press, Cambridge, 2006. xvi+417 pp.
- [16] Tao, T., *Topics in Random Matrix Theory*, American Mathematical Society, 2012.
- [17] Vershik, A. M. and Kerov, S., Asymptotics of the Plancherel measure of the symmetric group and the limiting form of Young tableaux, *Doklady AN SSSR* 233 (1977), no. 6, 1024–1027; English translation: *Soviet Mathematics Doklady* 18 (1977), 527–531.