

A NOTE ON BMO-MARTINGALES

By N.Kazamaki and T.Sekiguchi

(風巻 紀彦、関口 健)

Toyama University

(富山大・理)

Let $(\Omega, \mathcal{F}, P; (\mathcal{F}_t))$ be a probability system where the filtration $(\mathcal{F}_t)$ satisfies the usual conditions. We now consider two important subclasses of BMO, namely the class L^∞ of all bounded martingales and the class H^∞ of all martingales with bounded quadratic variation. There is not always an inclusion relation between these classes. In fact, if $B=(B_t)$ is a one dimensional Brownian motion starting at 0, then $(B_{t \wedge 1}) \in H^\infty \setminus L^\infty$ clearly. On the other hand, the process B stopped at τ , where $\tau = \min\{t: |B_t|=1\}$, belongs to $L^\infty \setminus H^\infty$ (see [2]).

Our aim is to prove the following.

THEOREM. Suppose the sample continuity of any martingale adapted to $(\mathcal{F}_t)$. Then the following properties are equivalent:

- (a) $BMO = L^\infty$.
- (b) $BMO = H^\infty$.
- (c) $\mathcal{F}_t = \mathcal{F}_0$ for every t .

Furthermore, excepting such a trivial case, $H^\infty \cup L^\infty$ is not dense in BMO.

Recall that BMO is the class of those uniformly integrable martingales M which satisfy $\|M\|_{BMO} = \sup_T \|E[(M_\infty - M_{T-})^2 | \mathcal{F}_T]^{1/2}\|_\infty < \infty$

where the supremum is taken over all stopping times T . It is well-known that BMO is Banach space with the norm $\|M\|_{\text{BMO}}$.

PROOF. Firstly, we shall establish the implication (a) $\rightarrow$ (c). In order to see (c), it suffices to prove that for any martingale M , almost all sample functions of M are constant, and by using the usual stopping argument we may assume $M \in \text{BMO}$. Suppose now $\text{BMO} = L^\infty$. Then the two norms $\|M\|_{\text{BMO}}$ and $\|M\|_\infty$ on BMO are equivalent by the Closed Graph Theorem. So there exists a constant $C > 0$, depending only on M , such that $\|K \cdot M\|_\infty < C$ for any predictable process $K = (K_t, \mathcal{F}_t)$ with $|K| \leq 1$. Here $K \cdot M$ denotes the stochastic integral of K relative to M . We show below that the negation of (c) causes a contradiction. If we deny (c), then there exist $t > 0$ and a partition $\Delta: 0 = t(0) < t(1) < \dots < t(n) = t$ of $[0, t]$ such that $P(A) > 0$ where $A = \{\sum_{j=1}^n |M_{t(j)} - M_{t(j-1)}| > 2C\}$. Let now

$$B_{j, \varepsilon(j)} = \begin{cases} \{M_{t(j)} - M_{t(j-1)} \geq 0\} & \text{if } \varepsilon(j) = 1, \\ \{M_{t(j)} - M_{t(j-1)} < 0\} & \text{if } \varepsilon(j) = -1 \end{cases}$$

for $j = 1, 2, \dots, n$. Since $A = \bigcup_{1 \leq j \leq n, \varepsilon(j) = \pm 1} A \cap B_{1, \varepsilon(1)} \cap \dots \cap B_{n, \varepsilon(n)}$, we have for some $\varepsilon^*(j) (1 \leq j \leq n)$

$$P(A \cap B_{1, \varepsilon^*(1)} \cap \dots \cap B_{n, \varepsilon^*(n)}) > 0.$$

Then the process K defined by $K_s = \sum_{j=1}^n \varepsilon^*(j) I_{[t(j-1), t(j)]}(s)$ is a predictable process with $|K| \leq 1$, so that $\|K \cdot M\|_\infty \leq C$ must follow. On the contrary, we find

$$(K \cdot M)_t = \sum_{j=1}^n |M_{t(j)} - M_{t(j-1)}| > 2C$$

on the set $A \cap B_{1,\varepsilon^*(1)} \cap \dots \cap B_{n,\varepsilon^*(n)}$. Thus (a) implies (c). The implication (c) $\rightarrow$ (b) is trivial. Finally, to prove the implication (b) $\rightarrow$ (a), let us suppose $BMO \neq L^\infty$. Then by the result of Dellacherie, Meyer and Yor L^∞ is not dense in BMO, and further Kazamaki and Shiota have recently shown in [2] that the BMO-closure of L^∞ contains H^∞ . Therefore, combining these results, we have $BMO \neq H^\infty$. That is, the contraposition of the implication (b) $\rightarrow$ (a) is established. The latter half of the theorem follows at the same time. This completes the proof.

We now exemplify that H^∞ as well as L^∞ is not always closed in BMO under the same assumption as in the theorem. For that, consider the identity mapping S of R_+ onto R_+ . Let μ be the probability measure on R_+ defined by $\mu(S \in dx) = \sqrt{2/\pi} e^{-x^2/2} dx$ and $\mathbb{G}_t$ be the μ -completion of the Borel field generated by $S \wedge t$. Then $(R_+, \mathbb{G}, \mu; (\mathbb{G}_t))$, where $\mathbb{G} = \bigvee_{t \geq 0} \mathbb{G}_t$, is a probability system. Clearly, S is a stopping time over $(\mathbb{G}_t)$. We next consider in the usual way a probability system $(\Omega, \mathfrak{F}, P; (\mathfrak{F}_t))$ by taking the product of the system $(R_+, \mathbb{G}, \mu; (\mathbb{G}_t))$ with another system $(\Omega', \mathfrak{F}', P'; (\mathfrak{F}'_t))$ which carries a one dimensional Brownian motion $B = (B_t)$ starting at 0. Then the filtration $(\mathfrak{F}_t)$ satisfies the usual conditions and S is also a stopping time over this filtration. Let M denote the process B stopped at S . It is a continuous martingale over $(\mathfrak{F}_t)$ such that $\langle M \rangle_t = t \wedge S$ where $\langle M \rangle$ denotes the continuous increasing process associated with M . We first verify $M \in BMO$. Since $\{S > t\}$ is an $\mathfrak{F}_t$ -atom, we have

$$E[\langle M \rangle_\infty - \langle M \rangle_t | \mathcal{F}_t] = E[S - t | \mathcal{F}_t] 1_{\{t < S\}} \\ \leq (\int_t^\infty e^{-x^2/2} dx)^{-1} (\int_t^\infty (x-t) e^{-x^2/2} dx),$$

which converges to 0 as $t \rightarrow \infty$. That is, there is a constant $C > 0$ such that $E[\langle M \rangle_\infty - \langle M \rangle_t | \mathcal{F}_t] \leq C$ for every t . In our setting, this yields that $E[\langle M \rangle_\infty - \langle M \rangle_T | \mathcal{F}_T] \leq C$ for every stopping time T . Thus $M \in \text{BMO}$. As a matter of course, we have $M \in H^\infty$. Secondly, let $M^{(n)} = B^{n \wedge S}$ ($n=1, 2, \dots$). Then $M^{(n)} \in H^\infty$. Since $\langle M^{(n)} - M \rangle_t = t \wedge S - t \wedge n \wedge S$, we find

$$E[\langle M^{(n)} - M \rangle_\infty - \langle M^{(n)} - M \rangle_t | \mathcal{F}_t] \\ \leq (\int_{t \vee n}^\infty e^{-x^2/2} dx)^{-1} (\int_{t \vee n}^\infty (x - t \vee n) e^{-x^2/2} dx),$$

from which $M^{(n)}$ converges in BMO to M as $n \rightarrow \infty$. Consequently, $M \in \overline{H^\infty} \setminus H^\infty$.

REFERENCES

- [1] Dellacherie, C., Meyer, P.A. and Yor, M. (1978). Sur certaines propriétés des espaces de Banach H^1 et BMO, Sémin. de Prob. XII, Lecture Notes in Math. 649 98-11
- [2] Kazamaki, K. and Shiota, Y. (1985). Remarks on the class of continuous martingales with bounded quadratic variation, Tohoku Math. J., 37 101-106