

(1) Title: Ecological functions of persistent Japanese cedar litter in structuring stream macroinvertebrate assemblages

(2) Name of authors: Masaru Sakai, Yoshihiro Natuhara, Keitaro Fukushima, Ayumi Imanishi, Kensuke Imai, Makoto Kato

Addresses:

Masaru Sakai, Yoshihiro Natuhara, Ayumi Imanishi, and Makoto Kato: Graduate School of Global Environmental Studies, Kyoto University, Yoshida-honmachi, Sakyo-ku, Kyoto, 606-8501, JAPAN

Keitaro Fukushima: Field Science Education and Research Center, Kyoto University, Oiwake-cho, Sakyo-ku, Kyoto, 606-8501, JAPAN

Kensuke Imai: Department of Human Environments and Architectural Design, Osaka University of Human Sciences, 1-4-1 Shojaku, Settsu city, Osaka, 566-8501, JAPAN

(3) Information of corresponding author (Masaru Sakai):

Address: Yoshida-honmachi, Sakyo-ku, Kyoto, 606-8501, Japan

Telephone / Fax number: +81-75-753-6849 / +81-75-753-6849

E-mail address: m.sakai@fw3.ecs.kyoto-u.ac.jp

(4) Article type: Original articles

Subject area: Biology and ecology

Field: Entomology

(5) Number of text pages: 20

(6) Number of tables: 3

Number of figures: 3

Abstract

Stream macroinvertebrate assemblages are expected to be affected by the abundance and constitution of litter from surrounding forests. We compared forest floor cover, overland flow, stream environment, and stream macroinvertebrate assemblages between the catchments of a Japanese cedar plantation (CP) and a primary deciduous forest (DF). Both systems experience excessive deer browsing. Understory vegetation cover was higher in the DF than in the CP in summer, although cover was low (<20%), likely because of excessive deer browsing. Litter cover was much higher in the CP than in the DF in summer as a result of the long abscission period, low breakdown rate, and low dispersal rate of Japanese cedar litter compared to deciduous litter. Monthly overland flow was always lower in the CP than in the DF, and substrate size was smaller in the DF stream. In the CP, cedar litter accumulated in the stream, probably because of its low breakdown rate and morphology, and abundant shredder taxa characterized the macroinvertebrate assemblage. In contrast, abundant burrower taxa characterized the macroinvertebrate assemblage in the DF stream. These results imply that Japanese cedar litter functions in structuring the macroinvertebrate assemblage by supplying persistent food resources for detritivores, and by buffering fine sedimentation via overland flow under excessive deer browsing.

Key words

Denudation • Forest floor cover • Overland flow • Sediment runoff • Shredders

52 **Introduction**

53

54 In lotic ecosystems, allochthonous organic matter such as terrestrial litter and
55 autochthonous organic matter such as algae are vital basic food resources (Dodds 2002).
56 Litter from riparian forests constitutes a higher proportion of available food for
57 macroinvertebrates in headwater streams than in downstream areas (Vannote et al. 1980).
58 Riparian forests influence stream ecosystems by providing terrestrial invertebrates as
59 food for fish (Nakano and Murakami 2001), controlling water temperature (Sugimoto et
60 al. 1997; Richardson and Danehy 2007), purifying water by filtering it through complex
61 networks of soil and roots (Peterjohn and Correll 1984; Pinay and Decamps 1988), and
62 creating diverse habitats via woody debris (Inoue and Nakano 1998; Richardson and
63 Danehy 2007).

64 Forest type (e.g., deciduous broad-leaf, evergreen coniferous) is one of the main
65 terrestrial conditions within catchments, and ecological properties tend to vary with
66 forest type. The biomass and diversity of understory vegetation are generally higher in
67 deciduous forests (Barbier et al. 2008), whereas annual litter input (Gregory et al. 1991)
68 and standing stock of accumulated litter (Augusto et al. 2002) tend to be higher in
69 coniferous systems. These ecological properties affect hydrological processes on slopes
70 (Sidle et al. 2007) as well as stream environments and macroinvertebrates (Gregory et al.
71 1991). For example, Friberg (1997) noted that both the standing stock of detritus and
72 the abundance of shredders in streams are greater in coniferous than in deciduous
73 catchments.

74 Nearly 27% of the land in Japan is covered by evergreen coniferous plantations
75 (Nagaike et al. 2006). Japanese cedar (*Cryptomeria japonica*) is one of the most

important plantation species in Japan (Ota 2007), covering about 4.5 million hectares. In contrast to deciduous trees, litterfall of Japanese cedar continues from winter to early spring, and cedar litter has a much narrower dispersal range (Kanasashi and Hattori 2011). Japanese cedar litter breaks down at a slower rate than deciduous litter in both terrestrial and aquatic systems (Nakane 1995; Hisabae et al. 2010). Therefore, litter is expected to remain on the forest floor and in streams for longer periods in Japanese cedar plantations than in deciduous forests. This may result in particular stream macroinvertebrate assemblages characteristic of cedar plantations (Yoshimura and Maeto 2006; Yoshimura 2007).

We studied the effects of Japanese cedar litter accumulation on stream macroinvertebrate assemblages by comparing forest floor cover, overland flow, stream environment, and macroinvertebrate assemblages between a Japanese cedar plantation and a deciduous forest. Both of the systems experience excessive deer browsing characterized by the denudation of vegetation. We expected to find characteristic differences in macroinvertebrate assemblages between these systems, particularly because cedar litter provides better protection against soil erosion and should help reduce erosion of denuded forest floors.

Materials and methods

Study site

The study area is located at the Ashiu Forest Research Station of the Field Science Education and Research Center, Kyoto University (35°20'N, 135°45'E; Fig. 1). Average

annual precipitation at the site was 2298 mm and mean annual temperature was 11.9°C from 1976 to 2005. Maximum snow depth in winter generally exceeds 2 m, and snow cover typically exists from mid-December to early April. The geological components of the area are sandstone, mudstone, and shale of the Tanba Belt of the Mesozoic. Most of the soil is brown forest soil. Excessive deer browsing has caused drastic decreases in the abundance and diversity of understory plants since the late 1990s (Kato and Okuyama 2004; Tanaka et al. 2008). Soil erosion via denudation by deer has increased sediment runoff and subsequent sandy sedimentation of streambeds in deciduous forests (Sakai et al. 2012).

The study was conducted in a 74-year-old Japanese cedar plantation catchment (CP; area, 1.33 ha) and a primary deciduous forest catchment (DF; area, 1.66 ha), both draining first-order streams of the Yura River (Fig. 1). The two catchments were located close to one another. The DF comprised primary cool-temperate deciduous forest and was dominated by *Fagus crenata* (dominance in basal area, 74.0%), *Acer mono* (10.6%), *Acer japonicum* (5.0%), and *Carpinus tschonoskii* (2.1%), whereas the CP was dominated by *Cryptomeria japonica* (85.3%), *Quercus crispula* (7.8%), and *Aesculus turbinata* (2.3%). The two streams were topographically similar and typical colluvial valleys (Bisson et al. 2006), where riffle-pool structures are rarely observed. A 30-m study reach in the lower section of each stream was established for surveying macroinvertebrates. The two reaches had perennial flow, and similar channel slope, width, and depth (Table 1).

Forest floor cover and overland flow

Because forest floor cover is a good predictor of soil erosion potential (e.g., Lyon and Sagers 1998; Wear et al. 1998; Heartsill-Scalley and Aide 2003), we evaluated coverage of understory vegetation and litter as well as water runoff along the slopes of each catchment to examine the effect of soil inflow on stream substrate and macroinvertebrate assemblages, following Sakai et al. (2012). One belt transect (1-m wide) was established from the valley floor to 20 m up the upper slope on the right and left banks of each catchment. These transects were located in areas where the most overland flow entering the streams was likely to be generated. Coverage (%) of vegetation within 30 cm of the ground was visually estimated by one individual in 1 × 2-m plots equally spaced along each transect (10 plots per bank). The survey was conducted in June and August 2008. In August 2011, 10 quadrats (40 × 40 cm) were also randomly located on the lower, middle, and upper slopes of each catchment, and litter coverage (%) within the quadrats was visually estimated by one individual. We also evaluated the standing stock of litter in the riparian zone. We collected litter from four randomly located quadrats (40 × 40 cm) along the 30-m study reach in each catchment in August and November 2009. Collected litter was dried at 60°C for 1 week and then weighed.

Plots for measuring water runoff (0.5 × 2.0 m) were established on the midslopes of each catchment near the vegetation transect (Fig. 1). The coverage of understory vegetation and litter in the plots was representative of other midslope locations for each catchment. Plastic borders were inserted about 5 cm into the soil along all sides of the plots, and a trough was inserted several centimeters into the soil (parallel to the slope

direction) to collect storm runoff. Runoff from these plots was routed to a rain gauge (Davis Instruments, Rain Collector II) to estimate discharge per month. Overland flow was monitored from July to November 2010.

Stream environment

Channel slope was measured at 6 points at 5 m intervals using a laser surveying instrument (Laser Technology, Trupulse 200), and stream width was measured at 7 transects located at 5 m intervals in each study reach in July 2010. Water depth and current velocity were also measured at 10 points in the centers of the streams in each study reach in July 2010. The current velocities of the surface and bottom water (about 0–2 cm above the streambed) were calculated by averaging three repeated 5-s measurements from an electromagnetic current meter (Kenek, VE10). The measurements were done under base-flow conditions. Litter coverage (%) in six quadrats (40 × 40 cm) randomly located on the streambed in each study reach was visually estimated by one individual in August 2011.

To evaluate grain-size distribution, we randomly collected three samples of stream substrates (1000 ml) by shovel, according to Grost et al. (1991) and Zweig and Rabení (2001), in each study reach. The substrate samples were sieved into 10 grain sizes: >31.5, 31.5–16, 16–8, 8–4, 4–2, 2–1, 1–0.5, 0.5–0.25, 0.25–0.125, and 0.125–0.063 mm. The substrates of each size were dried at 105°C for 24 h and weighed, and we determined the effective grain size (D_{50}) of each substrate sample. This survey was conducted in December 2010.

To evaluate the amount of organic matter of fine deposition, we collected one

sample of sandy deposition (300 ml) by shovel in each study reach in July 2010. The samples were sieved into six fractions: >2, 2–1, 1–0.5, 0.5–0.25, 0.25–0.125, and 0.125–0.063 mm, and dried at 105°C for 24 h. Organic material was decomposed by adding hydrogen peroxide at 90°C for 3 days according to Dane and Topp (2002) and samples were then reweighed. The amount of organic matter was calculated for each fraction.

Hemispherical photographs were taken at 6 points at 6 m intervals in each study reach in August 2008 and April 2009 to determine light conditions above the streambed. Photographs were taken with a digital camera (Nikon, E995) equipped with a fish-eye lens (Nikon, FC-E8) that was fixed horizontally at 1 m above the streambed. Photosynthetically active radiation (PAR; $\text{mol m}^{-2} \text{day}^{-1}$) was estimated from the photographs, and we determined relative photosynthetically active radiation (rPAR %) to the open environment using Gap Light Analyzer version 2.0 (Frazer et al. 1999).

Periphyton

Because fine sediment may affect macroinvertebrates by reducing periphyton, algal abundance was estimated in each study reach in August, September, and October 2008. Periphyton was sampled from four randomly-collected submerged rocks (maximum diameter, 13–26 cm) by brushing 4 cm^2 of the upper surface of the rocks and filtering the water through glass microfiber filters (Whatman, GF/F). These samples were ground in 90% acetone and then centrifugally separated (Hitachi, CF16RXII) after being dried and frozen. The absorbance of the periphyton samples was measured at 630, 647, 664,

and 750 nm (Hitachi, U-1800). The amount of chlorophyll *a* was calculated by substituting the measured values into the formula of Jeffrey and Humphrey (1975).

Macroinvertebrates

Macroinvertebrates were sampled every month from May to November 2008 and in April 2009 in each study reach. Four samples per stream were collected using a Surber net (25 × 25 cm, 1-m length, 0.5-mm mesh) on each sampling occasion. Sampling points were randomly selected among locations where the channel width was >50 cm. Samples were preserved in 70% ethanol in the field and returned to the laboratory.

Macroinvertebrates were sorted by eye and identified to the lowest possible taxonomic level using a stereomicroscope (Nikon, SMZ800), following Kawai and Tanida (2005) and Merritt et al. (2008). Identified macroinvertebrates were classified by life-form type (burrowers, clingers, crawlers, side-swimmers, or swimmers) and functional feeding group (collector-filterers, collector-gatherers, grazers, predators, or shredders) based on Takemon (2005), Merritt et al. (2008), Kobayashi et al. (2010) and our unpublished data.

Statistical analyses

Student's *t*-tests were used to determine the differences between the CP and DF in understory coverage on the forest floor (June, August), litter cover on the forest floor (lower, middle, upper slope), litter quantity in the riparian zone (August, November), channel slope, stream width, water depth, current velocity (surface, bottom), litter

coverage in the stream, D₅₀ of stream substrate, rPAR (April, August), and algal abundance (August, September, October). A Friedman test was used to test for differences between the CP and DF in monthly overland flow during the monitoring period.

To test for differences between the CP and DF in the total abundance and taxa richness of macroinvertebrates and the abundance of each of the most dominant taxa (>1% of the individuals in all samples), a two-way analysis of variance was performed using stream ($n = 2$) and month ($n = 8$) as factors. When the stream $\times$ month interaction was significant, the difference between streams in each month was tested using Fisher's protected least significant difference test.

Proportional data were arcsine-transformed, and abundance data were log-transformed to normalize distributions and standardize variance structures prior to statistical analyses.

Results

Forest floor cover and overland flow

Vegetation cover on hillslopes was two to three times lower in the CP than in the DF in June and August, whereas litter cover at any position on the hillslopes was three to five times higher in the CP than in the DF in August (Table 1). Litter in the riparian zone was also significantly more abundant in the CP than in the DF in August, although not in November (Table 1). Monthly overland flow was significantly lower in the CP than in the DF from July to November (Fig. 2, $P < 0.05$).

Stream environment

Mean channel slope, stream width, and water depth in July were largely similar between the CP and DF (Table 1). Current velocities of both surface and bottom water were slower in the CP than in the DF in July (Table 1). Litter cover in streams was more than 30 times higher in the CP than in the DF in August (Table 1 and Fig. 3). The D_{50} of stream substrate was significantly higher in the CP than in the DF in December (Table 1). Fine particulate organic matter (>0.25 mm) was more abundant in the CP than in the DF in July (Table 1). Except for rPAR in April, which was lower in the CP, rPAR and algal abundance did not differ significantly between the catchments.

Macroinvertebrates

A total of 4465 macroinvertebrate individuals representing 100 taxa in 13 orders, 49 families, and 78 genera were collected. Total abundance and taxa richness of the samples did not differ between the CP and DF, or among months (Table 2), although the stream $\times$ month interaction was significant; total abundance in November and taxa richness in October were higher in the DF than in the CP.

Forest type had a significant effect on the abundance of 13 of the 26 dominant taxa. The abundances of *Gammarus nipponensis*, *Nemoura* spp., *Amphinemura* spp., *Geothelphusa dehaani*, and *Lepidostoma crassicorne* were significantly greater in the CP than in the DF (Table 2). The abundances of Lumbriculidae gen. spp., *Caroperla pacifica*, *Ecdyonurus tigris*, Chironominae spp., Ceratopogonidae gen. spp., *Psilotreta*

kisoensis, *Niponiella limbatella*, and Tabanidae gen. spp. were significantly greater in the DF than in the CP (Table 2). The site $\times$ month interaction was significant for *Ga. nipponensis*, *Nemoura* spp., *E. tigris*, *Amphinemura* spp., Chironominae spp., *L. crassicornis*, *Epoicocladus* sp.1, Ceratopogonidae gen. spp., and *P. kisoensis* (Table 2). If a significant difference between the streams was detected in some months, *Ga. nipponensis*, *Nemoura* spp., *Amphinemura* spp., and *L. crassicornis* were always more abundant in the CP, whereas *Epoicocladus* sp.1, Ceratopogonidae gen. spp., and *P. kisoensis* were always more abundant in the DF (Table 3). *E. tigris* and Chironominae spp. were more abundant in the CP in April but in the DF in other months (Table 3).

Discussion

In summer, vegetative cover on hillslopes was lower in the CP than in the DF, whereas litter cover at any position along the hillslope and standing stock of litter in the riparian zone were higher in the CP. The lower amount of understory vegetative cover in the CP may be attributed to the low light intensity at the forest floor beneath the dense canopy of cedar plantations (Katagiri et al. 1985). However, understory cover was very low (<20%) even in the DF compared to an adjacent deciduous forest where deer are excluded (Sakai et al. 2012). The forest floor in the CP would be covered with more abundant litter at least in summer because Japanese cedar litter exhibits a long period of abscission, low breakdown rate, and low dispersal compared to deciduous litter (Nakane 1995; Inagaki et al. 2004; Kanasashi and Hattori 2011). Overland flow runoff was always lower in the CP. Generally, understory vegetation and litter protect the soil infiltration capacity against rain drop impact (Onda and Yukawa 1994) and prevent

overland flow from discharging soil (Miura et al. 2002; Sidle et al. 2007; Gomi et al. 2008). Our results suggest that an abundance of litter is more important than understory cover in buffering runoff of overland flow in forests experiencing excessive deer browsing.

Litter cover in the streambed was also higher in the CP than in the DF in summer. Japanese cedar litter breaks down at a slower rate than deciduous litter in streams (Hisabae et al. 2010) as well as on the forest floor, and accumulates as complex structures composed of shoot or twig litter within streams. These structures persist for a prolonged period, probably buffering current velocity in the CP. The D_{50} of substrate was higher in the CP. Sakai et al. (2012) reported that higher runoff of overland flow induced fine sedimentation in the streambed in the DF compared to the deer-excluded catchment. The reduced overland flow protected the CP stream from fine sedimentation. Although fine sedimentation may affect periphyton (Yamada and Nakamura 2002), algal abundance did not differ between the CP and DF.

We observed differences in total abundance and taxa richness of macroinvertebrate assemblages between the DF and CP, but only in some seasons. In contrast, we observed differences in the structure of macroinvertebrate assemblages between the streams throughout the year. Yoshimura and Maeto (2006) demonstrated that the structure, but not the abundance or taxa richness, of macroinvertebrate assemblages in streams varied between a Japanese cedar plantation and a natural broad-leaf forest. In Europe, several studies have documented negative effects of coniferous plantations on the abundance, diversity, and biomass of macroinvertebrate assemblages (Ormerod et al. 1993; Thomsen and Friberg 2002). However, Collier and Halliday (2000) reported higher richness and diversity of macroinvertebrate assemblages in coniferous plantations

313 compared to deciduous forests in New Zealand and suggested that slower breakdown
314 rates of conifer litter provide increased habitat for invertebrates.

315 All dominant shredder taxa (*Ga. nipponensis*, *Nemoura* spp., *Amphinemura* spp., and
316 *L. crassicornis*) were more abundant in the CP than in the DF. The strong dominance of
317 *Ga. nipponensis* has also been reported in other streams that drain Japanese cedar
318 plantations in Japan (Hisabae et al. 2010). Japanese cedar litter is generally considered
319 an unsuitable food for invertebrates because of its toughness and low nutritive quality
320 (Hisabae et al. 2010). However, cedar litter should provide a constantly available food
321 resource for shredder invertebrates due to its long period of abscission, low breakdown
322 rate, and low dispersal. The advantage of slow-decaying litter for shredders has also
323 been reported by Friberg and Jacobsen (1994). The shredders that were abundant in the
324 CP suggest that Japanese cedar litter supplies persistent and abundant organic matter. In
325 contrast, the abundances of burrowers (Lumbriculidae gen. spp., *Epoicocladus* sp.1,
326 Ceratopogonidae gen. spp., and Tabanidae gen. spp.) were greater in the DF, where the
327 increased level of fine sediments should provide more suitable habitats for burrower
328 invertebrates.

329 In conclusion, the macroinvertebrate assemblages in a first-order stream draining a
330 Japanese cedar plantation differed from those in a first-order stream draining a
331 deciduous forest under excessive deer browsing. Our findings suggest that Japanese
332 cedar litter functions in structuring macroinvertebrate assemblages by supplying
333 persistent food resources for detritivores, and by buffering fine sedimentation via
334 overland flow. The latter function might also be important in systems that have not
335 experienced excessive deer browsing because litter may prevent soil erosion more
336 effectively than understory vegetation (Miura et al. 2002).

Acknowledgements

We thank Dr. Yasuhiro Takemon for helping with aquatic macroinvertebrate identification, and Dr. Hideaki Miyashita for helping with chlorophyll measurements of periphyton. In addition, we are indebted to the staff of Ashiu Forest Research Station, for enabling us to conduct this research. Invaluable field assistance was provided by academic colleagues. Considerable advice was provided by the two anonymous reviewers. Part of this study was funded by ProNatura Fund and JSPS Research Fellowships for Young Scientists.

References

- Augusto L, Ranger J, Binkley D, Rothe A (2002) Impact of several common tree species of European temperate forests on soil fertility. *Annu For Sci* 59:233-253
- Barbier S, Gosselin F, Balandier P (2008) Influence of tree species on understory vegetation diversity and mechanisms involved - A critical review for temperate and boreal forests. *For Ecol Manage* 254:1-15
- Bisson PA, Montgomery DR, Buffington JM (2006) Valley segments, stream reaches, and channel units. In: Hauer FR, Lamberti GA (eds) *Methods in stream ecology*. 2nd edition, Academic Press, pp23-49
- Collier KJ, Halliday JN (2000) Macroinvertebrate-wood associations during decay of plantation pine in New Zealand pumice-bed streams: stable habitat or trophic subsidy. *J N Am Benthol Soc* 19:94-111

361 Dane JH, Topp GC (2002) Methods of soil analysis part 4: physical method. Soil
 362 Science Society of America Book series 5, Madison, WI
 363 Dodds WK (2002) Freshwater ecology: Concepts and environmental applications.
 364 Academic Press
 365 Frazer GW, Canham CD, Lertzman KP (1999) Gap light analyzer (GLA) imaging
 366 software to extract canopy structure and gap light transmission indices from
 367 true-colour fisheye photographs: User manual and program documentation, v. 2.0.
 368 Simon Frazer University / Institute of Ecosystem Studies, Burnaby, Canada /
 369 Millbrook, NY
 370 Friberg N (1997) Benthic invertebrate communities in six Danish forest streams: impact
 371 of forest type on structure and function. *Ecography* 20:19-28
 372 Friberg N, Jacobsen D (1994) Feeding plasticity of two detritivore-shredders. *Freshw*
 373 *Biol* 32:133-142
 374 Gomi T, Sidle RC, Ueno M, Miyata S, Kosugi K (2008) Characteristics of overland
 375 flow generation on steep forested hillslopes of central Japan. *J Hydrol* 361: 275-290.
 376 Gregory SV, Swanson FJ, Mckee WA, Cummins KW (1991) An ecosystem perspective
 377 of riparian zones - Focus on links between land and water. *Bioscience* 41:540-551
 378 Grost RT, Hubert WA, Wesche TA (1991) Field comparison of three devices used to
 379 sample substrate in small streams. *N Am J Fish Manage* 11:347-351
 380 Heartsill-Scalley T, Aide TM (2003) Riparian vegetation and stream condition in a
 381 tropical agriculture – secondary forest mosaic. *Ecol Appl* 13:225-234
 382 Hisabae M, Sone S, Inoue M (2010) Breakdown and macroinvertebrate colonization of
 383 needle and leaf litter in conifer plantation streams in Shikoku, southwestern Japan. *J*
 384 *For Res* 16:108-115

385 Inagaki Y, Miura S, Kohzu A (2004) Effects of forest type and stand age on litterfall
 386 quality and soil N dynamics in Shikoku district, southern Japan. *For Ecol Manage*
 387 202:107-117
 388 Inoue M and Nakano S (1998) Effects of woody debris on the habitat of juvenile masu
 389 salmon (*Oncorhynchus masou*) in northern Japanese streams. *Freshw Biol* 40:1–16.
 390 Jeffrey SW, Humphrey GF (1975) New spectrophotometric equations for determining
 391 chlorophyll a, b, c1 and c2 in higher plants, algae, and natural phytoplankton.
 392 *Biochem Physiol Pflanz* 167:191-194
 393 Kanasashi T, Hattori S (2011) Seasonal variation in leaf-litter input and leaf dispersal
 394 distances to streams: the effect of converting broadleaf riparian zones to conifer
 395 plantations in central Japan. *Hydrobiologia* 661:145-161
 396 Katagiri S, Ishii H, Miyake N (1985) Studies on above ground biomass of understory
 397 vegetation in sugi (*Cryptomeria japonica* D. Don) stand. *Bull Fac Agric, Shimane*
 398 *Univ* 19:34-38 (in Japanese)
 399 Kato M, Okuyama Y (2004) Changes in the biodiversity of a deciduous forest
 400 ecosystem caused by an increase in the sika deer population at Ashiu, Japan. *Contr*
 401 *Biol Lab Kyoto Univ* 29:237-448
 402 Kawai T, Tanida K (2005) *Aquatic insects of Japan: Manual with keys and illustrations.*
 403 Tokai University Press, Kanagawa (in Japanese)
 404 Kobayashi S, Gomi T, Sidle RC, Takemon Y (2010) Disturbances structuring
 405 macroinvertebrate communities in steep headwater streams: relative importance of
 406 forest clearcutting and debris flow occurrence. *Can J Fish Aquat Sci* 67:427-444
 407 Lyon J, Sagers CL (1998) Structure of herbaceous plant assemblages in a forested
 408 riparian landscape. *Plant Ecol* 138:1-16

409 Merritt RW, Cummins KW, Berg MB (2008) An Introduction to the Aquatic Insects of
 410 North America. Kendall / Hunt Publishing Company, Iowa
 411 Miura S, Hirai K, Yamada T (2002) Transport rates of surface materials on steep
 412 forested slopes induced by raindrop splash erosion. J For Res 7:201-211
 413 Nagaike T, Masaki T, Ito S (2006) Special feature: ecology and management of conifer
 414 plantations in Japan: control of tree growth and maintenance of biodiversity. J For
 415 Res 11:215-216
 416 Nakane K (1995) Soil carbon cycling in a Japanese cedar (*Cryptomeria japonica*)
 417 plantation. For Ecol Manage 72:185-197
 418 Nakano S, Murakami M (2001) Reciprocal subsidies: dynamic interdependence
 419 between terrestrial and aquatic food webs. Proc Natl Acad Sci USA 98:166–170
 420 Onda Y, Yukawa N (1994) The influence of understories and litter layer on the
 421 infiltration of forested hillslope. In: Ohta T (ed) Proceedings of international
 422 symposium on forest hydrology. University of Tokyo, Japan, pp 107-114
 423 Ormerod SJ, Rundle SD, Lloyd EC, Douglas AA (1993) The influence of riparian
 424 management on the habitat structure and macroinvertebrate communities of upland
 425 streams draining plantation forests. J Appl Ecol 30:13-24
 426 Ota I (2007) A forest owners' cooperative in Japan: obtaining benefits of certification
 427 for small-scale forests. Unasylva 58:64-66
 428 Peterjohn WT, Correll DL (1984) Nutrient dynamics in an agricultural watershed:
 429 observations on the role of a riparian forest. Ecology 65:1466–1475
 430 Pinay G, Decamps H (1988) The role of riparian woods in regulation nitrogen fluxes
 431 between the alluvial aquifer and surface water: a conceptual model. Regul. Rivers
 432 Res Manage 2:507-516

433 Richardson JS, Danehy RJ (2007) A synthesis of the ecology of headwater streams and
 434 their riparian zones in temperate forests. *For Sci* 53:131-147
 435 Sakai M, Natuhara Y, Imanishi A, Imai K, Kato M (2012) Indirect effects of excessive
 436 deer browsing through understory vegetation on stream insect assemblages. *Popul*
 437 *Ecol* 54:65-74
 438 Sidle RC, Hirano T, Gomi T, Terajima T (2007) Hortonian overland flow from
 439 Japanese forest plantations – an aberration, the real thing, or something in between?
 440 *Hydrol Process* 21:3237-3247
 441 Sugimoto S, Nakamura F, Ito A (1997) Heat budget and statistical analysis of the
 442 relationship between stream temperature and riparian forest in the Toikanbetu River
 443 basin, northern Japan. *J For Res* 2:103-107
 444 Takemon Y (2005) Life-type concept and functional feeding groups of benthos
 445 communities as indicators of lotic ecosystem conditions. *Jpn J Ecol* 55:189-197 (in
 446 Japanese)
 447 Tanaka Y, Takatsuki S, Takayanagi A (2008) Decline of *Sasa palmata* community by
 448 grazing of Sika deer (*Cervus nippon*) at Ashiu Research Forest Station. *For Res*
 449 *Kyoto* 77:13-23 (in Japanese)
 450 Thomsen ST, Friberg N (2002) Growth and emergence of the stonefly *Leuctra nigra* in
 451 coniferous forest streams with contrasting pH. *Freshw Biol* 47:1159-1172
 452 Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE (1980) The river
 453 continuum concept. *Can J Fish Aquat Sci* 37:130-137
 454 Wear DN, Turner MG, Naiman RJ (1998) Land cover along an urban-rural gradient:
 455 implications for water quality. *Ecol Appl* 8:619-630
 456 Yamada H, Nakamura F (2002) Effect of fine sediment deposition and channel works on

periphyton biomass in the Makomanai River, northern Japan. Riv Res Appl
18:481-493

Yoshimura M (2007) Comparison of stream benthic invertebrate assemblages among
forest types in the temperate region of Japan. Biodivers Conserv 16:2137-2148

Yoshimura M, Maeto K (2006) Comparison of an aquatic invertebrate assemblage
between an old-growth natural forest and planted coniferous forest basins in a
Japanese temperate region: the Kuroson stream in the Shimanto River basin.
Landscape Ecol Eng 2:81-89

Zweig LD, Rabení CF (2001) Biomonitoring for deposited sediment using benthic
invertebrates: a test on 4 Missouri streams. J N Am Benthol Soc 20:643-657

Table 1 Comparisons of environmental characteristics between the CP and DF. Differences between the catchments in litter cover, understory vegetation cover, quantity of litter at the bottom of the slope, channel slope, litter cover in the stream, stream width, water depth, current velocity, D₅₀ of stream substrate, relative photosynthetically active radiation, and algal abundance were tested using Student's t-tests.

Variable		CP	DF	<i>t</i> value	Degrees of freedom	<i>P</i>
Forest type		Japanese cedar plantation	Deciduous forest			
Catchment area (ha)		1.31	1.66			
Cover of litter (%): Bottom slope		90.50 ± 2.33	25.75 ± 2.06	14.813	38	<0.0001
: Middle slope		84.50 ± 2.78	26.50 ± 3.44	10.842	38	<0.0001
: Upper slope		78.00 ± 2.20	13.05 ± 1.77	18.011	38	<0.0001
Cover of understory vegetation (%)	Jun.	6.18 ± 2.12	15.20 ± 4.29	2.307	38	0.027
	Aug.	5.50 ± 2.17	16.62 ± 3.99	2.865	38	0.007
Quantity of litter at bottom slope (g m ⁻²)	Aug.	802.87 ± 125.14	294.87 ± 73.23	3.504	6	0.013
	Nov.	876.72 ± 125.54	611.53 ± 210.73	1.081	6	0.321
Channel slope (degrees)		12.75 ± 1.57	11.13 ± 1.75	0.688	10	0.507
Cover of litter in stream (%)		66.00 ± 3.93	2.00 ± 1.11	15.138	10	<0.0001
Stream width (m)		1.00 ± 0.11	1.17 ± 0.27	0.591	12	0.571
Water depth (cm)		4.79 ± 0.29	4.44 ± 0.40	0.712	18	0.486
Current velocity (cm s ⁻¹): Surface water		10.83 ± 0.75	13.80 ± 0.53	3.220	18	0.002
: Bottom water		4.08 ± 0.46	12.19 ± 0.61	10.601	18	<0.0001
Effective grain size (D ₅₀) (cm)		15.94 ± 0.69	6.34 ± 0.86	8.726	4	0.001
Quantity of organic matter (mg ml ⁻¹): 2 mm		0.41	0.22			
: 1 mm		1.25	0.43			
: 0.5 mm		1.60	0.16			
: 0.25 mm		1.04	0.30			
: 0.125 mm		0.20	0.26			
: 0.063 mm		0.15	0.16			
Relative photosynthetically active radiation (%)	Apr.	16.15 ± 1.14	42.68 ± 1.50	13.760	10	<0.0001
	Aug.	19.41 ± 1.46	22.74 ± 2.59	1.096	10	0.299
Algal abundance (mg Chl. <i>a</i> cm ⁻²)	Aug.	0.24 ± 0.14	0.19 ± 0.07	0.303	6	0.772
	Sep.	0.21 ± 0.07	0.08 ± 0.01	1.749	6	0.179
	Oct.	0.10 ± 0.03	0.07 ± 0.02	1.189	6	0.279

Values are means ± standard errors.

Bold characters indicate statistical significance (< 0.05).

Table 2 Results of analyses of variance testing for effects of stream and month on total abundance, taxa richness, and abundance of the 26 most dominant macroinvertebrate taxa.

Variable	LFT	FFG	Mean		Factors					
					Stream		Month		Stream × month	
			CP	DF	F	P	F	P	F	P
Total abundance	-	-	72.28	67.25	0.63	0.432	1.11	0.374	3.22	0.007
Taxa richness	-	-	17.09	17.59	0.25	0.618	1.21	0.317	3.06	0.010
Abundance of each taxon										
<i>Gammarus nipponensis</i>	SS	S	16.34	0.25	211.94	<0.0001	4.64	0.001	4.02	0.002
<i>Ephemera japonica</i>	B	C/G	5.16	9.31	3.16	0.082	2.22	0.049	0.85	0.553
<i>Nemoura</i> spp.	Cr	S	8.84	4.31	10.47	0.002	11.37	<0.0001	2.74	0.018
<i>Paraleptophlebia japonica</i>	B	C/G	2.91	4.66	0.50	0.485	2.81	0.016	1.59	0.163
Lumbriculiidae gen. spp.	B	C/G	2.31	5.00	9.40	0.004	2.24	0.047	1.76	0.118
<i>Caroperla pacifica</i>	Cr	P	2.38	4.88	5.37	0.025	5.73	<0.0001	1.16	0.342
Orthocladinae spp.	B	C/G, G	1.81	3.75	0.89	0.352	0.24	0.406	0.89	0.526
<i>Ecdyonurus tigris</i>	Cr	G	1.63	3.38	8.34	0.006	4.53	0.001	4.82	0.000
<i>Amphinemura</i> spp.	Cr	S	3.97	0.91	14.84	0.000	4.98	0.000	2.41	0.034
Chironominae spp.	B	C/F, C/G	0.47	3.63	12.26	0.001	4.33	0.001	5.09	0.000
<i>Geothelphusa dehaani</i>	Cr	C/G	3.31	0.66	42.82	<0.0001	1.07	0.396	0.79	0.603
<i>Togoperla limbata</i>	Cr	P	1.97	1.75	0.76	0.388	1.27	0.284	0.90	0.515
<i>Lepidostoma crassicornae</i>	Cr	S	2.69	0.75	13.68	0.001	3.98	0.002	3.52	0.004
<i>Baetis yoshinensis</i>	S	G	1.59	1.34	0.14	0.707	1.81	0.108	1.41	0.224
Diamesinae spp.	B	C/G, G	1.00	1.88	3.63	0.063	10.10	<0.0001	1.20	0.320
<i>Hydropsyche</i> sp.	Cl	C/F, C/G	1.88	0.78	3.42	0.071	2.28	0.044	0.76	0.625
<i>Epoicocladus</i> sp.1	B	C/G	0.66	1.78	3.81	0.057	19.89	<0.0001	5.53	0.000
<i>Perissoneura paradoxa</i>	Cr	C/G	0.72	1.41	3.17	0.081	0.69	0.677	2.17	0.054
Ceratopogonidae gen. spp.	B	C/G	0.50	1.50	11.62	0.001	6.73	<0.0001	3.22	0.007
<i>Psilotreta kisoensis</i>	Cr	C/G	0.00	1.97	48.98	<0.0001	3.29	0.006	3.29	0.006
<i>Epoicocladus</i> sp.2	B	C/G	1.22	0.72	0.91	0.345	6.39	<0.0001	1.21	0.318
<i>Paralichas</i> sp.	B	C/G	0.91	1.00	0.18	0.674	1.08	0.078	0.66	0.706
<i>Niponiella limbatella</i>	Cr	P	0.38	1.47	17.05	0.000	5.91	<0.0001	0.65	0.717
Tabanidae gen. spp.	B	P	0.00	1.81	41.79	<0.0001	0.69	0.678	0.69	0.678
<i>Rhyacophila</i> sp.	Cr	P	0.97	0.66	1.39	0.244	0.62	0.734	0.89	0.525
<i>Dolophilodes</i> sp.DB	Cl	C/F, C/G	0.72	0.69	0.03	0.870	1.76	0.117	2.05	0.068

df = 1,48 for effects of stream, and 7,48 for month and stream × month.

Bold characters indicate statistical significance (< 0.05).

Life form type (LFT); B: Burrower, Cl: Clinger, Cr: Crawler, S: Swimmer, SS: Side swimmer.

Functional feeding group (FFG); C/F: Collector-filterer, C/G: Collector-gatherer, G: Grazer, P: Predator, S: Shredder.

Table 3 Results of Fisher's protected least significant difference test for difference between streams in each month in total abundance, taxa richness and abundance of the 9 macroinvertebrate taxa, in which interaction between stream and month in analysis of variance were significant. The stream (CP or DF) that showed greater abundance or richness is denoted if significant difference was detected ($P < 0.05$).

Variable	May	Jun.	Jul	Aug.	Sep.	Oct.	Nov.	Apr.
Total abundance	-	-	-	-	-	-	DF	-
Taxa richness	-	-	-	-	-	DF	-	-
Abundance of each taxon								
<i>Gammarus nipponensis</i>	CP	CP	CP	CP	CP	CP	CP	CP
<i>Nemoura</i> spp.	-	CP	-	CP	-	-	CP	-
<i>Amphinemura</i> spp.	-	CP	-	CP	-	-	-	-
<i>Lepidostoma crassicorne</i>	-	-	-	CP	-	-	CP	-
<i>Epoicocladius</i> sp.1	-	-	-	DF	DF	DF	-	-
Ceratopogonidae gen. spp.	-	-	DF	-	-	DF	-	-
<i>Psilotreta kisoensis</i>	-	-	DF	DF	DF	DF	-	DF
<i>Ecdyonurus tigris</i>	-	-	-	-	DF	DF	-	CP
Chironominae spp.	-	DF	DF	-	-	-	-	CP

Figure legends

Fig. 1 The study site in the Ashiu Research Forest Station, Kyoto Prefecture, Japan.

Fig. 2 Monthly runoff of overland flows (mm) for the cedar plantation and deciduous forest catchments.

Fig. 3 Landscapes (a, c) and close-up views of streams (b, d) in the cedar plantation and deciduous forest catchments.

Fig. 1 (Sakai et al.)





Fig. 2 (Sakai et al.)



Cedar plantation catchment (CP)



Deciduous forest catchment (DF)

