

Which Structural Rules Admit Cut Elimination? — An Algebraic Criterion (Excerpt)

照井 一成 国立情報学研究所

Kazushige Terui National Institute of Informatics

terui@nii.ac.jp

概要

This is an excerpt of our recent paper [Ter05]. See [Ter05] for the details.

1 Introduction

Gentzen's original sequent calculus contains three structural rules:

$$\begin{array}{ccc} \text{Exchange:} & \text{Weakening:} & \text{Contraction:} \\ \frac{\Gamma, \alpha, \beta, \Delta \Rightarrow \gamma}{\Gamma, \beta, \alpha, \Delta \Rightarrow \gamma} \mathbf{e} & \frac{\Gamma, \Delta \Rightarrow \gamma}{\Gamma, \alpha, \Delta \Rightarrow \gamma} \mathbf{w} & \frac{\Gamma, \alpha, \alpha, \Delta \Rightarrow \gamma}{\Gamma, \alpha, \Delta \Rightarrow \gamma} \mathbf{c} \end{array}$$

where α, β and γ stand for formulas and Γ and Δ stand for sequences of formulas (we only consider intuitionistic sequents in this paper). In addition, one can also consider other non-standard structural rules such as:

$$\begin{array}{cc} \text{Expansion (cf. [vB91]):} & \text{Mingle (cf. [OM64]):} \\ \frac{\Gamma, \alpha, \Delta \Rightarrow \gamma}{\Gamma, \alpha, \alpha, \Delta \Rightarrow \gamma} \mathbf{exp} & \frac{\Gamma, \Sigma, \Delta \Rightarrow \gamma \quad \Gamma, \Theta, \Delta \Rightarrow \gamma}{\Gamma, \Sigma, \Theta, \Delta \Rightarrow \gamma} \mathbf{min} \end{array}$$

(See also [HOS94, Kam02] for a detailed account.) Among them, some are harmless but others cause failure of cut elimination. In fact, the availability of cut elimination is very sensitive to the choice of structural rules:

- In general, sequent calculi with Contraction but without Exchange do not enjoy cut elimination. One way to recover cut elimination is to generalize Contraction to the one for *sequences* of formulas:

$$\frac{\Gamma, \Sigma, \Sigma, \Delta \Rightarrow \gamma}{\Gamma, \Sigma, \Delta \Rightarrow \gamma} \mathbf{seq-c}$$

- Expansion and Mingle are derivable from each other. However, Mingle admits cut elimination whereas Expansion does not.

In view of these intricacies, it is natural to look for some general criteria for a set of structural rules to admit cut elimination. The aim of this paper is to give such a criterion for cut elimination by using algebraic semantics.

We consider (the 0-free fragment of) *full Lambek calculus* ($\mathbf{FL}^+$, [Ono90, Ono94, Ono03]), i.e., intuitionistic logic without any structural rules, as our basic framework. We then introduce structural rules on $\mathbf{FL}^+$ in a general format. Residuated lattices are the algebraic structures corresponding to $\mathbf{FL}^+$ (see [JT02, Ono03]). In this setting, we introduce a criterion, called the propagation property, that can be stated both in syntactic and algebraic terminologies. It is a refinement of Girard's *naturality test*, which appears in an informal discussion in Appendix C.4 of [Gir99].

We then show that, for any set $\mathcal{R}$ of structural rules, the cut elimination theorem holds for $\mathbf{FL}^+$ enriched with $\mathcal{R}$ if and only if $\mathcal{R}$ satisfies the propagation property. To show the 'if' direction, the *phase structures* ([Abr90, Tro92, Ono94]) as well as Okada's cut elimination technique [Oka96, Oka99, Oka02] are essentially used.

As an application, we show that any set $\mathcal{R}$ of structural rules can be "completed" into another set $\mathcal{R}^*$, so that the cut elimination theorem holds for $\mathbf{FL}^+$ enriched with $\mathcal{R}^*$, while the provability remains the same.

2 Full Lambek Calculus and Structural Rules

The *formulas* of $\mathbf{FL}^+$ are built from propositional variables $a, b, c, \dots$ and constants 1 (unit), $\top$ (true) and $\perp$ (false) by using binary logical connectives $\cdot$ (fusion), $\backslash$ (right implication), $/$ (left implication), $\wedge$ (conjunction) and $\vee$ (disjunction). The set of formulas is denoted by $\mathcal{F}$. Small Greek letters $\alpha, \beta, \dots$ range over $\mathcal{F}$. For simplicity, we do not consider negation nor 0 in this paper. We use $\rightarrow$ as synonym for $\backslash$.

A *sequent* of $\mathbf{FL}^+$ is of the form $\alpha_1, \dots, \alpha_n \Rightarrow \beta$. Here, formulas $\alpha_1, \dots, \alpha_n$ are called *antecedents* and β is called a *succedent*. In the sequel, $\Gamma, \Delta, \dots$ stand for finite sequences of formulas, and $\emptyset$ stands for the empty sequence.

A sequent $\Gamma \Rightarrow \alpha$ is said to be *provable* in $\mathbf{FL}^+$ if it is derivable by using the inference rules in Figure 1. A formula α is *provable* if the sequent $\Rightarrow \alpha$ is provable. Given a (possibly infinite) set Ω of sequents, a sequent $\Gamma \Rightarrow \gamma$ is said to be *deducible* from Ω if $\Gamma \Rightarrow \gamma$ is provable in $\mathbf{FL}^+$ enriched with the additional axioms Ω (see [Ono94, Ono03] for more information).

$\frac{\Gamma \Rightarrow \alpha \quad \Delta_1, \alpha, \Delta_2 \Rightarrow \gamma}{\Delta_1, \Gamma, \Delta_2 \Rightarrow \gamma} \text{ cut}$	$\frac{}{\alpha \Rightarrow \alpha} \text{ init}$	$\frac{}{\Rightarrow 1} 1r$
$\frac{\Gamma_1, \alpha, \beta, \Gamma_2 \Rightarrow \gamma}{\Gamma_1, \alpha \cdot \beta, \Gamma_2 \Rightarrow \gamma} \cdot l$	$\frac{\Gamma \Rightarrow \alpha \quad \Delta \Rightarrow \beta}{\Gamma, \Delta \Rightarrow \alpha \cdot \beta} \cdot r$	$\frac{\Gamma_1, \Gamma_2 \Rightarrow \delta}{\Gamma_1, 1, \Gamma_2 \Rightarrow \delta} 1l$
$\frac{\Gamma \Rightarrow \alpha \quad \Delta_1, \beta, \Delta_2 \Rightarrow \delta}{\Delta_1, \Gamma, \alpha \setminus \beta, \Delta_2 \Rightarrow \delta} \setminus l$	$\frac{\alpha, \Gamma \Rightarrow \beta}{\Gamma \Rightarrow \alpha \setminus \beta} \setminus r$	$\frac{}{\Gamma_1, \perp, \Gamma_2 \Rightarrow C} \perp l$
$\frac{\Gamma \Rightarrow \alpha \quad \Delta_1, \beta, \Delta_2 \Rightarrow \delta}{\Delta_1, \beta / \alpha, \Gamma, \Delta_2 \Rightarrow \delta} / l$	$\frac{\Gamma, \alpha \Rightarrow \beta}{\Gamma \Rightarrow \beta / \alpha} / r$	$\frac{}{\Gamma \Rightarrow \top} \top r$
$\frac{\Gamma_1, \alpha, \Gamma_2 \Rightarrow \delta \quad \Gamma_1, \beta, \Gamma_2 \Rightarrow \delta}{\Gamma_1, \alpha \vee \beta, \Gamma_2 \Rightarrow \delta} \vee l$	$\frac{\Gamma \Rightarrow \alpha}{\Gamma \Rightarrow \alpha \vee \beta} \vee r_1$	$\frac{\Gamma \Rightarrow \beta}{\Gamma \Rightarrow \alpha \vee \beta} \vee r_2$
$\frac{\Gamma_1, \alpha, \Gamma_2 \Rightarrow \delta}{\Gamma_1, \alpha \wedge \beta, \Gamma_2 \Rightarrow \delta} \wedge l_1$	$\frac{\Gamma_1, \beta, \Gamma_2 \Rightarrow \delta}{\Gamma_1, \alpha \wedge \beta, \Gamma_2 \Rightarrow \delta} \wedge l_2$	$\frac{\Gamma \Rightarrow \alpha \quad \Gamma \Rightarrow \beta}{\Gamma \Rightarrow \alpha \wedge \beta} \wedge r$

图 1: Inference Rules of $\mathbf{FL}^+$

When it is necessary to indicate variables $a_1, \dots, a_m$ that might possibly occur in a formula α , we shall use the notation $\alpha[a_1, \dots, a_m]$, or $\alpha[\vec{a}]$ for short. The formula obtained from $\alpha[a_1, \dots, a_m]$ by substituting β_i for each a_i is denoted by $\alpha[\beta_1, \dots, \beta_m]$, or $\alpha[\vec{\beta}]$. Similar notation is used for sequences of formulas (and structural rules introduced below).

For $\Sigma \equiv \alpha_1, \dots, \alpha_n$ ($n \geq 1$), we define

$$\begin{aligned} * \Sigma &\equiv \alpha_1 \cdots \alpha_n, \\ \bigvee \Sigma &\equiv \alpha_1 \vee \cdots \vee \alpha_n. \end{aligned}$$

$\mathbf{FL}^+$ is entirely free from structural rules. Various systems of so-called sub-structural logics are obtained by enriching it with a suitable set of structural rules. Formally, a *structural rule* R is an $n + 1$ tuple $(\Theta_1; \dots; \Theta_n \triangleright \Theta_0)$, where $n \geq 1$ and each Θ_i is a finite sequence of variables, that satisfies the following condition:

(*) any variable occurring in $\Theta_1, \dots, \Theta_n$ also occurs in Θ_0 .

The last condition will be referred to as the *non-erasing condition*.

Let $R[\vec{a}]$ be a structural rule $(\Theta_1[\vec{a}]; \dots; \Theta_n[\vec{a}] \triangleright \Theta_0[\vec{a}])$, and $\vec{\beta}$ be a sequence of formulas. Then the result of substitution $R[\vec{\beta}] = (\Theta_1[\vec{\beta}]; \dots; \Theta_n[\vec{\beta}] \triangleright$

$\Theta_0[\vec{\beta}]$), is called an *instance* of R . When Φ is a set of formulas and formulas $\vec{\beta}$ belong to Φ , $R[\vec{\beta}]$ is called a Φ -*instance*. Each instance $R[\vec{\beta}]$ codifies an inference scheme of the form:

$$\frac{\Gamma, \Theta_1[\vec{\beta}], \Delta \Rightarrow \gamma \quad \dots \quad \Gamma, \Theta_n[\vec{\beta}], \Delta \Rightarrow \gamma}{\Gamma, \Theta_0[\vec{\beta}], \Delta \Rightarrow \gamma}$$

with Γ, Δ and γ arbitrary.

For example, the structural rules mentioned in the introduction can be formally specified as follows:

- **e**: $(a, b \triangleright b, a)$
- **w**: $(\emptyset \triangleright a)$
- **c**: $(a, a \triangleright a)$
- **exp**: $(a \triangleright a, a)$
- **min**: $\{(a_1, \dots, a_k; b_1, \dots, b_l \triangleright a_1, \dots, a_k, b_1, \dots, b_l) \mid 1 \leq k, 1 \leq l\}$
- **seq-c**: $\{(a_1, \dots, a_k, a_1, \dots, a_k \triangleright a_1, \dots, a_k) \mid 1 \leq k\}$

Notice that **min** and **seq-c** are specified by a countable set of structural rules.

Given a set $\mathcal{R}$ of structural rules, the system $\mathbf{FL}^+(\mathcal{R})$ is defined to be $\mathbf{FL}^+$ enriched with all instances of the additional structural rules $\mathcal{R}$. For instance, $\mathbf{FL}^+(\{\mathbf{e}\})$ amounts to $\mathbf{FL}_{\mathbf{e}}^+$ (intuitionistic linear logic without modality), while $\mathbf{FL}^+(\{\mathbf{e}, \mathbf{w}, \mathbf{c}\})$ is nothing but intuitionistic logic.

Due to the non-erasing condition, our structural rules satisfy the following property: any formula occurring in the upper sequents of a structural rule also occurs in the lower sequent. It follows that the cut elimination theorem always implies the subformula property.

Given a sequent, the *positive subformulas* and *negative subformulas* are defined as usual. We then have:

Lemma 2.1 *Let $\mathcal{R}$ be a set of structural rules. Suppose that $\mathbf{FL}^+(\mathcal{R})$ enjoys cut elimination. Then it satisfies the (polarized) subformula property: if a sequent $\Gamma \Rightarrow \alpha$ is provable in $\mathbf{FL}^+(\mathcal{R})$, then it has a derivation π in which only subformulas of $\Gamma \Rightarrow \alpha$ occur. Moreover, any antecedent (succedent, resp.) formula of a sequent in π is a negative (positive, resp.) subformula of $\Gamma \Rightarrow \alpha$.*

To study the properties of structural rules, it is convenient to represent them as formulas. Given a structural rule $R = (\Theta_1; \dots; \Theta_n \triangleright \Theta_0)$, define its *formula representation* $\hat{R}$ by

$$\hat{R} \equiv * \Theta_0 \rightarrow (* \Theta_1 \vee \dots \vee * \Theta_n).$$

For instance, $\hat{\mathbf{e}} \equiv b \cdot a \rightarrow a \cdot b$ and $\hat{\mathbf{w}} \equiv a \rightarrow 1$. The formula representation of $\mathbf{min}_1 = (a; b \triangleright a, b)$ is $a \cdot b \rightarrow a \vee b$.

If R is of the form $R[a_1, \dots, a_m]$ and $\alpha_1, \dots, \alpha_m$ belong to a set Φ of formulas, then $\hat{R}[\alpha_1, \dots, \alpha_m]$ is called a Φ -*instance* of $\hat{R}$. When $\mathcal{R}$ is a set of structural rules, $\hat{\mathcal{R}}$ denotes the set $\{\hat{R} \mid R \in \mathcal{R}\}$.

As expected, there is an instance-wise correspondence between structural rules and their formula representations:

Lemma 2.2 *Let $R[\vec{a}]$ be a structural rule. Then an instance $R[\vec{\alpha}]$ is derivable from $\hat{R}[\vec{\alpha}]$ and vice versa.*

3 Syntactic Propagation

Let us now introduce a syntactic version of the propagation property. To motivate the notion, consider the contrast between $\mathbf{FL}^+(\{\mathbf{c}\})$ and $\mathbf{FL}^+(\mathbf{seq-c})$. As is mentioned in the introduction, the former does not enjoy cut elimination. For instance, the cut below cannot be eliminated:

$$\frac{\frac{\frac{\alpha \Rightarrow \alpha}{\alpha, \beta \Rightarrow \alpha \cdot \beta} \quad \frac{\beta \Rightarrow \beta}{\alpha, \beta \Rightarrow \alpha \cdot \beta}}{\alpha, \beta \Rightarrow \alpha \cdot \beta} \quad \frac{\frac{\frac{\alpha \cdot \beta \Rightarrow \alpha \cdot \beta}{\alpha \cdot \beta, \alpha \cdot \beta \Rightarrow (\alpha \cdot \beta) \cdot (\alpha \cdot \beta)} \quad \frac{\alpha \cdot \beta \Rightarrow \alpha \cdot \beta}{\alpha \cdot \beta \Rightarrow (\alpha \cdot \beta) \cdot (\alpha \cdot \beta)}}{\alpha \cdot \beta \Rightarrow (\alpha \cdot \beta) \cdot (\alpha \cdot \beta)} \mathbf{c}}{\alpha, \beta \Rightarrow (\alpha \cdot \beta) \cdot (\alpha \cdot \beta)} \mathbf{cut}$$

On the other hand, if $\mathbf{c}$ is generalized to $\mathbf{seq-c}$, the cut can be easily eliminated:

$$\frac{\frac{\frac{\alpha \Rightarrow \alpha}{\alpha, \beta \Rightarrow \alpha \cdot \beta} \quad \frac{\beta \Rightarrow \beta}{\alpha, \beta \Rightarrow \alpha \cdot \beta}}{\alpha, \beta, \alpha, \beta \Rightarrow (\alpha \cdot \beta) \cdot (\alpha \cdot \beta)} \mathbf{seq-c}}{\alpha, \beta \Rightarrow (\alpha \cdot \beta) \cdot (\alpha \cdot \beta)}$$

Now our question is this: what is the *essential* difference between $\mathbf{c}$ and $\mathbf{seq-c}$? A distinctive feature of $\mathbf{seq-c}$ is that it *propagates from variable instances to fusion instances*. Namely, a fusion instance $(a \cdot b, a \cdot b \triangleright a \cdot b)$ is derivable from a variable instance $(a, b, a, b \triangleright a, b)$ as follows:

$$\frac{\frac{\Gamma, a \cdot b, a \cdot b, \Delta \Rightarrow \gamma}{\Gamma, a, b, a, b, \Delta \Rightarrow \gamma} \text{seq-c}}{\Gamma, a, b, \Delta \Rightarrow \gamma} \frac{\Gamma, a, b, \Delta \Rightarrow \gamma}{\Gamma, a \cdot b, \Delta \Rightarrow \gamma}$$

(Pedantically speaking, an instance $R[\vec{\alpha}] = (\Theta_1[\vec{\alpha}]; \dots; \Theta_n[\vec{\alpha}] \triangleright \Theta_0[\vec{\alpha}])$ is *derivable* from a set Ω of instances of some structural rules if for arbitrary Γ, Δ and γ , the sequent $\Gamma, \Theta_0[\vec{\alpha}], \Delta \Rightarrow \gamma$ is deducible from the sequents $\Gamma, \Theta_i[\vec{\alpha}], \Delta \Rightarrow \gamma$ for $1 \leq i \leq n$ in $\mathbf{FL}^+$ enriched with the rule instances Ω .)

In contrast, one can observe that **c** does not propagate to fusion instances.

Next, consider the contrast between $\mathbf{FL}^+(\{\mathbf{exp}\})$ and $\mathbf{FL}^+(\mathbf{min})$. The former does not enjoy cut elimination, as witnessed by:

$$\frac{\frac{\beta \Rightarrow \beta}{\beta \Rightarrow \alpha \vee \beta} \quad \frac{\frac{\overline{\alpha \Rightarrow \alpha}}{\alpha \Rightarrow \alpha \vee \beta} \quad \frac{\overline{\alpha \vee \beta \Rightarrow \alpha \vee \beta}}{\alpha \vee \beta, \alpha \vee \beta \Rightarrow \alpha \vee \beta} \text{exp}}{\alpha, \alpha \vee \beta \Rightarrow \alpha \vee \beta} \text{cut} \quad \frac{\alpha, \alpha \vee \beta \Rightarrow \alpha \vee \beta}{\alpha, \beta \Rightarrow \alpha \vee \beta} \text{cut}$$

Notice that one cannot obtain a cut-free proof even if **exp** is generalized to a sequence version as above. On the other hand, when **exp** is replaced with **min**, a cut-free proof is obtained:

$$\frac{\frac{\overline{\alpha \Rightarrow \alpha}}{\alpha \Rightarrow \alpha \vee \beta} \quad \frac{\overline{\beta \Rightarrow \beta}}{\beta \Rightarrow \alpha \vee \beta}}{\alpha, \beta \Rightarrow \alpha \vee \beta} \text{min}$$

Therefore, we may again ask what is the *essential* difference between **exp** and **min**. This time, our answer is that **min** *propagates from variable instances to disjunction instances*. Namely, a disjunction instance $(a_1 \vee b_1; a_2 \vee b_2 \triangleright a_1 \vee b_1, a_2 \vee b_2)$ is derivable from variable instances $(a_1; a_2 \triangleright a_1, a_2)$, $(a_1; b_2 \triangleright a_1, b_2)$, $(b_1; a_2 \triangleright b_1, a_2)$ and $(b_1; b_2 \triangleright b_1, b_2)$ as follows:

$$\frac{\frac{\Gamma, a_1 \vee b_1, \Delta \Rightarrow \gamma}{\Gamma, a_1, \Delta \Rightarrow \gamma} \quad \frac{\Gamma, a_2 \vee b_2, \Delta \Rightarrow \gamma}{\Gamma, a_2, \Delta \Rightarrow \gamma} \text{min} \quad \dots \quad \frac{\Gamma, a_1 \vee b_1, \Delta \Rightarrow \gamma}{\Gamma, b_1, \Delta \Rightarrow \gamma} \quad \frac{\Gamma, a_2 \vee b_2, \Delta \Rightarrow \gamma}{\Gamma, b_2, \Delta \Rightarrow \gamma} \text{min}}{\Gamma, a_1, a_2, \Delta \Rightarrow \gamma} \text{min} \quad \dots \quad \frac{\Gamma, b_1, b_2, \Delta \Rightarrow \gamma}{\Gamma, a_1 \vee b_1, a_2 \vee b_2, \Delta \Rightarrow \gamma} \text{min}$$

In contrast, **exp** does not propagate to disjunction instances.

These observations bring us to the following definition. A set $\mathcal{R}$ of structural rules satisfies the *syntactic propagation property* if the following holds:

- For every $R[a_1, \dots, a_m] \in \mathcal{R}$ and every $\Sigma_1, \dots, \Sigma_m$, where each Σ_i is a sequence of variables, both $R[*\Sigma_1, \dots, *\Sigma_m]$ and $R[\vee \Sigma_1, \dots, \vee \Sigma_m]$

are derivable from the Φ -instances of the structural rules in $\mathcal{R}$, where Φ is the set of variables occurring in $\Sigma_1, \dots, \Sigma_m$.

In view of Lemma 2.2, this is equivalent to say that

- the formulas $\hat{R}[*\Sigma_1, \dots, *\Sigma_m]$ and $\hat{R}[\bigvee \Sigma_1, \dots, \bigvee \Sigma_m]$ are deducible from the Φ -instances of the formulas in $\hat{\mathcal{R}}$.

The syntactic propagation property does not explicitly refer to, but is actually closely related to cut elimination. In fact, we have:

Proposition 3.1 *Let $\mathcal{R}$ be a set of structural rules. If $\mathbf{FL}^+(\mathcal{R})$ enjoys cut elimination, then $\mathcal{R}$ satisfies the syntactic propagation property.*

4 Residuated lattices and semantic propagation

An algebra $\mathbf{P} = \langle P, \wedge, \vee, \cdot, \backslash, /, 1 \rangle$ is called a (*bounded*) *residuated lattice* if

1. $\langle P, \wedge, \vee \rangle$ is a lattice with the greatest element $\top$ and the least element $\perp$.
2. $\langle P, \cdot, 1 \rangle$ is a monoid.
3. The operations $\backslash$ and $/$ are right and left residuals of $\cdot$. Namely, for any $x, y, z \in P$,

$$x \cdot y \leq z \iff x \leq z/y \iff y \leq x \backslash z.$$

(See [JT02, Ono03] for general introductions to residuated lattices.)

A *valuation* f on $\mathbf{P}$ maps each variable to an element of P . Given a set $X \subseteq P$, f is called an *X-valuation* if the range is a subset of X . As usual, f can be extended to a map from the formulas $\mathcal{F}$ to P as follows:

$$\begin{aligned} f(\dagger) &= \dagger && \text{for } \dagger \in \{\top, \perp, 1\}, \\ f(\alpha \star \beta) &= f(\alpha) \star f(\beta) && \text{for } \star \in \{\wedge, \vee, \cdot, \backslash, /\}. \end{aligned}$$

A formula α is said to be *true* under valuation f in $\mathbf{P}$ if $f(\alpha) \geq 1$. In particular, $\alpha \rightarrow \beta$, i.e., $\alpha \backslash \beta$ is true iff $f(\alpha) \leq f(\beta)$. A formula α is *valid* (*X-valid*, resp.) in $\mathbf{P}$ if it is true under all valuations (*X-valuations*, resp.) on $\mathbf{P}$.

The residuated lattices are algebraic models of $\mathbf{FL}^+$. In particular, the following strong form of soundness holds for them:

Lemma 4.1 *Let $\mathbf{P}$ be a residuated lattice and f be a valuation on it. If α is deducible from Φ and all formulas in Φ are true under f in $\mathbf{P}$, then α is also true under f .*

Given a set $\mathcal{R}$ of structural rules, an $\mathcal{R}$ -*residuated lattice* is a residuated lattice in which all formulas in $\widehat{\mathcal{R}}$ are valid. By the previous lemma, any formula provable in $\mathbf{FL}^+(\mathcal{R})$ is valid in all $\mathcal{R}$ -residuated lattices.

Coming back to the residuated lattices in general, we may observe that the monoid multiplication $\cdot$ is *continuous* in the following sense:

Lemma 4.2 *Let $q_0, \dots, q_m \in P$ and let*

$$\delta(p_1, \dots, p_m) = q_0 \cdot p_1 \cdot q_1 \cdots q_{m-1} \cdot p_m \cdot q_m,$$

for any $p_1, \dots, p_m \in P$. Let also $\tilde{\delta}(p) = \delta(p, \dots, p)$. Suppose that X is a subset of P for which $\bigvee X$ exists. We then have:

$$\tilde{\delta}(\bigvee X) = \bigvee_{Y \subseteq_{fin} X} \tilde{\delta}(\bigvee Y),$$

where $Y \subseteq_{fin} X$ holds iff Y is a finite subset of X .

Given $X \subseteq P$, the *multiplication closure* $\prod(X)$, the *join closure* $\coprod(X)$ and the *finite join closure* $\coprod_{fin}(X)$ are defined by

$$\begin{aligned} \prod(X) &= \{p_1 \cdots p_n \mid n \geq 0, p_1, \dots, p_n \in X\}, \\ \coprod(X) &= \{\bigvee Y \mid Y \subseteq X, \bigvee Y \text{ exists}\}, \\ \coprod_{fin}(X) &= \{\bigvee Y \mid Y \subseteq_{fin} X\}. \end{aligned}$$

A set $\mathcal{R}$ of structural rules satisfies the *semantic propagation property* if for any residuated lattice $\mathbf{P}$ and $X \subseteq P$, the following holds:

- if all formulas in $\widehat{\mathcal{R}}$ are X -valid, then they are also $\prod(\prod(X))$ -valid.

We have:

Proposition 4.3 *If a set $\mathcal{R}$ of structural rules satisfies the syntactic propagation property, it also satisfies the semantic propagation property.*

5 Phase structures and semantic cut elimination

We now introduce a special class of residuated lattices, sometimes called (intuitionistic noncommutative) *phase structures* (see [Abr90, Tro92, Ono94]). Let $\mathbf{M} = \langle M, \cdot, 1 \rangle$ be a monoid. Denote the powerset of M by $\wp(M)$, and define for $X, Y \in \wp(M)$,

$$X \bullet Y = \{x \cdot y \mid x \in X, y \in Y\}.$$

A function $C : \wp(M) \rightarrow \wp(M)$ is said to be a *closure operator* on $\wp(M)$ if for all $X, Y \in \wp(M)$,

1. $X \subseteq C(X)$,
2. $C(C(X)) \subseteq C(X)$,
3. $X \subseteq Y$ implies $C(X) \subseteq C(Y)$,
4. $C(X) \bullet C(Y) \subseteq C(X \bullet Y)$.

A set $X \in \wp(M)$ is *closed* if $X = C(X)$. The set of all closed sets in $\wp(M)$ is denoted by $\mathcal{C}_M$. Define for any closed sets $X, Y \in \mathcal{C}_M$ and for any family $\mathcal{X}$ of closed sets,

$$\begin{aligned} X \cup_C Y &= C(X \cup Y), \\ \bigcup_C \mathcal{X} &= C(\bigcup \mathcal{X}), \\ X \bullet_C Y &= C(X \bullet Y), \\ X \parallel Y &= \{y \mid \forall x \in X, x \cdot y \in Y\}, \\ Y \parallel X &= \{y \mid \forall x \in X, y \cdot x \in Y\}. \end{aligned}$$

We then have:

Lemma 5.1 *If $\mathbf{M}$ is a monoid and C is a closure operator on $\wp(M)$, then the algebra*

$$\mathbf{C}_M = \langle \mathcal{C}_M, \cap, \cup_C, \bullet_C, \parallel, \parallel, C(\{1\}) \rangle,$$

is a complete residuated lattice with infinite join $\bigcup_C$.

In every phase structure, the following hold:

1. $C(\{x \cdot y\}) = C(\{x\}) \bullet_C C(\{y\})$ for any $x, y \in M$,
2. $C(X) = \bigcup_C_{x \in X} C(\{x\})$ for any $X \subseteq M$.

As a consequence, phase structures satisfy the following remarkable property which plays a key role in connecting the semantic propagation property to cut elimination:

Lemma 5.2 *Suppose that $\mathbf{M}$ is finitely generated by a set A , i.e., any element x of M can be written as $y_1 \cdots y_n$ for some $y_1, \dots, y_n \in A$. Let $C'_A = \{C(\{y\}) \mid y \in A\}$. Then we have $C_M = \coprod (\prod (C'_A))$.*

We now describe a specific construction of a phase structure due to [Oka96, Oka99] (and slightly remedied by [OT99]), which is quite useful for proving the cut elimination theorem. (See also [BOJ01], where Okada's construction is reformulated as algebraic *quasi-completion* and *quasi-embedding*.)

Let $\mathcal{F}^*$ be the free monoid generated by the formulas $\mathcal{F}$ of $\mathbf{FL}^+$; the elements of $\mathcal{F}^*$ are sequences of formulas, the monoid multiplication is concatenation, and the unit element is the empty sequence $\emptyset$.

Let us fix a set $\mathcal{R}$ of structural rules. The operator C is defined on the basis of *cut-free* provability in $\mathbf{FL}^+(\mathcal{R})$:

$$\begin{aligned} \llbracket \Gamma _ \Delta \Rightarrow \gamma \rrbracket &= \{ \Sigma \mid \Gamma, \Sigma, \Delta \Rightarrow \gamma \text{ is cut-free provable in } \mathbf{FL}^+(\mathcal{R}) \}, \\ \mathcal{D} &= \{ \llbracket \Gamma _ \Delta \Rightarrow \gamma \rrbracket \mid \Gamma, \Delta, \gamma \text{ arbitrary} \}, \\ C(X) &= \bigcap_{X \subseteq Y \in \mathcal{D}} Y. \end{aligned}$$

Then one can show that C is indeed a closure operator on $\wp(\mathcal{F}^*)$ (for an arbitrary $\mathcal{R}$). Hence by Lemma 5.1, the algebra

$$\mathbf{C}_{\mathcal{F}^*} = \langle \mathcal{C}_{\mathcal{F}^*}, \cap, \cup_C, \bullet_C, \backslash, /, C(\{\emptyset\}) \rangle$$

is a residuated lattice.

Let f_0 be a valuation on $\mathbf{C}_{\mathcal{F}^*}$ defined by $f_0(a) = C(\{a\})$. In this setting, we have *Okada's lemma*:

Lemma 5.3 *For every formula α , $\alpha \in f_0(\alpha) \subseteq \llbracket _ \Rightarrow \alpha \rrbracket$. In particular, for every sequent $\Gamma \Rightarrow \alpha$, if $(*\Gamma) \rightarrow \alpha$ is true under f_0 , then $\Gamma \Rightarrow \alpha$ is cut-free provable in $\mathbf{FL}^+(\mathcal{R})$.*

It is worth noting that Okada's lemma holds independently of which structural rules $\mathcal{R}$ we adopt. It only concerns with the properties of logical inference rules. What depends on the choice of $\mathcal{R}$ is the following:

Lemma 5.4 *If $\mathcal{R}$ satisfies the semantic propagation property, then $\mathbf{C}_{\mathcal{F}^*}$ is an $\mathcal{R}$ -residuated lattice.*

We have thus arrived at:

Proposition 5.5 *If $\mathcal{R}$ satisfies the semantic propagation property, then $\mathbf{FL}^+(\mathcal{R})$ enjoys cut elimination.*

By putting Propositions 3.1, 4.3 and 5.5 together, we obtain our main theorem:

Theorem 5.6 *Let $\mathcal{R}$ be a set of structural rules. Then the following are equivalent:*

1. $\mathbf{FL}^+(\mathcal{R})$ enjoys cut elimination.
2. $\mathcal{R}$ satisfies the syntactic propagation property.
3. $\mathcal{R}$ satisfies the semantic propagation property.

6 Completion of Structural Rules

Recall that Contraction **c** can be generalized to its sequence version **seq-c** without changing provability so that the cut elimination theorem holds for $\mathbf{FL}^+(\mathbf{seq-c})$. We say that **c** can be *completed* into **seq-c**. Likewise, Expansion **exp** can be completed into Mingle **min**. The completion techniques implicitly used there are by no means specific to **c** and **exp**. In fact, we can show that an arbitrary set of structural rules can be completed by using those techniques.

Theorem 6.1 *Given a set $\mathcal{R}$ of structural rules, one can obtain another set $\mathcal{R}^*$ of structural rules such that the following hold.*

- $\mathbf{FL}^+(\mathcal{R})$ and $\mathbf{FL}^+(\mathcal{R}^*)$ are equivalent.
- $\mathcal{R}^*$ satisfies the syntactic propagation property. Hence $\mathbf{FL}^+(\mathcal{R}^*)$ enjoys cut-elimination.

To prove this, we use our characterization of cut elimination by the syntactic propagation property.

Acknowledgements. We are indebted to Jean-Yves Girard, who suggested to the author a possible linkage between his test and cut elimination in 1999, and thus motivated the current work. Our thanks are also due to Shun'ichi Amano, Nicolas Galatos, Makoto Kanazawa, Hiroakira Ono, Takafumi Sakurai, Kentaro Sato and Hiroki Takamura for various comments and stimulating discussions.

参考文献

- [Abr90] V. Michele Abrusci. Non-commutative intuitionistic linear propositional logic. *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik*, 36:297–318, 1990.
- [BOJ01] F. Belardinelli, H. Ono, and P. Jipsen. Algebraic aspects of cut elimination. *Studia Logica*, 68:1–32, 2001.
- [De93] K. Došen and P. Schröder-Heister (eds). *Substructural Logics*. Oxford University Press, 1993.
- [GO] N. Galatos and H. Ono. Substructural logics over FL I: Algebraization, parametrized local deduction theorem and interpolation. Draft.
- [Gir87] J.-Y. Girard. Linear logic. *Theoretical Computer Science*, 50:1–102, 1987.
- [Gir95] J.-Y. Girard. Linear logic: Its syntax and semantics. In J.-Y. Girard, Y. Lafont, and L. Regnier, editors, *Advances in Linear Logic*, pages 1–42. Cambridge University Press, 1995. Proceedings of the Workshop on Linear Logic, Ithaca, New York, June 1993.
- [Gir99] J.-Y. Girard. On the meaning of logical rules I: syntax vs. semantics. In U. Berger and H. Schwichtenberg, editors, *Computational Logic*, pages 215 – 272. Heidelberg Springer-Verlag, 1999.
- [HOS94] R. Hori, Hiroakira Ono, and Harold Schellinx. Extending intuitionistic linear logic with knotted structural rules. *Notre-Dame Journal of Formal Logic*, 35(2):219–242, 1994.
- [JT02] J. Jipsen and C. Tsinakis. A survey of residuated lattices. In J. Martinez, editor, *Ordered Algebraic Structures*, pages 19–56. Kluwer Academic Publishes, 2002.
- [Kam02] N. Kamide. Substructural logics with mingle. *Journal of Logic, Language and Information*, 11(2):227–249, 2002.
- [OM64] M. Ohnishi and K. Matsumoto. A system for strict implication. *Annals of the Japan Association for Philosophy of Science*, 2:183–188, 1964.

- [Oka96] M. Okada. Phase semantics for higher order completeness, cut-elimination and normalization proofs (extended abstract). In J.-Y. Girard, M. Okada, and A. Scedrov, editors, *ENTCS (Electronic Notes in Theoretical Computer Science) Vol.3: A Special Issue on the Linear Logic 96, Tokyo Meeting*. Elsevier-ENTCS, 1996.
- [Oka99] M. Okada. Phase semantic cut-elimination and normalization proofs of first- and higher-order linear logic. *Theoretical Computer Science*, 227:333–396, 1999.
- [Oka02] M. Okada. A uniform semantic proof for cut-elimination and completeness of various first and higher order logics. *Theoretical Computer Science*, 281:471–498, 2002.
- [OT99] M. Okada and K. Terui. The finite model property for various fragments of intuitionistic linear logic. *Journal of Symbolic Logic*, 64(2):790–802, 1999.
- [Ono90] H. Ono. Structural rules and a logical hierarchy. In P. P. Petkov, editor, *Mathematical Logic*, pages 95–104. Plenum Press, 1990. Proceedings of the Summer School and Conference on Mathematical Logic, honourably dedicated to the 90th Anniversary of Arend Heyting (1898–1980), Chaika, Bulgaria, 1988.
- [Ono94] H. Ono. Semantics for substructural logics. In K. Došen and P. Schröder-Heister, editors, *Substructural logics*, pages 259–291. Oxford University Press, 1994.
- [Ono98] H. Ono. Proof-theoretic methods for nonclassical logic – an introduction. In M. Takahashi, M. Okada, and M. Dezani-Ciancaglini, editors, *Theories of Types and Proofs*, pages 207–254. MSJ Memoirs vol.2, Mathematical Society of Japan, 1998.
- [Ono03] H. Ono. Substructural logics and residuated lattices — an introduction. *Trends in Logic*, 20:177–212, 2003.
- [Ter05] K. Terui. Which structural rules admit cut elimination? — An algebraic criterion. Submitted, 2005. Available at <http://research.nii.ac.jp/~terui>.
- [Tro92] A. S. Troelstra. *Lectures on Linear Logic*. CSLI Lecture Notes 29, Center for the Study of Language and Information, Stanford, California, 1992.

- [vB91] J. van Benthem. *Language in Action: Categories, Lambdas and Dynamic Logic*. (Studies in Logic 130). North-Holland, Amsterdam, 1991.