

REVIEW PAPER

The molecular architecture of the plant nuclear pore complex

Kentaro Tamura and Ikuko Hara-Nishimura*

Department of Botany, Graduate School of Science, Kyoto University, Kyoto 606
8502, Japan

*To whom correspondence should be addressed.

E-mail: ihnishi@gr.bot.kyoto-u.ac.jp

Running title: Plant NPC structure

Abbreviations: FG, phenylalanine-glycine; Nup, nucleoporin; NPC, nuclear pore
complex

21 **Abstract**

22 The nucleus contains the cell's genetic material, which directs cellular activity via
23 gene regulation. The physical barrier of the nuclear envelope needs to be permeable
24 to a variety of macromolecules and signals. The most prominent gateways for the
25 transport of macromolecules are the nuclear pore complexes (NPCs). The NPC is
26 the largest multiprotein complex in the cell, and is composed of multiple copies of
27 approximately 30 different proteins called nucleoporins. Although much progress has
28 been made in dissecting the NPC structure in vertebrates and yeast, the molecular
29 architecture and physiological function of nucleoporins in plants remains poorly
30 understood. In this review, we summarize the current knowledge regarding the plant
31 NPC proteome and address structural and functional aspects of plant nucleoporins,
32 which support the fundamental cellular machinery.

33

34

Ultrastructure of the plant NPC

Because the nuclear pore complexes (NPCs) are the largest macromolecular complexes in cells, early studies of the complexes in plants were performed using electron microscopy. More than 40 years ago, Yoo and Bayley (1967) reported that nuclear pores in the pea plant resembled those described in various animal cells, and comprised 1 to 3 central granules (or possibly tubules), which were surrounded by an annulus. It was estimated that nuclear pores occupied a maximum of 38% of the nuclear envelope area. Roberts and Northcote (1970) used a freeze-etch technique to reveal the high-resolution structure of NPCs in sycamore and bean. They showed that, as in other organisms, the plant NPC is octagonally symmetrical around its cylindrical axis. They also determined that the plant NPC ($1150 \times 640 \text{ \AA}$) is larger than its yeast counterpart ($960 \times 350 \text{ \AA}$), but smaller than that in vertebrates ($1450 \times 800 \text{ \AA}$). These studies provided the first model of higher plant NPCs, which was based on observation of vertical sections and a rough three-dimensional structure.

Fiserova *et al.* (2009) used an in-lens field emission scanning electron microscope (feSEM), a type of high-resolution SEM, to gain further insight into NPC structure. Using cultured tobacco cells, the NPC structure of the nuclear membrane was visualized on both the cytoplasmic and nucleoplasmic sides. They clearly demonstrated that, in both logarithmic and stationary phase cells, NPCs are non-randomly distributed over the nuclear envelope (as previously observed in other higher eukaryotes). The density of the NPCs was largely unchanged during cell growth, but NPCs became predominantly organized into rows in stationary phase cells. Interestingly, different NPC conformations, which were related to different cell stages, were observed. Logarithmic phase cells, which are metabolically active and undergo rapid cell division, contained NPCs with a larger inner pore diameter, which may be capable of rapid and effective transport. By contrast, stationary phase and senescent cells contained NPCs with smaller inner pore diameters; internal filaments

62 were observed within the pores, which emerged from the base of each subunit and
63 were directed toward the NPC center. These results suggest that the NPC may
64 differentially regulate transport activity and specificity by changing its component
65 parts. Importantly, such a conformational change was also observed in *Xenopus*
66 (Goldberg *et al.*, 1997) and *Drosophila* (Kiseleva *et al.*, 2001), suggesting that the
67 mechanisms underlying NPC differentiation are conserved across eukaryotes.

68 Direct involvement of the inner nuclear envelope protein in NPC anchoring
69 and positioning was suggested by studies in vertebrate cells (Lenz-Bohme *et al.*,
70 1997; Liu *et al.*, 2000; Maeshima *et al.*, 2006). The distribution of NPCs on the
71 nuclear envelope in vertebrates correlates with the distribution of lamins. NPCs in
72 tobacco are closely linked to a filamentous structure on the inner nuclear membrane
73 (Fiserova *et al.*, 2009; Fiserova and Goldberg, 2010), and the organization and
74 dimensions of these filaments resemble the arrangement of the nuclear lamina in
75 *Xenopus* oocytes (Goldberg *et al.*, 2008). Although no lamin homologues have been
76 identified in plants, the plant NPC might be anchored on the nuclear envelope in the
77 same way as in vertebrates. Nuclear matrix constituent protein1 (NMCP1), which is a
78 long coiled-coil protein localized at the nuclear rim, is considered to be the best
79 candidate of plant lamin-like protein (Masuda *et al.*, 1997; Boruc *et al.*, 2012).
80 Mutants of two NMCP1 homologues in *Arabidopsis*, *little nuclei1* (*linc1*) and *linc2*,
81 show reduced nuclear size and an altered nuclear structure (Dittmer *et al.*, 2007).
82 Thus, NMCP1/LINC is thought to determine nuclear organization in plants (Dittmer *et al.*,
83 2007). Moreover, *Arabidopsis* SUN (Sad-1/UNC-84)-domain proteins, which are
84 inner nuclear envelope proteins, have been isolated and characterized (Graumann
85 *et al.*, 2010; Oda and Fukuda, 2011). A mammal SUN-domain protein is known to
86 interact with the NPC and likely regulates NPC distribution across the nuclear
87 surface (Liu *et al.*, 2007). It will be necessary to determine how plant NPCs interact
88 with these proteins on the inner nuclear envelope if we are to better understand NPC
89 function, positioning, assembly, and disassembly.

90

91 **Attempts to identify NPC components using proteomics**

92 *Yeast and vertebrate NPC proteomes*

93 The first comprehensive proteomics study of NPCs was performed in yeast
94 (*Saccharomyces cerevisiae*) (Rout *et al.*, 2000). There are several advantages to
95 working with yeast nuclei: yeast have the highest NPC/nuclear volume ratio of any
96 organism (Maul, 1977) and, unlike the nuclei in mammalian cells, they do not have a
97 lamina connecting the NPCs to other structures and/or protein complexes. Rout and
98 Blobel (1993) prepared a highly enriched NPC fraction from yeast spheroplasts after
99 several rounds of sucrose density gradient centrifugation. The fraction was then
100 subjected to three different HPLC separation techniques followed by SDS-PAGE to
101 identify the individual proteins associated with the NPC (Rout *et al.*, 2000).
102 Matrix-associated laser desorption ionization time-of-flight (MALDI-TOF) and
103 MALDI-ion trap tandem mass spectrometry identified a total of 174 proteins, of which
104 34 were previously uncharacterized ORFs. The uncharacterized ORFs and putative
105 nucleoporins were epitope-tagged and their subcellular localizations were analyzed
106 by immunofluorescence and immunoelectron microscopy to determine the position
107 and stoichiometry of each nucleoporin within the NPC. In all, 29 nucleoporins and 11
108 NPC-associated proteins, which serve as transport factors, were identified.

109 Two years after this yeast NPC proteomics study, Cronshaw *et al.* (2002)
110 reported the first vertebrate NPC proteomics data. They developed a fractionation
111 procedure that yielded highly enriched nuclear envelopes from rat liver nuclei. After
112 removing the chromatin, the nuclear membranes and their associated proteins were
113 extracted by incubation in Triton X-100 and SDS. This procedure yielded intact NPCs
114 embedded in the lamina. Then, a zwitterionic detergent was used to specifically
115 solubilize the NPCs and release the monomeric nucleoporins. The solubilized
116 proteins were separated by HPLC and SDS-PAGE and analyzed by using both

MALDI-quadrupole-quadrupole time-of-flight (MALDI-QqTOF) and MALDI-ion trap spectrometry. The uncharacterized proteins were then expressed as GFP fusions in HeLa cells to investigate their localization. This work identified and classified 29 nucleoporins and 18 NPC-associated proteins (Cronshaw *et al.*, 2002).

The plant NPC proteome

Knowledge of the individual components and overall structure of NPCs in plants largely lagged behind that of vertebrate and yeast NPCs. With a few exceptions, nucleoporin homologues could not be identified in plants using homology-based approaches (Meier, 2006). Several nuclear proteomics studies were reported, but they only identified a few nucleoporins (Pendle *et al.*, 2005; Aki and Yanagisawa, 2009). It was largely unknown which nucleoporins comprised plant NPCs. Because a protocol for the biochemical isolation of plant NPCs has not been developed, it is difficult to perform studies similar to those done in yeast and vertebrates. To overcome these problems, interactive proteomics was used to identify plant NPC components (Tamura *et al.*, 2010). Transgenic plants were generated, which expressed GFP-tagged mRNA export factor1 (RAE1), a known nucleoporin in *Arabidopsis*. The nucleoporins were then purified by immunoprecipitating the extracts from transgenic plants with an anti-GFP antibody. A linear ion trap mass spectrometer (LTQ-Orbitrap) was used to identify a total of 200 proteins in the immunoprecipitates. By comparing these with a database containing metazoan nucleoporins and performing expression studies of GFP fusions, 24 proteins were classified as nucleoporins (Tamura *et al.*, 2010).

To obtain more information about plant NPCs, Tamura *et al.* (2010) selected other nucleoporins identified in the RAE1-GFP immunoprecipitates and used them as bait for further rounds of NPC purification. Finally, five cycles of interactive proteomic analysis were performed. This procedure identified at least 30 putative

nucleoporins, 22 of which had not been previously annotated. This work also demonstrated that the interactive proteomic approach is a very powerful technique, which can be used to comprehensively identify the individual components of macromolecular complexes in plants.

Domain architecture

Proteomic analysis and x-ray crystallography revealed the detailed protein structure of individual nucleoporins. It was estimated that 38% of all nucleoporin amino acid residues contain an α -solenoid fold, 29% contain Phe-Gly repeats (FG repeats), and 16% contain β -propeller folds. Other individual fold types accounted for less than 5% of the total nucleoporin pool (Devos *et al.*, 2006). The small number of predicted fold types within the nucleoporin proteins and their similar internal symmetries suggest that the bulk of the NPC structures evolved through a series of gene duplications and divergences from a simple precursor set of only a few proteins. The predicted structure of individual *Arabidopsis* nucleoporins are summarised in Fig. 1.

Phe-Gly repeats

There are an estimated 128 FG domains, which harbour thousands of total FG repeats, within any given yeast NPC (Rout *et al.*, 2000). The FG repeats interact with nuclear transport receptors, providing a selective barrier to the diffusion of macromolecules (Radu *et al.*, 1995; Patel *et al.*, 2007). The FG repeats appear to be localized toward the inside of the NPC (Rout *et al.*, 2000). It is therefore reasonable to suggest that nucleoporins rich in FG repeats (FG nucleoporins) coat the central pore surface, providing interaction domains for transport receptors within the central pore. Several studies investigated how these FG repeats function as a transport barrier. It appears that the FG repeats are intrinsically unfolded, and contain short

clusters of hydrophobic amino acids (e.g., FXFG or GLFG) separated by hydrophilic spacers. Because of flexibility of FG-repeat domains, it was proposed that phenylalanine-mediated inter-repeat interactions cross-linked the FG repeat domains into elastic and reversible hydrogels (Frey *et al.*, 2006). The hydrogel-like properties of the nuclear pores were thought to function as a molecular sieve. Systematic and large-scale analyses of the FG nucleoporins in yeast revealed that two forms are present in NPCs (Patel *et al.*, 2007). FG nucleoporins anchored at the NPC centre (central FG nucleoporins) form a cohesive meshwork of filaments via hydrophobic interactions. By contrast, FG nucleoporins anchored at the NPC periphery (cytoplasmic and nucleoplasmic FG nucleoporins) are generally noncohesive. Therefore, a two-gate model of NPC architecture is proposed, which comprises a central diffusion gate formed by a meshwork of cohesive FG nucleoporins and a peripheral gate formed by repulsive FG nucleoporins (Patel *et al.*, 2007; Strambio-De-Castillia *et al.*, 2010).

Coiled coils

The nuclear basket within the NPC is generally constructed from large coiled-coil proteins: NUCLEAR PORE ANCHOR (NUA) in plants (Xu *et al.*, 2007a), MYOSIN-LIKE PROTEIN1/2 (Mlp1/2) in yeast (Rout *et al.*, 2000), and TRANSLOCATED PROMOTER REGION (Tpr) in vertebrates (Cordes *et al.*, 1997). Coiled coils mediate protein-protein interactions, implying that the nuclear basket serves as a recruitment platform that brings various factors together within the nucleus. The yeast FG-nucleoporin complex, Nsp1-Nup82-Nup49 complex (Nup62-Nup58-Nup54 complex in plants), is held together by coiled-coil interactions (Bailer *et al.*, 2001) and tethered to the NPC scaffold via the N-terminal coiled-coil region of NUCLEOPORIN-INTERACTING COMPONENT OF 96 kDa (Nic96). The FG nucleoporins in plants also possess a coiled-coil domain and are thus assumed

also to form a complex (Nup62-Nup58-Nup54 complex). However, Nup93, a plant homologue of yeast Nic96, has no obvious coiled-coil motif (Tamura *et al.*, 2010), implying that the plant FG-nucleoporin complex is tethered to the NPC scaffold via other interactions.

α -solenoids

The α -solenoid fold comprises a two- or three- α helix unit, which is repeatedly stacked to form an elongated domain with the N- and C-termini at opposite ends of the molecule (Brohawn *et al.*, 2009). Approximately one-third of nucleoporins contain α -solenoid domains (Devos *et al.*, 2006). The outer and inner rings (scaffold nucleoporins) are dominated by an evenly distributed meshwork of α -solenoid domains, which is expected to facilitate the formation of a flexible fold. This allows large conformational changes without breaking protein-protein interactions, accommodates nucleocytoplasmic transport cargoes of different sizes, and promotes malleability of the nuclear envelope (Alber *et al.*, 2007a). α -solenoid folds are also common in large protein assemblies, such as those found in clathrin-, COPI-, and COPII-coated vesicles (Devos *et al.*, 2004). In coated vesicles, clathrin-like and adaptor proteins form the structural scaffold of the protein coat, which surrounds the membrane of the vesicle in a lattice-like fashion (Fotin *et al.*, 2004a; Fotin *et al.*, 2004b); a role that maybe similar to that played by the scaffold nucleoporins in the NPC. Therefore, it was proposed that the core structures of NPCs and vesicle-coating complexes have a common evolutionary origin (Devos *et al.*, 2004; Devos *et al.*, 2006).

β -propellers

The β -propeller is a disc-like structure assembled from structural modules, known as

223 blades, circularly-arranged around a central channel (Chen *et al.*, 2011). In general,
224 each blade consists of 4–10 repeats of a four-stranded antiparallel β -sheet motif. The
225 central channel of the propeller fold is usually funnel-like in shape, with a wider
226 bottom opening, which serves as entry point to the active site. A set of nucleoporins
227 was initially identified as β -propellers based on sequence analysis. In yeast, only
228 SECRETORY13 (SEC13) and SEC13 HOMOLOGUE1 (SEH1) contain the signature
229 WD-40 repeat motif, and were among the very first β -propellers to be identified (Pryer
230 *et al.*, 1993). Since then, other nucleoporins containing WD-40 repeats have been
231 identified as β -propeller nucleoporins. Berke *et al.* (2004) experimentally determined
232 the crystal structure of the N-terminal domain of human Nup133. They revealed that
233 it consists of a seven-bladed β -propeller, which provides a molecular platform that
234 mediates multiple interactions with other proteins (Berke *et al.*, 2004). After this
235 structure was solved, additional noncanonical β -propeller domains within the NPC
236 were identified. The β -propellers are widely assumed to serve as protein-protein
237 interaction sites (Chen *et al.*, 2011). Peripheral β -propeller domains function to recruit
238 accessory proteins, whereas those located centrally within the NPC bind
239 subcomplexes (Strambio-De-Castillia *et al.*, 2010; Grossman *et al.*, 2012).

240 The α -solenoid and β -propeller nucleoporins are thought to act as the
241 structural scaffold of the NPC. Both the α -solenoid and the β -propeller folds provide
242 extensive solvent-accessible surfaces, which appear well-suited for binding other
243 proteins. Moreover, the α -solenoid and β -propeller folds seem to be quite robust and
244 can tolerate significant variation in amino acid sequence while still retaining the core
245 structure, thereby allowing optimization interactions with many different partners
246 (Devos *et al.*, 2006).

248 **NPC architecture**

249 Electron microscopy studies have dissected the NPC structure (Callan and Tomlin,

1950; Yoo and Bayley, 1967; Roberts and Northcote, 1970). These studies show that the general morphology of the NPC is conserved among eukaryotes. Using a different approach, Alber *et al.* (2007) performed a computational analysis to determine a detailed architectural map of the yeast NPC. They combined a diverse set of immunoelectron microscopic, crystallographic, and proteomic data to generate the map with an estimated 5 nm resolution, which agreed with a large body of complementary data for both vertebrates and yeast (Alber *et al.*, 2007a; Alber *et al.*, 2007b; Brohawn *et al.*, 2009). Although the primary sequence homology between nucleoporins from different model organisms is low, the high conservation of the overall shape and predicted fold types suggests that this NPC map can be applied to the NPCs of other organisms. Therefore, we compared a dataset derived from each plant nucleoporin with those from yeast and vertebrate nucleoporins and generated an architectural map of plant NPCs (Fig. 2). According to this model, nucleoporins can be subdivided into five classes: transmembrane ring, core scaffold (inner ring, outer ring, and linker), cytoplasmic filaments, nuclear basket, and central FG (Alber *et al.*, 2007a; Alber *et al.*, 2007b; Brohawn *et al.*, 2009; Grossman *et al.*, 2012).

Transmembrane ring nucleoporins

Transmembrane nucleoporins are thought to anchor the NPC to the pore membrane and bind the assembled complex to the nuclear envelope. In plants, two transmembrane nucleoporins, GLYCOPROTEIN OF 210 kDa (Gp210) (Pom152 in yeast) (Gerace *et al.*, 1982; Greber *et al.*, 1990) and NUCLEAR DIVISION CYCLE1 (NDC1) (Wozniak *et al.*, 1994), constitute an outer transmembrane ring. In addition to these proteins, yeast and vertebrates possess their own unique membrane proteins, PORE MEMBRANE PROTEIN OF 34 kDa (Pom34) (Rout *et al.*, 2000) and Pom121 (Hallberg *et al.*, 1993), respectively. In vertebrates, Gp210 serves a fundamental role in NPC disassembly and is phosphorylated during nuclear envelope breakdown

(Galy *et al.*, 2008). An *Arabidopsis gp210* knockout mutant was identified as an embryo defective mutant (EMB3012) (<http://www.seedgenes.org/index.html>) (Meinke *et al.*, 2008), which arrested at the proglobular stage during embryogenesis. This suggests that plant Gp210 plays an essential role in embryonic development. Human NDC1 is involved in NPC assembly and is required for targeting the heritable disease-associated nucleoporin, Aladin, to the NPC (Kind *et al.*, 2009; Yamazumi *et al.*, 2009). It would be interesting to determine whether a plant Aladin also tethers to the NPC via NDC1.

Cytoplasmic filaments and the nuclear basket

Two characteristic peripheral NPC components, the cytoplasmic filaments and the nuclear basket, are localized asymmetrically within the cytoplasm and nucleoplasm of the NPC, respectively. These structures play a role in specific interactions that can serve as docking sites for transport complexes at both cytoplasmic and nucleoplasmic sides. The cytoplasmic filaments are composed primarily of two FG nucleoporins, Nup214 and CG1 (Kraemer *et al.*, 1995; Rout *et al.*, 2000). Although these nucleoporins are not well-characterized in plants, those in yeast provide sites at which mRNA export factors can mature messenger ribonucleoprotein particles (Folkmann *et al.*, 2011). The biggest nucleoporin in vertebrates (Nup358) is localized in the cytoplasmic filaments, and tethers RanGAP (RanGTPase-activating protein) to the NPC to facilitate transportin-dependent nuclear import (Hutten *et al.*, 2009). By contrast, plants and yeast have no Nup358 homologues in their genomes (Grossman *et al.*, 2012). Plants use the WIP-WIT complex (discussed below) to anchor RanGAP on the nuclear envelope (Xu *et al.*, 2007b; Zhao *et al.*, 2008), whereas yeast RanGAP is localized to the cytosol (Hopper *et al.*, 1990).

The nuclear basket is composed of eight elongated filaments, which protrude ~60–80 nm from the nuclear face of the NPC into the nucleoplasm and

converge on a distal ring structure (Strambio-De-Castillia *et al.*, 2010). Each nuclear basket consists of one nucleoporin (NUA in plants), which has long coiled coil domains, and two FG nucleoporins (Nup136/Nup1 and Nup50 in plants). Human Tpr (NUA in plants) is thought to constitute the central architectural element that forms the scaffold of the nuclear basket, whereas Nup153 (Nup136/Nup1 in plants) binds to the nuclear coaxial ring linking the NPC core structures to Tpr (Krull *et al.*, 2004). It is reported that both *Arabidopsis* NUA and Nup136 are involved in mRNA export (Xu *et al.*, 2007a; Tamura *et al.*, 2010), as suggested by a study of vertebrate Tpr (Shibata *et al.*, 2002). Importantly, *Arabidopsis nua* and *nup136* mutant plants display similar phenotypes, including early flowering and low fertility, indicating that NUA and Nup136 function together during plant development. In *Xenopus*, Nup153 (Nup136/Nup1 in plants) is the only nucleoporin that interacts with lamin B, which supports the nuclear structure. Although no lamin homologue has been identified in plants, it is reported that *Arabidopsis* nuclei can change shape, depending on the amount of accumulated Nup136 (Tamura *et al.*, 2010; Tamura and Hara-Nishimura, 2011). This implies that Nup136/Nup1 is the nucleoporin that links the NPC to the nuclear lamina in plants.

Central FG nucleoporins

Central FG nucleoporins, also known as barrier nucleoporins, show a transverse orientation with respect to the central channel of the NPC and form a selective barrier that mediates nucleocytoplasmic transport. Native FG repeat regions have an unfolded structure, allowing multiple low-affinity and high-specificity interactions with transport factors (Strawn *et al.*, 2004). In fact, *in vitro* studies show that all FG nucleoporins appear to have the ability to interact with at least one transport receptor (Ryan and Wentz, 2000). Bayliss *et al.* (2000) determined the crystal structure of the human transport receptor, importin β , and identified its two FG repeat binding sites. It

is estimated that there are approximately 160 transport factor binding sites within each NPC (Grossman *et al.*, 2012), which allow the simultaneous binding of multiple transport factors within a single NPC.

Five central FG nucleoporins (Nup98, Nup62, Nup58, Nup54, and Nup35) have been identified in plants and are well conserved in other organisms. Of these, only Nup62 has been characterized in *Arabidopsis* (Zhao and Meier, 2011). *Arabidopsis* Nup62 interacts with nuclear transport factor2 (NTF2), as previously observed in yeast (Nsp1) (Clarkson *et al.*, 1997) and vertebrates (Nup62) (Percipalle *et al.*, 1997). Knockdown of Nup62 transcripts results in a severe dwarf phenotype and early flowering, indicating an important function of Nup62 at different stages of plant development. Further analysis of central FG nucleoporins will be required to identify how the different FG nucleoporins function during the plant life cycle. This will increase our understanding of the specific interactions between transport factors and NPCs in plants.

Outer ring, inner ring, and linker nucleoporins (scaffold nucleoporins)

Scaffold nucleoporins are composed of three groups of nucleoporin subcomplexes: the outer ring, the inner ring, and the linker. They connect membrane nucleoporins to central FG nucleoporins thereby bridging the anchoring transmembrane layer and the barrier layer of the NPC. Because scaffold nucleoporins form the rigid skeleton of the NPC, most remain incorporated within the NPC during the entire life of a cell (D'Angelo *et al.*, 2009). Scaffold nucleoporins are thought to play a key role in maintaining the stability of the nuclear envelope by ensuring co-planarity of the outer and inner surfaces (Alber *et al.*, 2007a). Therefore, it is proposed that a major function of the outer rings is to facilitate the smooth transition of the pore membrane into the inner and outer nuclear envelopes (Alber *et al.*, 2007a).

The largest and most evolutionarily conserved subcomplex, comprising

several nucleoporins, is located at the outer rings and is known as the Nup107/160 subcomplex in plants and vertebrates and the Nup84 subcomplex in yeast. In plants, it is composed of eight nucleoporins: Nup160, Nup133, Nup107, Nup96, Nup85, Nup43, SEC13, and SEH1 (Fig. 2) (Xu and Meier 2007; Tamura *et al.*, 2010; Wiermer *et al.*, 2012). Electron microscopic and biochemical analyses revealed that the yeast Nup84 subcomplex has a 40-nm long, Y-shaped, triskelion-like structure (Siniosoglou *et al.*, 2000; Kampmann and Blobel, 2009). This structure has two hinge regions that are conformationally flexible, which allows the NPC scaffold component to change its structure to enable large cargoes to pass through the central channel (Kampmann and Blobel, 2009). The scaffold nucleoporins located in the central part of the main channel are called inner ring nucleoporins and linker nucleoporins. In plants, the inner ring consists of Nup205, Nup188, Nup155, and Nup35 (Fig. 2) (Tamura *et al.*, 2010). Similar to the outer ring nucleoporins, they contain α -solenoid and β -propeller folds and exhibit the typical structural scaffolding motif. The linker nucleoporins are attached between the outer and inner rings (Alber *et al.*, 2007a), and include Nup93 and Nup88. They act as a bridge between the core scaffold and the FG nucleoporins.

Genetic studies have characterized several plant scaffold nucleoporin mutants. *Arabidopsis nup96* and *nup88* mutants were isolated during a genetic screen aimed at identifying downstream components responsible for resistance (R) protein activation (Zhang and Li, 2005; Cheng *et al.*, 2009). Both mutants showed deficiencies in innate immunity and, possibly, an impairment in nuclear import of proteins, which is involved in the plant response to pathogens. Furthermore, Wiermer *et al.* (2012) isolated knockout mutants for each component of the Nup107/160 complex and investigated plant immune responses. Of the seven mutated nucleoporins, only *nup160* and *seh1* caused impaired immune responses. These results suggest that Nup160, Nup96, and SEH1 within the Nup107/160 subcomplex are important for defense signaling (Wiermer *et al.*, 2012). The other Nup107/160

subcomplex proteins, Nup133 and Nup75, are required for fungal and rhizobial colonization in *Lotus japonicus* (Kanamori *et al.*, 2006; Saito *et al.*, 2007). Moreover, *Arabidopsis nup160* and *nup96* mutants show altered hormonal and temperature responses and an early flowering phenotype (Dong *et al.*, 2006; Parry *et al.*, 2006). Further work is required to identify the underlying molecular mechanisms and functions of the plant Nup107/160 subcomplex in various signaling pathways.

NPC-associated proteins

TREX-2

The TREX-2 (transcription-coupled export 2) complex comprises SUPPRESSOR OF ACTIN3 (SAC3), Tho2/Hpr1 PHENOTYPE1 (THP1), SL GENE UPSTREAM OF ySa1 (SUS1), and CELL DIVISION CYCLE31 (CDC31) proteins, and is involved in mRNA export in yeast (Kohler and Hurt, 2007). The TREX-2 complex anchors at the inner side of the NPC via Nup1 (Nup136/Nup1 in plants) and Nup60 (Nup50 in plants) (Fischer *et al.*, 2002). SUS1 also interacts with the SAGA (Spt, Ada, Gcn5 and acetyltransferase histone acetyltransferase) complex, a large transcription initiation complex that catalyzes histone acetylation and de-ubiquitylation. The TREX-2 complex is thought to functionally couple SAGA-dependent gene expression to mRNA export on the inner side of the NPC (Rodriguez-Navarro *et al.*, 2004). Lu *et al.* (2010) identified TREX2 components in *Arabidopsis*, including THP1, SUS1, SAC3s, and CDC31s, and showed that the Nup136/Nup1-THP1 interaction links the TREX-2 complex to the nuclear basket (Fig. 3). Using yeast two-hybrid assays and a bimolecular fluorescence complementation assay, they also showed that THP1 interacts with DSS1, a subunit of the 26S proteasome regulatory particle (Lu *et al.*, 2010). Consistent with this finding, yeast SDD1 was identified as a functional component of the TREX-2 complex (Mannen *et al.*, 2008), and is required for nuclear export of specific sets of mRNA. These results suggest that plants utilize

an evolutionarily conserved system for mRNA export through the NPC.

ESD4

In *Arabidopsis*, EARLY IN SHORT DAYS4 (ESD4) functionally interacts with NUA, a major component of the nuclear basket (Xu *et al.*, 2007a). Genetic analysis indicates that NUA and ESD4 might act via a shared pathway involved in various aspects of plant development. ESD4 encodes a SUMO (small ubiquitin-related modifier conjugates)-specific protease and is localized on the nuclear envelope (Murtas *et al.*, 2003) (Fig. 3). In both *esd4* and *nua* mutants, the accumulation of SUMO conjugates increases, suggesting that ESD4 and NUA are involved in SUMO homeostasis and in regulating nucleocytoplasmic transport in plants. The nuclear basket components, Mlp1/2 in yeast and Nup153 in humans, tether the SUMO-conjugating enzymes, UBIQUITIN-LIKE PROTEIN1 (Ulp1) and SENTRIN-SPECIFIC PROTEASE2 (SEN2), respectively (Zhang *et al.*, 2002; Panse *et al.*, 2003). The yeast *ulp1* mutant displays altered protein sumoylation patterns and increased pre-mRNA leakage into the cytoplasm. These results suggest that desumoylation within the NPC might be a key regulatory event that prevents inappropriate pre-mRNA export in different species (Lewis *et al.*, 2007).

WPP DOMAIN-INTERACTING PROTEIN (WIP) and WPP DOMAIN-INTERACTING TAIL-ANCHORED PROTEIN (WIT) complexes

In vertebrates, RanGAP, which plays an important role in nucleocytoplasmic transport, is anchored to the NPC via Nup358 (Strambio-De-Castillia *et al.*, 2010). This interaction is mediated by a unique C-terminal domain within RanGAP, and is dependent upon sumoylation. By contrast, plant NPCs lack a Nup358 homologue, and all known plant RanGAPs contain a unique N-terminal domain called the WPP

domain (a highly conserved Trp-Pro-Pro motif), which is necessary and sufficient for nuclear envelope targeting (Meier, 2000). Two types of plant-specific nuclear envelope proteins, WIPs and WITs, were isolated and identified as RanGAP-anchoring proteins (Xu *et al.*, 2007b; Zhao *et al.*, 2008). Although there is no direct evidence that WIP and WIT proteins physically interact with the NPC, it is assumed that they have a functional connection (Fig. 3). Recently, *Arabidopsis* WIP was reported to function as a KASH (Klarsicht/ANC-1/Syne Homology)-domain protein, which interacts with a SUN-domain protein (Zhou *et al.*, 2012). This SUN-KASH bridge is necessary for maintaining the elongated nuclear shape of epidermal cells, indicating that WIPs are versatile and play different roles.

Conclusion

Since the identification of the plant NPC proteome, much progress has been made in clarifying the molecular components and architecture of the NPC. Despite conserved functional similarities between NPCs from plants and other organisms, structural differences exist. It is necessary to determine how these structural variations and protein sequence differences contribute to the functional organization of the plant NPC. Clearly, reverse genetic studies and high-resolution imaging techniques are needed if we are to understand the function and structure of each nucleoporin in detail.

References

- Aki T, Yanagisawa S.** 2009. Application of rice nuclear proteome analysis to the identification of evolutionarily conserved and glucose-responsive nuclear proteins. *J Proteome Res* **8**, 3912-3924.
- Alber F, Dokudovskaya S, Veenhoff LM, Zhang W, Kipper J, Devos D, Suprpto A, Karni-Schmidt O, Williams R, Chait BT, Sali A, Rout MP.** 2007a. The molecular architecture of the nuclear pore complex. *Nature* **450**, 695-701.
- Alber F, Dokudovskaya S, Veenhoff LM, Zhang W, Kipper J, Devos D, Suprpto A, Karni-Schmidt O, Williams R, Chait BT, Rout MP, Sali A.** 2007b. Determining the architectures of macromolecular assemblies. *Nature* **450**, 683-694.
- Bailer SM, Balduf C, Hurt E.** 2001. The Nsp1p carboxy-terminal domain is organized into functionally distinct coiled-coil regions required for assembly of nucleoporin subcomplexes and nucleocytoplasmic transport. *Mol Cell Biol* **21**, 7944-7955.
- Bayliss R, Littlewood T, Stewart M.** 2000. Structural basis for the interaction between FxFG nucleoporin repeats and importin-beta in nuclear trafficking. *Cell* **102**, 99-108.
- Berke IC, Boehmer T, Blobel G, Schwartz TU.** 2004. Structural and functional analysis of Nup133 domains reveals modular building blocks of the nuclear pore complex. *J Cell Biol* **167**, 591-597.
- Boruc J, Zhou X, Meier I.** 2012 Dynamics of the plant nuclear envelope and nuclear pore. *Plant Physiol* **158**. 78-86.
- Brohawn SG, Partridge JR, Whittle JR, Schwartz TU.** 2009. The nuclear pore complex has entered the atomic age. *Structure* **17**, 1156-1168.

- Callan HG, Tomlin SG.** 1950. Experimental studies on amphibian oocyte nuclei. I. Investigation of the structure of the nuclear membrane by means of the electron microscope. *Proc R Soc Lond B Biol Sci* **137**, 367-378.
- Chen CK, Chan NL, Wang AH.** 2011. The many blades of the beta-propeller proteins: conserved but versatile. *Trends Biochem Sci* **36**, 553-561.
- Cheng YT, Germain H, Wiermer M, Bi D, Xu F, Garcia AV, Wirthmueller L, Despres C, Parker JE, Zhang Y, Li X.** 2009. Nuclear pore complex component MOS7/Nup88 is required for innate immunity and nuclear accumulation of defense regulators in Arabidopsis. *Plant Cell* **21**, 2503-2516.
- Clarkson WD, Corbett AH, Paschal BM, Kent HM, McCoy AJ, Gerace L, Silver PA, Stewart M.** 1997. Nuclear protein import is decreased by engineered mutants of nuclear transport factor 2 (NTF2) that do not bind GDP-Ran. *J Mol Biol* **272**, 716-730.
- Cordes VC, Reidenbach S, Racjwitz HR, Franke WW.** 1997. Identification of protein p270/Tpr as a constitutive component of the nucleoporecomplex-attachedintranuclear laments. *J Cell Biol* **136**, 515-529.
- Cronshaw JM, Krutchinsky AN, Zhang W, Chait BT, Matunis MJ.** 2002. Proteomic analysis of the mammalian nuclear pore complex. *J Cell Biol* **158**, 915-927.
- D'Angelo MA, Raices M, Panowski SH, Hetzer MW.** 2009. Age-dependent deterioration of nuclear pore complexes causes a loss of nuclear integrity in postmitotic cells. *Cell* **136**, 284-295.
- Devos D, Dokudovskaya S, Alber F, Williams R, Chait BT, Sali A, Rout MP.** 2004. Components of coated vesicles and nuclear pore complexes share a common molecular architecture. *PLoS Biol* **2**, e380.
- Devos D, Dokudovskaya S, Williams R, Alber F, Eswar N, Chait BT, Rout MP, Sali**

- A.** 2006. Simple fold composition and modular architecture of the nuclear pore complex. *Proc Natl Acad Sci U S A* **103**, 2172-2177.
- Dittmer TA, Stacey NJ, Sugimoto-Shirasu K, Richards EJ.** 2007 LITTLE NUCLEI genes affecting nuclear morphology in *Arabidopsis thaliana*. *Plant Cell* **19**, 2793–2803
- Dong CH, Hu X, Tang W, Zheng X, Kim YS, Lee BH, Zhu JK.** 2006. A putative *Arabidopsis* nucleoporin, AtNUP160, is critical for RNA export and required for plant tolerance to cold stress. *Mol Cell Biol* **26**, 9533-9543.
- Fischer T, Strasser K, Racz A, Rodriguez-Navarro S, Oppizzi M, Ihrig P, Lechner J, Hurt E.** 2002. The mRNA export machinery requires the novel Sac3p-Thp1p complex to dock at the nucleoplasmic entrance of the nuclear pores. *Embo J* **21**, 5843-5852.
- Fiserova J, Goldberg MW.** 2010. Relationships at the nuclear envelope: lamins and nuclear pore complexes in animals and plants. *Biochem Soc Trans* **38**, 829-831.
- Fiserova J, Kiseleva E, Goldberg MW.** 2009. Nuclear envelope and nuclear pore complex structure and organization in tobacco BY-2 cells. *Plant J* **59**, 243-255.
- Folkmann AW, Noble KN, Cole CN, Wentz SR.** 2011. Dbp5, Gle1-IP6 and Nup159: a working model for mRNP export. *Nucleus* **2**, 540-548.
- Fotin A, Cheng Y, Grigorieff N, Walz T, Harrison SC, Kirchhausen T.** 2004a. Structure of an auxilin-bound clathrin coat and its implications for the mechanism of uncoating. *Nature* **432**, 649-653.
- Fotin A, Cheng Y, Sliz P, Grigorieff N, Harrison SC, Kirchhausen T, Walz T.** 2004b. Molecular model for a complete clathrin lattice from electron cryomicroscopy. *Nature* **432**, 573-579.
- Frey S, Richter RP, Gorlich D.** 2006. FG-rich repeats of nuclear pore proteins form a

- three-dimensional meshwork with hydrogel-like properties. *Science* **314**, 815-817.
- Galy V, Antonin W, Jaedicke A, Sachse M, Santarella R, Haselmann U, Mattaj J.** 2008. A role for gp210 in mitotic nuclear-envelope breakdown. *J Cell Sci* **121**, 317-328.
- Gerace L, Ottaviano Y, Kondor-Koch C.** 1982. Identification of a major polypeptide of the nuclear pore complex. *J Cell Biol* **95**, 826-837.
- Goldberg MW, Wiese C, Allen TD, Wilson KL.** 1997. Dimples, pores, star-rings, and thin rings on growing nuclear envelopes: evidence for structural intermediates in nuclear pore complex assembly. *J Cell Sci* **110**, 409-420.
- Goldberg MW, Huttenlauch I, Hutchison CJ, Stick R.** 2008. Filaments made from A- and B-type lamins differ in structure and organization. *J Cell Sci* **121**, 215-225.
- Graumann K, Runions J, Evans DE.** 2010 Characterization of SUN-domain proteins at the higher plant nuclear envelope. *Plant J* **61**, 134–144
- Greber UF, Senior A, Gerace L.** 1990. A major glycoprotein of the nuclear pore complex is a membrane-spanning polypeptide with a large luminal domain and a small cytoplasmic tail. *EMBO J.* **9**, 1495-1502.
- Grossman E, Medalia O, Zwerger M.** 2012. Functional architecture of the nuclear pore complex. *Annu Rev Biophys* **41**, 557-584.
- Hallberg E, Wozniak RW, Blobel G.** 1993. An integral membrane protein of the pore membrane domain of the nuclear envelope contains a nucleoporin-like region. *J Cell Biol* **122**, 513-521.
- Hopper AK, Traglia HM, Dunst RW.** 1990. The yeast RNA1 gene product necessary for RNA processing is located in the cytosol and apparently excluded from the nucleus. *J Cell Biol* **111**, 309-321.

- Hutten S, Walde S, Spillner C, Hauber J, Kehlenbach RH.** 2009. The nuclear pore component Nup358 promotes transportin-dependent nuclear import. *J Cell Sci* **122**, 1100-1110.
- Kampmann M, Blobel G.** 2009. Three-dimensional structure and flexibility of a membrane-coating module of the nuclear pore complex. *Nat Struct Mol Biol* **16**, 782-788.
- Kanamori N, Madsen LH, Radutoiu S, Frantescu M, Quistgaard EM, Miwa H, Downie JA, James EK, Felle HH, Haaning LL, Jensen TH, Sato S, Nakamura Y, Tabata S, Sandal N, Stougaard J.** 2006. A nucleoporin is required for induction of Ca²⁺ spiking in legume nodule development and essential for rhizobial and fungal symbiosis. *Proc Natl Acad Sci U S A* **103**, 359-364.
- Kind B, Koehler K, Lorenz M, Huebner A.** 2009. The nuclear pore complex protein ALADIN is anchored via NDC1 but not via POM121 and GP210 in the nuclear envelope. *Biochem Biophys Res Commun* **390**, 205-210.
- Kiseleva E, Rutherford S, Cotter LM, Allen TD, Goldberg MW.** 2001. Steps of nuclear pore complex disassembly and reassembly during mitosis in early *Drosophila* embryos. *J Cell Sci* **114**, 3607-3618.
- Kohler A, Hurt E.** 2007. Exporting RNA from the nucleus to the cytoplasm. *Nat Rev Mol Cell Biol* **8**, 761-773.
- Kraemer DM, Strambio-de-Castillia C, Blobel G, Rout MP.** 1995. The essential yeast nucleoporin NUP159 is located on the cytoplasmic side of the nuclear pore complex and serves in karyopherin-mediated binding of transport substrate. *J Biol Chem* **270**, 19017–19021.
- Krull S, Thyberg J, Bjorkroth B, Rackwitz HR, Cordes VC.** 2004. Nucleoporins as components of the nuclear pore complex core structure and Tpr as the architectural element of the nuclear basket. *Mol Biol Cell* **15**, 4261-4277.

- Lenz-Bohme B, Wismar J, Fuchs S, Reifegerste R, Buchner E, Betz H, Schmitt B.** 1997. Insertional mutation of the *Drosophila* nuclear lamin Dm0 gene results in defective nuclear envelopes, clustering of nuclear pore complexes, and accumulation of annulate lamellae. *J Cell Biol* **137**, 1001-1016.
- Lewis A, Felberbaum R, Hochstrasser M.** 2007. A nuclear envelope protein linking nuclear pore basket assembly, SUMO protease regulation, and mRNA surveillance. *J Cell Biol* **178**, 813-827.
- Liu J, Rolef Ben-Shahar T, Riemer D, Treinin M, Spann P, Weber K, Fire A, Gruenbaum Y.** 2000. Essential roles for *Caenorhabditis elegans* lamin gene in nuclear organization, cell cycle progression, and spatial organization of nuclear pore complexes. *Mol Biol Cell* **11**, 3937-3947.
- Liu Q, Pante N, Misteli T, Elsagga M, Crisp M, Hodzic D, Burke B, Roux KJ.** 2007. Functional association of Sun1 with nuclear pore complexes. *J Cell Biol* **178**, 785–798.
- Lu Q, Tang X, Tian G, Wang F, Liu K, Nguyen V, Kohalmi SE, Keller WA, Tsang EW, Harada JJ, Rothstein SJ, Cui Y.** 2010. Arabidopsis homolog of the yeast TREX-2 mRNA export complex: components and anchoring nucleoporin. *Plant J* **61**, 259-270.
- Maeshima K, Yahata K, Sasaki Y, Nakatomi R, Tachibana T, Hashikawa T, Imamoto F, Imamoto N.** 2006. Cell-cycle-dependent dynamics of nuclear pores: pore-free islands and lamins. *J Cell Sci* **119**, 4442-4451.
- Mannen T, Andoh T, Tani T.** 2008. Dss1 associating with the proteasome functions in selective nuclear mRNA export in yeast. *Biochem Biophys Res Commun* **365**, 664-671.
- Masuda K, Xu ZJ, Takahashi S, Ito A, Ono M, Nomura K, Inoue M.** 1997. Peripheral framework of carrot cell nucleus contains a novel protein predicted to exhibit a

- long alpha-helical domain. *Exp Cell Res* **232**, 173–181.
- Maul GG.** 1977. Nuclear pore complexes. Elimination and reconstruction during mitosis. *J Cell Biol* **74**, 492-500.
- Meier I.** 2000. A novel link between ran signal transduction and nuclear envelope proteins in plants. *Plant Physiol* **124**, 1507-1510.
- Meier I.** 2006. Composition of the plant nuclear envelope: theme and variations. *Journal of Experimental Botany* **58**, 27-34.
- Meinke D, Muralla R, Sweeney C, Dickerman A.** 2008. Identifying essential genes in *Arabidopsis thaliana*. *Trends Plant Sci* **13**, 483-491.
- Murtas G, Reeves PH, Fu Y-F, Bancroft I, Dean C, Coupland G.** 2003. A nuclear protease required for flowering-time regulation in *Arabidopsis* reduces the abundance of SMALL UBIQUITIN-RELATED MODIFIER conjugates. *Plant Cell* **15**, 2308-2319.
- Oda Y, Fukuda H.** 2011 Dynamics of *Arabidopsis* SUN proteins during mitosis and their involvement in nuclear shaping. *Plant J* **66**, 629–641
- Panse VG, Kuster B, Gerstberger T, Hurt E.** 2003. Unconventional tethering of Ulp1 to the transport channel of the nuclear pore complex by karyopherins. *Nat Cell Biol* **5**, 21-27.
- Parry G, Ward S, Cernac A, Dharmasiri S, Estelle M.** 2006. The *Arabidopsis* SUPPRESSOR OF AUXIN RESISTANCE proteins are nucleoporins with an important role in hormone signaling and development. *Plant Cell* **18**, 1590-1603.
- Patel S, Rose A, Meulia T, Dixit R, Cyr RJ, Meier I.** 2004. *Arabidopsis* WPP-domain proteins are developmentally associated with the nuclear envelope and promote cell division. *Plant Cell* **16**, 3260-3273.
- Patel SS, Belmont BJ, Sante JM, Rexach MF.** 2007. Natively unfolded nucleoporins

- gate protein diffusion across the nuclear pore complex. *Cell* **129**, 83-96.
- Pendle AF, Clark GP, Boon R, Lewandowska D, Lam YW, Andersen J, Mann M, Lamond AI, Brown JWS, Shaw PJ.** 2005. Proteomic analysis of the Arabidopsis nucleolus suggests novel nucleolar functions. *Mol Biol Cell* **16**, 260-269.
- Percipalle P, Clarkson WD, Kent HM, Rhodes D, Stewart M.** 1997. Molecular interactions between the importin alpha/beta heterodimer and proteins involved in vertebrate nuclear protein import. *J Mol Biol* **266**, 722-732.
- Pryer NK, Salama NR, Schekman R, Kaiser CA.** 1993. Cytosolic Sec13p complex is required for vesicle formation from the endoplasmic reticulum in vitro. *J Cell Biol* **120**, 865-875.
- Radu A, Moore MS, Blobel G.** 1995. The peptide repeat domain of nucleoporin Nup98 functions as a docking site in transport across the nuclear pore complex. *Cell* **81**, 215-222.
- Roberts K, Northcote DH.** 1970. Structure of the nuclear pore in higher plants. *Nature* **228**, 385-386.
- Rodriguez-Navarro S, Fischer T, Luo MJ, Antunez O, Brettschneider S, Lechner J, Perez-Ortin JE, Reed R, Hurt E.** 2004. Sus1, a functional component of the SAGA histone acetylase complex and the nuclear pore-associated mRNA export machinery. *Cell* **116**, 75-86.
- Rout MP, Blobel G.** 1993. Isolation of the yeast nuclear pore complex. *J Cell Biol* **123**, 771-783.
- Rout MP, Aitchison JD, Suprpto A, Hjertaas K, Zhao Y, Chait BT.** 2000. The yeast nuclear pore complex: composition, architecture, and transport mechanism. *J Cell Biol* **148**, 635-651.

- Ryan KJ, Wente SR.** 2000. The nuclear pore complex: a protein machine bridging the nucleus and cytoplasm. *Curr Opin Cell Biol* **12**, 361-371.
- Saito K, Yoshikawa M, Yano K, Miwa H, Uchida H, Asamizu E, Sato S, Tabata S, Imaizumi-Anraku H, Umehara Y, Kouchi H, Murooka Y, Szczylowski K, Downie JA, Parniske M, Hayashi M, Kawaguchi M.** 2007. NUCLEOPORIN85 is required for calcium spiking, fungal and bacterial symbioses, and seed production in *Lotus japonicus*. *Plant Cell* **19**, 610-624.
- Shibata S, Matsuoka Y, Yoneda Y.** 2002. Nucleocytoplasmic transport of proteins and poly(A) RNA in reconstituted Tpr-less nuclei in living mammalian cells. *Genes Cells* **7**, 421–434.
- Siniosoglou S, Lutzmann M, Santos-Rosa H, Leonard K, Mueller S, Aebi U, Hurt E.** 2000. Structure and assembly of the Nup84p complex. *J Cell Biol* **149**, 41-54.
- Strambio-De-Castillia C, Niepel M, Rout MP.** 2010. The nuclear pore complex: bridging nuclear transport and gene regulation. *Nat Rev Mol Cell Biol* **11**, 490-501.
- Strawn LA, Shen T, Shulga N, Goldfarb DS, Wente SR.** 2004. Minimal nuclear pore complexes define FG repeat domains essential for transport. *Nat Cell Biol* **6**, 197-206.
- Tamura K, Hara-Nishimura I.** 2011. Involvement of the nuclear pore complex in morphology of the plant nucleus. *Nucleus* **2**, 168-172.
- Tamura K, Fukao Y, Iwamoto M, Haraguchi T, Hara-Nishimura I.** 2010. Identification and characterization of nuclear pore complex components in *Arabidopsis thaliana*. *Plant Cell* **22**, 4084-4097.
- Wiermer M, Cheng YT, Imkampe J, Li M, Wang D, Lipka V, Li X.** 2012. Putative members of the *Arabidopsis* Nup107-160 nuclear pore sub-complex contribute to

- pathogen defense. *Plant J* **70**, 796-808.
- Wozniak RW, Blobel G, Rout MP.** 1994. POM152 is an integral protein of the pore membrane domain of the yeast nuclear envelope. *J Cell Biol* **125**, 31–42.
- Xu XM, Meier I.** 2008. The nuclear pore comes to force. *Trends Plant Sci* **13**, 20-27.
- Xu XM, Rose A, Muthuswamy S, Jeong SY, Venkatakrishnan S, Zhao Q, Meier I.** 2007a. NUCLEAR PORE ANCHOR, the Arabidopsis homolog of Tpr/Mlp1/Mlp2/megator, is involved in mRNA export and SUMO homeostasis and affects diverse aspects of plant development. *Plant Cell* **19**, 1537-1548.
- Xu XM, Meulia T, Meier I.** 2007b. Anchorage of plant RanGAP to the nuclear envelope involves novel nuclear-pore-associated proteins. *Curr Biol* **17**, 1157-1163.
- Yamazumi Y, Kamiya A, Nishida A, Nishihara A, Iemura S, Natsume T, Akiyama T.** 2009. The transmembrane nucleoporin NDC1 is required for targeting of ALADIN to nuclear pore complexes. *Biochem Biophys Res Commun* **389**, 100-104.
- Yoo BY, Bayley ST.** 1967. The structure of pores in isolated pea nuclei. *J Ultrastruct Res* **18**, 651-660.
- Zhang H, Saitoh H, Matunis MJ.** 2002. Enzymes of the SUMO modification pathway localize to filaments of the nuclear pore complex. *Mol Cell Biol* **22**, 6498-6508.
- Zhang Y, Li X.** 2005. A putative nucleoporin 96 Is required for both basal defense and constitutive resistance responses mediated by suppressor of npr1-1, constitutive 1. *Plant Cell* **17**, 1306-1316.
- Zhao Q, Meier I.** 2011. Identification and characterization of the Arabidopsis FG-repeat nucleoporin Nup62. *Plant Signal Behav* **6**, 330-334.
- Zhao Q, Brkljacic J, Meier I.** 2008. Two distinct interacting classes of nuclear envelope-associated coiled-coil proteins are required for the tissue-specific

nuclear envelope targeting of Arabidopsis RanGAP. *Plant Cell* **20**, 1639-1651.


Zhou X, Graumann K, Evans DE, Meier I. 2012. Novel plant SUN-KASH bridges are involved in RanGAP anchoring and nuclear shape determination. *J Cell Biol* **196**, 203-211.

Figure legends

Fig. 1. Summary of *Arabidopsis* nucleoporins. The structural motifs that appear next to each nucleoporin refer to the yeast protein folds predicted by Devos *et al.* (2006). Protein fold illustrations were derived from PDB ID: 1D7M, 1DXZ, 1XKS 2Q5X, 2RFO, and 3T97. AGI code: *Arabidopsis* genome initiative code.

Fig. 2. Molecular architecture of the NPC based on the model proposed by Alber *et al.* (2007a). The nucleoporins (Nups) are grouped according to their location. The NPC is divided into seven component groups. The symmetrical core is composed of the outer ring Nups, the linker Nups, the inner ring Nups, the transmembrane ring Nups, and the central FG Nups. The asymmetric parts of the pore are formed by the cytoplasmic FG Nups and filaments and the nuclear FG Nups and basket.

Fig. 3. NPC-associated proteins in plants. The plant NPC associates with many proteins, including the TREX-2 complex, ESD4, and the RanGAP-WIT-WIP complex. This allows the NPC to participate in diverse cellular functions.



	Name	AGI code
	Nup160	At1g33410
	Nup155	At1g14850
	Nup133	At2g05120
	Nup107	At3g14120
	Nup96	At1g80680
	Nup75	At4g32910
	Nup205	At5g51200
	Nup188	At4g38760
	Nup93	At2g41620/At3g57350
	Nup43	At4g30840
	SEH1	At1g64350
	SEC13	At2g30050
	Aladin	At3g56900
	RAE1	At1g80670
	Nup88	At5g05680
	NUA	At1g79280
	Nup136/Nup1	At3g10650
	Nup98	At1g10390/At1g59660
	Nup50	At1g52380/At3g15970
	CG1	At1g75340
	Nup62	At2g45000
	Nup58	At4g37130
	Nup54	At1g24310
	Nup214	AAD10642
	gp210	At5g40480
	NDC1	At1g73240
	Elys/HOS1	At2g39810
	GLE1	At1g13120
	Nup35	At3g16310

Figure 1. Tamura and Hara-Nishimura




Figure 2. Tamura and Hara-Nishimura




Figure 3. Tamura and Hara-Nishimura