

AZUMAYA ALGEBRAS AND SKEW POLYNOMIAL RINGS

Shuichi IKEHATA

Department of Mathematics, Okayama University

This note is an abstract of the author's paper [1] and includes some improvements of the results in it.

Throughout this note, every ring has identity 1, its subring contains 1, and every module over a ring is unital. A ring homomorphism means such one sending 1 to 1. In what follows, B will represent a ring, ρ an automorphism of B , D a ρ -derivation of B (i.e. an additive endomorphism of B such that $D(ab) = D(a)\rho(b) + aD(b)$ for all $a, b \in B$). Let $R = B[X; \rho, D]$ be the skew polynomial ring in which the multiplication is given by $aX = X\rho(a) + D(a)$ ($a \in B$). By $R_{(0)}$, we denote the set of all monic polynomials in R with $gR = Rg$.

A ring extension B/A is called to be separable if the B - B -homomorphism of $B \otimes_A B$ onto B defined by $a \otimes b \rightarrow ab$ splits, and B/A is called to be H-separable if $B \otimes_A B$ is B - B -isomorphic to a direct summand of a finite direct sum of copies of B . As

is well known, an H-separable extension is separable. A polynomial g in $R_{(0)}$ is called to be separable (resp. H-separable) if R/gR is a separable (resp. H-separable) extension of B . Moreover, a ring extension B/A is called to be G-Galois if there exists a finite group G of automorphisms of B such that $A = B^G$ (the fixed ring of G in B) and $\sum_i x_i \sigma(y_i) = \delta_{1,\sigma}$ ($\sigma \in G$) for some finite $x_i, y_i \in B$.

We shall use the following conventions:

$U(B)$ = the set of all invertible elements in B .

u_l (resp. u_r) = the left (resp. right) multiplication effected by $u \in B$, $B_l = \{u_l \mid u \in B\}$.

$B^0 = \{a \in B \mid \rho(a) = a\}$, $B^D = \{a \in B \mid D(a) = 0\}$.

1. H-separable polynomials. In our study, H-separable polynomials in skew polynomial rings play important rôles. Therefore, this section is devoted to giving some results concerning H-separable polynomials. Throughout, let $f = X^m + X^{m-1}a_{m-1} + \dots + Xa_1 + a_0$ be in $B[X; \rho, D]$ and $m \geq 2$. First, we state the following which is easily obtained from the result of Miyashita [2, Theorem 1.9].

Theorem 1.1. Let f be in $R_{(0)}$, and $I = fR$. If f is an H-separable polynomial in R , then there exist $y_i, z_i \in R$ with $\deg y_i < m$ and $\deg z_i < m$ such that $ay_i = y_i a$, $\rho^{m-1}(a)z_i = z_i a$ ($a \in B$) and

$\sum_i y_i x^{m-1} z_i \equiv 1 \pmod{I}$, $\sum_i y_i x^k z_i \equiv 0 \pmod{I}$ ($0 \leq k \leq m-2$), and conversely.

By virtue of Theorem 1.1, we have the following

Proposition 1.2. Let f be in $R_{(0)} = B[X; \rho]_{(0)}$. If f is H -separable in R , then $a_0 \in U(B)$, $\rho(a_0) = a_0$, $\rho^m = (a_0^{-1})_L (a_0)_R$, and $f = X^m + a_0$. Moreover, $\{g \in R \mid g \text{ is } H\text{-separable}\} = \{X^m + b_0 \mid b_0 \in U(Z \cap B^0) a_0\}$, where Z is the center of B .

Proposition 1.3. Let f be in $R_{(0)} = B[X; D]_{(0)}$. If f is H -separable in R , then B is of prime characteristic p , and f is a p -polynomial of the form $\sum_{j=0}^e x^{p^j} b_{j+1} + b_0$ ($p^e = m$). Moreover, $\{g \in R \mid g \text{ is } H\text{-separable}\} = \{\sum_{j=0}^e x^{p^j} b_{j+1} + \beta \mid \beta - b_0 \in Z \cap B^D\}$.

2. Azumaya algebras induced by $B[X; \rho]$. Throughout this section, B will mean a commutative ring, ρ an automorphism of B , G the cyclic group generated by ρ , $A = B^G = B$, and $R = B[X; \rho]$.

Theorem 2.1. Let $f = X^m + X^{m-1} a_{m-1} + \dots + X a_1 + a_0$ be in $R_{(0)}$, and $S = R/fR$. Then, f is H -separable in R if and only if S is an Azumaya A -algebra. When this is the case, there holds that B/A is G -Galois, the order of G is m , $f = X^m + a_0$, and $a_0 \in U(A)$.

Theorem 2.2. The following conditions are equivalent:

- (a) B/A is a G -Galois extension with G of order m .
- (b) $R_{(0)}$ contains an H -separable polynomial of degree m .
- (c) $R_{(0)}$ contains a polynomial f of degree m such that R/fR is an Azumaya A -algebra.
- (d) $\{g \in R \mid g \text{ is } H\text{-separable}\} = \{X^m + a \mid a \in U(A)\}$.

When this is the case, for every $a \in U(A)$, B is a maximal commutative A -subalgebra of $R/(X^m + a)R$, $(R/(X^m + a)R) \otimes_A B \cong B \otimes_A (R/(X^m + a)R) \cong M_m(B)$, and moreover, if $m \in U(A)$ then $A[X]/(X^m + a)A[X]$ is a separable splitting ring for $R/(X^m + a)R$.

Theorem 2.3. Assume that R contains an H -separable polynomial of degree $m \geq 2$. For $f \in R_{(0)}$, the following conditions are equivalent:

- (a) f is separable in R .
- (b) $f = g(X^m)$ or $Xg(X^m)$ for some $g(t)$ in $A[t]_{(0)}$ such that $g(t)$ is separable in $A[t]$ and the constant term of $g(t)$ is in $U(A)$.
- (c) R/fR is a separable A -algebra.

3. Azumaya algebras induced by $B[X;D]$. Throughout this section, B will mean a commutative ring, D a derivation of B , $A = B^D$ and $R = B[X;D]$.

Theorem 3.1. Let $f \in R_{(0)}$, $\deg f = m$, and $S = R/fR$. Then the following conditions are equivalent:

- (a) f is H -separable in R .
- (b) S is an Azumaya A -algebra.
- (c) There exist $y_i, z_i \in B$ such that $\sum_i D^{m-1}(y_i)z_i = 1$ and $\sum_i D^k(y_i)z_i = 0$ ($0 \leq k \leq m-2$).

Theorem 3.2. The followings are equivalent:

- (a) ${}_A B$ is a finitely generated projective module of rank m and $\text{Hom}({}_A B, {}_A B) = B[D]$ (the subring generated by B_ℓ and D).
- (b) R contains an H -separable polynomial f of degree m .
- (c) $R_{(0)}$ contains a polynomial f of degree m such that R/fR is an Azumaya A -algebra.
- (d) $R_{(0)}$ contains a polynomial f of degree m , and there exist $y_i, z_i \in B$ such that $\sum_i D^{m-1}(y_i)z_i = 1$ and $\sum_i D^k(y_i)z_i = 0$ ($0 \leq k \leq m-2$).

When this is the case, for any H -separable polynomial f , there holds the following:

- (1) $R = B[X;D]$ is an Azumaya $A[f]$ -algebra such that $B[f]$ is a maximal commutative $A[f]$ -sub-

algebra of R with $B[f] \otimes_{A[f]} R \cong R \otimes_{A[f]} B[f] \cong M_m(B[f])$.

(2) B is a maximal commutative A -subalgebra of R/fR with $B \otimes_A (R/fR) \cong (R/fR) \otimes_A B \cong M_m(B)$.

Theorem 3.3. Assume that R contains an H -separable polynomial f . Let $\Psi : A[t] \rightarrow R$ be defined by $\Psi(g_0(t)) = g_0(f)$.

(a) Ψ induces a one-to-one correspondence between $A[t]_{(0)}$ and $R_{(0)}$.

(b) For $g_0 \in A[t]_{(0)}$, g_0 is separable in $A[t]$ if and only if $R/\Psi(g_0)R$ is a separable A -algebra, and moreover, $\Psi(g_0)$ is H -separable in R if and only if $\deg g_0 = 1$.

References

- [1] S. Ikehata: Azumaya algebras and skew polynomial rings, Math. J. Okayama Univ., 23(1981), 19 - 32.
- [2] Y. Miyashita: On a skew polynomial ring, J. Math. Soc. Japan 31(1979), 317 - 330.