

ON THE NORM CONVERGENCE OF THE TROTTER-KATO PRODUCT FORMULA WITH ERROR BOUND

金沢大理 一瀬 孝 (Takashi Ichinose*)

岡山大理 田村 英男 (Hideo Tamura**)

*Department of Mathematics, Faculty of Science, Kanazawa Univ.

** Department of Mathematics, Faculty of Science, Okayama Univ.

Abstract. The norm convergence of the Trotter-Kato product formula with error bound is shown for the semigroup generated by that operator sum of two nonnegative selfadjoint operators A and B which is selfadjoint.

1. Introduction and Result

It is well-known ([23], [15]; [19]) that the Trotter-Kato product formula for the selfadjoint semigroup holds in strong operator topology. Namely, when A and B are nonnegative selfadjoint operators in a Hilbert space $\mathcal{H}$ with domains $D[A]$ and $D[B]$, then

$$\text{s-lim}_{n \rightarrow \infty} (e^{-tB/2n} e^{-tA/n} e^{-tB/2n})^n = \text{s-lim}_{n \rightarrow \infty} (e^{-tA/n} e^{-tB/n})^n = e^{-tC}, \quad (1.1)$$

if C is the form sum $A+B$ which is selfadjoint, or, in particular, if the operator sum $A+B$ is essentially selfadjoint on $D[A] \cap D[B]$ with C its closure. The convergence is uniform on each compact t -interval in the closed half line $[0, \infty)$.

The aim of this note is to briefly announce our recent results on its operator-norm convergence with error bound. In [12] we have shown

Theorem 1.1. *If A and B are nonnegative selfadjoint operators in $\mathcal{H}$ with domains $D[A]$ and $D[B]$ and if their operator sum $C := A+B$ is selfadjoint on $D[C] = D[A] \cap D[B]$, then the product formula in operator norm holds with error bound:*

$$\begin{aligned} \|(e^{-tB/2n} e^{-tA/n} e^{-tB/2n})^n - e^{-tC}\| &= O(n^{-1/2}), \\ \|(e^{-tA/n} e^{-tB/n})^n - e^{-tC}\| &= O(n^{-1/2}), \quad n \rightarrow \infty. \end{aligned} \quad (1.2)$$

The convergence is uniform on each compact t -interval in the open half line $(0, \infty)$, and further, if C is strictly positive, uniform on the closed half line $[T, \infty)$ for every fixed $T > 0$.

One of the typical examples of such a selfadjoint operator $C = A + B$ is the Schrödinger operator

$$H = -\frac{1}{2}\Delta + P|x|^{-1} + D|x|^2 + E|x|^{2000}$$

in $L^2(\mathbf{R}^3)$, where P , D and E are nonnegative constants.

Remark 1.1 The first result of such a norm convergence of the Trotter–Kato product formula (1.1) was proved by Rogava [20] in the abstract case under an additional condition that B is A -bounded, with error bound $O(n^{-1/2} \log n)$. The next was by Helffer [5] for the Schrödinger operators $H = H_0 + V \equiv -\frac{1}{2}\Delta + V(x)$ with C^∞ nonnegative potentials $V(x)$, roughly speaking, growing at most of order $O(|x|^2)$ for large $|x|$ with error bound $O(n^{-1})$. Each of these two results is independent of the other.

Then under some stronger or more general conditions, several further results are obtained. As for the abstract case, a better error bound $O(n^{-1} \log n)$ than Rogava's is obtained by Ichinose–Tamura [11] (cf. [9]) when B is A^α -bounded for some $0 < \alpha < 1$, even though the $B = B(t)$ may be t -dependent, and by Neidhardt–Zagrebnov [16], [17] (cf. [18]) when B is A -bounded with relative bound less than 1. As for the Schrödinger operators, a different proof to Helffer's result was obtained by Dia-Schatzman [2]. Further, more general results were proved for continuous nonnegative potentials $V(x)$, roughly speaking, growing of order $O(|x|^\rho)$ for large $|x|$ with $\rho > 0$, together with error bounds dependent on the power ρ (for instance, of order $O(n^{-2/\rho})$, if $\rho \geq 2$), by Ichinose–Takanobu [6] (cf. [7]), Doumeki–Ichinose–Tamura [3], Ichinose–Tamura [10], Decombes–Dia [1] and others, although the primary purpose of most of these papers was to prove rather a norm estimate between the Kac transfer operator and its corresponding Schrödinger semigroup. The Schrödinger operators treated in [6] and [3] may even involve bounded magnetic fields $\nabla \times A(x)$: $H = H_0(A) + V \equiv \frac{1}{2}(-i\nabla - A(x))^2 + V(x)$. In [7] and [8] the relativistic Schrödinger operator was also dealt with.

It should be noted (see [4], [21]) that in all these cases of the Schrödinger operators the sum $H = H_0 + V$ (resp. $H = H_0(A) + V$) is selfadjoint on the domain $D[H] = D[H_0] \cap D[V]$ (resp. $D[H] = D[H_0(A)] \cap D[V]$).

Thus the present theorem not only extends Rogava's result, but also can extend and contain all the results mentioned above, inclusive better error bounds in some cases.

Remark 1.2. Unless the sum $A + B$ is selfadjoint on $D[A] \cap D[B]$, the norm convergence of the Trotter–Kato product formula does not always hold, even though the sum is essentially selfadjoint there and B is A -form-bounded with relative bound less than 1. A counterexample is due to Hiroshi Tamura [22].

The theorem also holds with the exponential function e^{-s} replaced by real-valued, Borel measurable functions f and g on $[0, \infty)$ satisfying that

$$0 \leq f(s) \leq 1, \quad f(0) = 1, \quad f'(0) = -1, \quad (1.3)$$

that for every small $\varepsilon > 0$ there exists a positive constant $\delta = \delta(\varepsilon) < 1$ such that

$$f(s) \leq 1 - \delta(\varepsilon), \quad s \geq \varepsilon, \quad (1.4)$$

and that, for some fixed constant κ with $1 < \kappa \leq 2$,

$$[f]_\kappa := \sup_{s>0} s^{-\kappa} |f(s) - 1 + s| < \infty, \quad (1.5)$$

and the same for g . Of course, the functions $f(s) = e^{-s}$ and $f(s) = (1 + k^{-1}s)^{-k}$ with $k > 0$ are examples of functions having these properties.

Theorem 1.2. *If $3/2 \leq \kappa \leq 2$, it holds in operator norm that*

$$\begin{aligned} \| [g(tB/2n)f(tA/n)g(tB/2n)]^n - e^{-tC} \| &= O(n^{-1/2}), \\ \| [f(tA/n)g(tB/n)]^n - e^{-tC} \| &= O(n^{-1/2}), \quad n \rightarrow \infty. \end{aligned} \quad (1.6)$$

2. Outline of Proof

To proving the theorem, it is crucial to show the following operator-norm version of Chernoff's theorem with error bounds. The case without error bounds was noted by Neidhardt-Zagrebnov [18].

Lemma. *Let C be a nonnegative selfadjoint operator in a Hilbert space $\mathcal{H}$ and let $\{F(t)\}_{t \geq 0}$ be a family of selfadjoint operators with $0 \leq F(t) \leq 1$. Define $S_t = t^{-1}(1 - F(t))$. Then in the following two assertions, for $0 < \alpha \leq 1$, (a) implies (b).*

(a)

$$\|(1 + S_t)^{-1} - (1 + C)^{-1}\| = O(t^\alpha), \quad t \downarrow 0. \quad (2.1)$$

(b) *For any $\delta > 0$ with $0 < \delta \leq 1$,*

$$\|F(t/n)^n - e^{-tC}\| = \delta^{-2} t^{-1+\alpha} e^{\delta t} O(n^{-\alpha}), \quad n \rightarrow \infty, \quad (2.2)$$

for all $t > 0$.

Therefore, for $0 < \alpha < 1$ (resp. $\alpha = 1$), the convergence in (2.2) is uniform on each compact t -interval in the open half line $(0, \infty)$ (resp. in the closed half line $[0, \infty)$).

Moreover, if C is strictly positive, i.e. $C \geq \eta$ for some constant $\eta > 0$, the error bound on the right-hand side of (2.2) can also be replaced by $(1 + 2/\eta)^2 t^{-1+\alpha} O(n^{-\alpha})$, so that, for $0 < \alpha < 1$ (resp. $\alpha = 1$), the convergence in (2.2) is uniform on the closed half line $[T, \infty)$ for every fixed $T > 0$ (resp. on the whole closed half line $[0, \infty)$).

Sketch of Proof of Lemma.

Put

$$F(t/n)^n - e^{-tC} = (F(t/n)^n - e^{-tS_{t/n}}) + (e^{-tS_{t/n}} - e^{-tC}).$$

For the first term on the right we have by the spectral theorem

$$\|F(t/n)^n - e^{-tS_{t/n}}\| = \|F(t/n)^n - e^{-n(1-F(t/n))}\| \leq e^{-1} n^{-1},$$

$$0 \leq e^{-n(1-\lambda)} - \lambda^n \leq e^{-1}/n, \quad \text{for } 0 \leq \lambda \leq 1.$$

For the second term, we use

$$\begin{aligned} & (1 + S_\varepsilon)^{-1} [e^{-t(\delta+S_\varepsilon)} - e^{-t(\delta+C)}] (1 + C)^{-1} \\ &= \int_0^t e^{-(t-s)(\delta+S_\varepsilon)} [(1 + S_\varepsilon)^{-1} - (1 + C)^{-1}] e^{-s(\delta+C)} ds \\ &= \int_0^{t/2} + \int_{t/2}^t. \end{aligned}$$

where $0 < \delta \leq 1$ and $\varepsilon > 0$, to bound these two integrals on the right by $(\delta^2 t)^{-1} e^{\delta t} O(\varepsilon^\alpha)$. Taking $\varepsilon = t/n$, we have

$$\|e^{-tS_{t/n}} - e^{-tC}\| \leq (\delta^2 t)^{-1} e^{\delta t} O((t/n)^\alpha) = \delta^{-2} t^{-1+\alpha} e^{\delta t} O(n^{-\alpha}).$$

Sketch of Proof of Theorems 1.1 and 1.2.

First note that since $C = A + B$ is itself selfadjoint and so a closed operator, by the closed graph theorem there exists a constant a such that

$$\|(1 + A)u\| + \|(1 + B)u\| \leq a\|(1 + C)u\|, \quad u \in D[C] = D[A] \cap D[B].$$

The proof of the theorem is divided into two cases, (a) the symmetric product case

$$F(t) = e^{-tB/2} e^{-tA} e^{-tB/2}, \quad (2.3)$$

and (b) the non-symmetric product case

$$G(t) = e^{-tA} e^{-tB}. \quad (2.4)$$

(a) In the symmetric case we put

$$S_t = t^{-1}(1 - F(t)) = t^{-1}(1 - e^{-tB/2} e^{-tA} e^{-tB/2})$$

and use Lemma to show that

$$\|(1 + S_t)^{-1} - (1 + C)^{-1}\| = O(t^{1/2}), \quad t \downarrow 0.$$

Put

$$A_t = t^{-1}(1 - e^{-tA}), \quad B_t = t^{-1}(1 - e^{-tB}), \quad C_t = t^{-1}(1 - e^{-tC}).$$

We have

$$\begin{aligned} 1 + S_t &= 1 + A_t + B_{t/2} - \frac{t}{4} B_{t/2}^2 + \frac{t^2}{4} B_{t/2} A_t B_{t/2} - \frac{t}{2} (A_t B_{t/2} + B_{t/2} A_t) \\ &= K_t^{1/2} (1 + Q_t) K_t^{1/2}, \end{aligned}$$

$$K_t = 1 + A_t + B_{t/2} - \frac{t}{4}B_{t/2}^2 \geq 1,$$

$$Q_t = \frac{t^2}{4}K_t^{-1/2}B_{t/2}A_tB_{t/2}K_t^{-1/2} - \frac{t}{2}K_t^{-1/2}(A_tB_{t/2} + B_{t/2}A_t)K_t^{-1/2}.$$

Then we can show

$$\|(1 + Q_t)^{-1}\| \leq 2/(3 - \sqrt{5}), \quad (2.5)$$

$$\|(1 + S_t)^{-1}K_t^{1/2}\| = \|K_t^{-1/2}(1 + Q_t)^{-1}\| \leq 2/(3 - \sqrt{5}). \quad (2.6)$$

Then we have

$$\begin{aligned} & (1 + S_t)^{-1} - (1 + C)^{-1} \\ &= (1 + S_t)^{-1}[A + B - (A_t + B_{t/2} - \frac{t}{4}B_{t/2}(1 - tA_t)B_{t/2} \\ & \quad - \frac{t}{2}(A_tB_{t/2} + B_{t/2}A_t))](1 + C)^{-1} \\ &= (1 + S_t)^{-1}(A - A_t)(1 + C)^{-1} + (1 + S_t)^{-1}(B - B_{t/2})(1 + C)^{-1} \\ & \quad + (1 + S_t)^{-1}[\frac{t}{4}B_{t/2}(1 - tA_t)B_{t/2} + \frac{t}{2}(A_tB_{t/2} + B_{t/2}A_t)](1 + C)^{-1} \\ &\equiv R_1(t) + R_2(t) + R_3(t). \end{aligned} \quad (2.7)$$

We can show the bounds

$$\|R_i(t)\| \leq ct^{1/2}, \quad i = 1, 2, 3, \quad (2.8)$$

with some constant $c > 0$. For instance, we can get the bound for $R_1(t)$, via the expression

$$\begin{aligned} R_1(t) &= [(1 + S_t)^{-1}K_t^{1/2}][K_t^{-1/2}(1 + A_t)^{1/2}] \\ &\quad \times [(1 + A_t)^{-1/2} - (1 + A_t)^{1/2}(1 + A)^{-1}](1 + A)(1 + C)^{-1} \end{aligned}$$

by (2.6) and the spectral theorem

$$\|R_1(t)\| \leq \frac{2}{3-\sqrt{5}}a\|(1 + A_t)^{-1/2} - (1 + A_t)^{1/2}(1 + A)^{-1}\| \leq ct^{1/2}.$$

(b) The non-symmetric case will follow from the symmetric case. We use the commutator argument to observe that

$$\begin{aligned} \|G(t/n)^n - F(t/n)^n\| &= \|(e^{-tA/n}e^{-tB/n})^n - (e^{-tB/2n}e^{-tA/n}e^{-tB/2n})^n\| \\ &= O(1/n). \end{aligned}$$

3. The Final Result

In a recent preprint [14], we have shown that if $\kappa = 2$, then Theorem 1.2 holds with optimal error bound $O(n^{-1})$. Further, the convergence is uniform on each compact t -interval in the closed half line $[0, \infty)$, and further, if C is strictly positive, uniform on the whole closed half line $[0, \infty)$.

The idea of proof is simply to iterate the resolvent equation of the first identity in (2.5) with help of its adjoint form to get

$$\begin{aligned} & (1 + S_t)^{-1} - (1 + C)^{-1} \\ &= ((1 + C)^{-1} + [(1 + S_t)^{-1} - (1 + C)^{-1}](C - S_t)(1 + C)^{-1} \\ &= (1 + C)^{-1}(C - S_t)(1 + C)^{-1} + [(C - S_t)(1 + C)^{-1}]^*(1 + S_t)^{-1}(C - S_t)(1 + C)^{-1} \\ &\equiv R'_1(t) + R'_2(t). \end{aligned}$$

Then by the same arguments together with (2.6) we can show the bounds

$$\|R'_i(t)\| = O(t), \quad i = 1, 2.$$

Therefore it turns out that the product formula (1.2) in Theorem 1.1 holds, now with ultimate error bound $O(n^{-1})$, properly extending and containing all the known previous related results.

Finally, we comment about optimality of the error bound $O(n^{-1})$. We know that if both A and B are bounded operators, then we have, in the symmetric product case (2.3), $\|F(t/n)^n - e^{-tC}\| = O(n^{-2})$, while, in the non-symmetric product case (2.4), $\|G(t/n)^n - e^{-tC}\| = O(n^{-1})$. But also in the symmetric product case, we can give an example of two unbounded selfadjoint operators A and B whose operator sum $C = A + B$ is selfadjoint on $D[A] \cap D[B]$ such that $\|F(t/n)^n - e^{-tC}\| \geq L(t)n^{-1}$, with a positive continuous function $L(t)$ of $t > 0$ independent of n .

Part of the present results also was briefly announced in [13].

References

- [1] S. Deombes and B. O. Dia: *An operator-theoretic proof of an estimate on the transfer operator*, J. Functional Analysis, **165**, 240–257 (1999)
- [2] B. O. Dia and M. Schatzman: *An estimate of the Kac transfer operator*, J. Functional Analysis, **145**, 108–135 (1997)
- [3] A. Doumeki, T. Ichinose and Hideo Tamura: *Error bound on exponential product formulas for Schrödinger operators*, J. Math. Soc. Japan, **50**, 359–377 (1998)
- [4] D. Guibourg: *Inégalités maximales pour l'opérateur de Schrödinger*, C. R. Acad. Sci. Paris, **316**, Série I Math., 249–252 (1993)
- [5] B. Helffer: *Around the transfer operator and the Trotter–Kato formula*, Operator Theory: Advances and Appl., **78**, 161–174 (1995)
- [6] T. Ichinose and S. Takanobu: *Estimate of the difference between the Kac operator and the Schrödinger semigroup*, Commun. Math. Phys., **186**, 167–197 (1997)

- [7] T. Ichinose and S. Takanobu: *The norm estimate of the difference between the Kac operator and the Schrödinger semigroup: A unified approach to the nonrelativistic and relativistic cases*, Nagoya Math. J., **149**, 53–81 (1998)
- [8] T. Ichinose and S. Takanobu: *The norm estimate of the difference between the Kac operator and the Schrödinger semigroup II: The general case including the relativistic case*, Electronic Journal of Probability, **5**, Paper no. 5, 1–47 (2000)
- [9] T. Ichinose and Hideo Tamura: *Error estimates in operator norm for Trotter–Kato product formula*, Integr. Equat. Oper. Theory, **27**, 195–207 (1997)
- [10] T. Ichinose and Hideo Tamura, *Error bound in trace norm for Trotter–Kato product formula of Gibbs semigroups*, Asymptotic Analysis, **17**, 239–266 (1998)
- [11] T. Ichinose and Hideo Tamura: *Error estimates in operator norm of exponential product formulas for propagators of parabolic evolution equations*, Osaka J. Math., **35**, 751–770 (1998)
- [12] T. Ichinose and Hideo Tamura: *The norm convergence of the Trotter–Kato product formula with error bound*, to appear in Commun. Math. Phys.
- [13] T. Ichinose and Hideo Tamura: *On the norm convergence of the Trotter–Kato product formula with error bound*, to appear in Proceedings of Partial Differential Equations, Clausthal, July 24–28, 2000
- [14] T. Ichinose, Hideo Tamura, Hiroshi Tamura and V. A. Zagrebnov: *Note on the paper “The norm convergence of the Trotter–Kato product formula with error bound” by Ichinose and Tamura*, to appear
- [15] T. Kato: *Trotter’s product formula for an arbitrary pair of self-adjoint contraction semigroups*, In: Topics in Functional Analysis (G.C. Rota, ed.), Academic Press, New York, 1978, pp.185–195
- [16] H. Neidhardt and V. Zagrebnov: *On error estimates for the Trotter–Kato product formula*, Lett. Math. Phys., **44**, 169–186 (1998)
- [17] H. Neidhardt and V. Zagrebnov: *Fractional powers of selfadjoint operators and Trotter–Kato product formula*, Integr. Equat. Oper. Theory, **35**, 209–231 (1999)
- [18] H. Neidhardt and V. Zagrebnov: *Trotter–Kato product formula and operator-norm convergence*, Commun. Math. Phys., **205**, 129–159 (1999)
- [19] M. Reed and B. Simon: *Methods of Modern Mathematical Physics I: Functional Analysis*, Revised and enlarged ed., Academic Press, New York, London 1980
- [20] Dzh. L. Rogava: *Error bounds for Trotter-type formulas for self-adjoint operators*, Functional Analysis and Its Applications, **27**, 217–219 (1993)
- [21] Z. Shen: *L^p estimates for Schrödinger operators with certain potentials*, Ann. Inst. Fourier, Grenoble, **45**, 513–546 (1995); *Estimates in L^p for magnetic Schrödinger operators*, Indiana Univ. Math. J., **45**, 817–841 (1996)
- [22] Hiroshi Tamura: *A remark on operator-norm convergence of Trotter–Kato product formula*, Integr. Equat. Oper. Theory, **37**, 350–356 (2000)
- [23] H. F. Trotter: *On the product of semi-groups of operators*, Proc. Amer. Math. Soc., **10**, 545–551 (1959)