

On the integral closures of certain ideals generated by regular sequences

千葉大自然科学研究科 西田 康二 (Koji Nishida)

1 Introduction

The purpose of this report is to introduce a notion of equimultiplicity for filtrations in local rings. We will apply it's theory for computation of the integral closures of certain ideals generated by regular sequences.

Throughout this report A is a d -dimensional local ring with the maximal ideal $\mathfrak{m}$ and a family of ideals $\mathcal{F} = \{F_n\}_{n \in \mathbb{Z}}$ is a filtration in A , which means (i) $F_n \supseteq F_{n+1}$ for all $n \in \mathbb{Z}$, (ii) $F_0 = A$, $F_1 \neq A$ and (iii) $F_m F_n \subseteq F_{m+n}$ for all $m, n \in \mathbb{Z}$. We can define the following graded algebras associated to a filtration $\mathcal{F}$.

$$\begin{aligned} R(\mathcal{F}) &= \sum_{n \geq 0} F_n t^n \subseteq A[t], \\ R'(\mathcal{F}) &= \sum_{n \in \mathbb{Z}} F_n t^n \subseteq A[t, t^{-1}] \text{ and} \\ G(\mathcal{F}) &= R'(\mathcal{F})/t^{-1}R'(\mathcal{F}) = \oplus_{n \geq 0} F_n/F_{n+1}, \end{aligned}$$

where t is an indeterminate. These algebras are respectively called the Rees algebra of $\mathcal{F}$, the extended Rees algebra of $\mathcal{F}$ and the associated graded ring of $\mathcal{F}$. We always assume that $R(\mathcal{F})$ is Noetherian and $\dim R(\mathcal{F}) = d + 1$.

2 The analytic spread of a filtration

We set $\ell(\mathcal{F}) = \dim A/\mathfrak{m} \otimes_A R(\mathcal{F})$ and call it the analytic spread of $\mathcal{F}$. It is easy to see that $\ell(\mathcal{F}) = \dim A/\mathfrak{m} \otimes_A G(\mathcal{F})$. We say that a system of elements $a_1, \dots, a_r$ in A is a reduction of $\mathcal{F}$, if the following condition (*) is satisfied.

- (*) There exist $m_i > 0$ for all $1 \leq i \leq r$ such that $a_i \in F_{m_i}$ and $F_n = \sum_{i=1}^r a_i F_{n-m_i}$ for all $n \gg 0$.

This condition is equivalent to saying that we have a module-finite extension

$$A[a_1 t^{m_1}, \dots, a_r t^{m_r}] \subseteq R(\mathcal{F})$$

of rings. If $a_1, \dots, a_r$ is a reduction of $\mathcal{F}$, then obviously we have $\ell(\mathcal{F}) \leq r$. We say that a reduction $a_1, \dots, a_r$ of $\mathcal{F}$ is minimal, if $\ell(\mathcal{F}) = r$. We always have a minimal reduction for any filtration $\mathcal{F}$ (It is not necessary to assume that the residue field is infinite).

By the definition of filtration, we have $\sqrt{F_n} = \sqrt{F_1}$ for all $n \geq 1$, and so $\text{ht}_A F_n$ is constant for $n \geq 1$. We denote this number by $\text{ht}_A \mathcal{F}$. Then the following inequality always holds:

$$\text{ht}_A \mathcal{F} \leq \ell(\mathcal{F}) \leq \dim A.$$

We say that $\mathcal{F}$ is equimultiple, if $\text{ht}_A \mathcal{F} = \ell(\mathcal{F})$. If $\mathcal{F}$ is equimultiple and $a_1, \dots, a_r$ is a minimal reduction of $\mathcal{F}$, the number m_i in (*) must coincide to

$$\deg_{\mathcal{F}} a_i := \max\{n \mid a_i \in F_n\}$$

for all $1 \leq i \leq r$.

Example 2.1 Let $\mathfrak{p}$ be a prime ideal in A such that $\dim A/\mathfrak{p} = 1$. Let $F_n = \mathfrak{p}^{(n)}$ for all $n \in \mathbb{Z}$, where $\mathfrak{p}^{(n)}$ denotes the n -th symbolic power of $\mathfrak{p}$. If $R(\mathcal{F})$ is Noetherian, then $\mathcal{F}$ is equimultiple.

Proof. Because $R(\mathcal{F})$ is Noetherian, there exists a positive integer k such that $\mathfrak{p}^{(kn)} = [\mathfrak{p}^{(k)}]^n$ for all $n \in \mathbb{Z}$. This means the k -th Veronesean subring $R(\mathcal{F})^{(k)} = \sum_{n \geq 0} \mathfrak{p}^{(kn)} t^{kn}$ is isomorphic to $R(\mathfrak{p}^{(k)})$ and $\text{depth } A/[\mathfrak{p}^{(k)}]^n = 1$ for all $n \geq 1$. Then the extension

$$R(\mathfrak{p}^{(k)}) \subseteq R(\mathcal{F})$$

is module-finite and $\ell(\mathfrak{p}^{(k)}) = d - 1$ by Burch's inequality. Let $a_1, \dots, a_{d-1}$ be a minimal reduction of $\mathfrak{p}^{(k)}$. Then the extension

$$A[a_1 t^k, \dots, a_{d-1} t^k] \subseteq R(\mathcal{F})^{(k)}$$

is module-finite, and so

$$A[a_1, \dots, a_{d-1}] \subseteq R(\mathcal{F})$$

is also module-finite.

Example 2.2 Let J be an ideal in A generated by a subsystem of parameters $a_1, \dots, a_s$ for A . Let $\mathcal{F}$ be a filtration such that $J^n \subseteq F_n \subseteq \overline{J^n}$ for all $n \in \mathbb{Z}$. If $R(\mathcal{F})$ is Noetherian, then $\mathcal{F}$ is equimultiple and $a_1, \dots, a_s$ is a minimal reduction of $\mathcal{F}$.

Proof. Obviously, $\text{ht}_A \mathcal{F} = s$. As $J^n \subseteq F_n$ for all $n \in \mathbb{Z}$, $R(\mathcal{F})$ contains $A[a_1 t, \dots, a_s t]$. Moreover, as $F_n \subseteq \overline{J^n}$ for all $n \in \mathbb{Z}$, $R(\mathcal{F})$ is integral over $A[a_1 t, \dots, a_s t]$. Because $R(\mathcal{F})$ is Noetherian, we see that the extension

$$A[a_1 t, \dots, a_s t] \subseteq R(\mathcal{F})$$

is module-finite.

For a prime ideal $\mathfrak{p}$ in A containing F_1 , we set $\mathcal{F}_{\mathfrak{p}} = \{F_n A_{\mathfrak{p}}\}_{n \in \mathbb{Z}}$, which is a filtration in $A_{\mathfrak{p}}$. Obviously, $\ell(\mathcal{F}_{\mathfrak{p}}) \leq \ell(\mathcal{F})$ for any prime ideal $\mathfrak{p}$ in A containing F_1 .

3 Cohen-Macaulay property of the graded rings associated to equimultiple filtrations

Theorem 3.1 *Let A be a quasi-unmixed local ring. If $\mathcal{F}$ is equimultiple, then we have*

$$a(G(\mathcal{F})) = \max\{a(G(\mathcal{F}_{\mathfrak{p}})) \mid \mathfrak{p} \in \text{Assh}_A A/F_1\}$$

Theorem 3.2 *Let A be a Cohen-Macaulay ring. Let $\mathcal{F}$ be an equimultiple filtration. We set $s = \text{ht}_A \mathcal{F}$. Then the following conditions are equivalent:*

- (1) $G(\mathcal{F})$ is a Cohen-Macaulay ring.
- (2) $G(\mathcal{F}_{\mathfrak{p}})$ is Cohen-Macaulay for all $\mathfrak{p} \in \text{Assh}_A A/F_1$ and there exists a minimal reduction $a_1, \dots, a_s$ of $\mathcal{F}$ such that $A/(a_1, \dots, a_s) + F_n$ is Cohen-Macaulay for all $1 \leq n \leq a(G(\mathcal{F})) + \sum_{i=1}^s \deg_{\mathcal{F}} a_i$.

When this is the case, for any minimal reduction $b_1, \dots, b_s$ of $\mathcal{F}$, $A/(b_1, \dots, b_s) + F_n$ is Cohen-Macaulay for all $n \geq 1$ and

$$R(\mathcal{F}) = A[\{b_i t^{\deg_{\mathcal{F}} b_i}\}_{1 \leq i \leq s}, \{F_n t^n\}_{1 \leq n \leq a(G(\mathcal{F})) + \sum_{i=1}^s \deg_{\mathcal{F}} b_i}].$$

Corollary 3.3 *Let A be a Cohen-Macaulay ring. Let I be an equimultiple ideal. Then the following conditions are equivalent:*

- (1) $G(I)$ is a Cohen-Macaulay ring.
- (2) $G(I_{\mathfrak{p}})$ is Cohen-Macaulay for all $\mathfrak{p} \in \text{Assh}_A A/I$ and there exists a minimal reduction J of I such that $A/J + I^n$ is Cohen-Macaulay for all $1 \leq n \leq r_J(I)$.

4 Integral closures of certain ideals

Applying the results in section 3, we can prove the following assertions.

Example 4.1 Let $A = k[[X, Y, Z]]$ be the formal power series ring over a field k . Suppose that the ideal generated by the maximal minors of the matrix

$$\begin{pmatrix} X^\alpha & Y^{\beta'} & Z^{\gamma'} \\ Y^\beta & Z^\gamma & X^{\alpha'} \end{pmatrix}$$

is a prime ideal, where $\alpha, \beta, \gamma, \alpha', \beta'$ and γ' are all positive integers. We put $a = Z^{\gamma+\gamma'} - X^{\alpha'}Y^{\beta'}$, $b = X^{\alpha+\alpha'} - Y^\beta Z^{\gamma'}$ and $c = Y^{\beta+\beta'} - X^\alpha Z^\gamma$. Let $J = (a, b)A$. Then we have

$$\overline{J^n} = J^{n-1} \cdot (a, b, \{X^i Z^j c \mid i, j \geq 0 \text{ and } i/\alpha' + j/\gamma' \geq 1\})A$$

for all $n \geq 1$ and $\overline{R(J)}$ is a Cohen-Macaulay ring.

Example 4.2 Let $A = k[[X, Y, Z, W]]$ be the formal power series ring over a field k . Let α, β and γ be positive integers such that $0 < \alpha \leq \beta \leq \gamma$. We set

$$a = X^{\alpha+\ell} - Y^\beta W, b = Y^{\beta+m} - Z^\gamma W, c = Z^{\gamma+1} - X^\alpha W \text{ and } d = W^3 - X^\ell Y^m Z,$$

where $\ell = \gamma + \beta - 2\alpha + 1$ and $m = 2\gamma - \beta - \alpha + 1$. It is easy to see that a, b, c is a regular sequence in A . Let $J = (a, b, c)A$. Then we have

$$\begin{aligned} \overline{J} &= J + (\{X^i Y^j Z^k d \mid i/\alpha + j/\beta + k/\gamma \geq 1\})A, \\ \overline{J^2} &= \overline{J}^2 + (X^i Y^j Z^k d^2 \mid i/2\alpha + j/2\beta + k/2\gamma \geq 1)A \text{ and} \\ \overline{J^n} &= \overline{J}^{n-2} \cdot \overline{J^2} \text{ for } n \geq 3. \end{aligned}$$

Moreover $\overline{R(J)}$ is a Cohen-Macaulay ring.

参考文献

- [1] L. Burch, *Codimension and analytic spread*, Proc. Camb. Philos. Soc. **72** (1972), 369 - 373.
- [2] S. Goto and K. Nishida, *The Cohen-Macaulay and Gorenstein Rees algebras associated to filtrations*, Mem. Amer. Math. Soc. **526** (1994).
- [3] M. Herrmann, S. Ikeda and U. Orbanz, *Equimultiplicity and blowing up*, Springer-Verlag, Berlin Heidelberg New York London Paris Tokyo, 1988.
- [4] J. Herzog, *Generators and relations of abelian semigroups and semigroup rings*, Manuscripta Math. **3** (1970), 175 - 193.

- [5] C. Huneke, *Complete ideals in two-dimensional regular local rings*, Math. Sci. Res. Inst. Publ. **15** (1989), 325 – 337.
- [6] J. Lipman and B. Teissier, *Pseudo-rational local rings and a theorem of Briançon-Skoda about integral closure of ideals*, Michigan Math. J. **28** (1981), 97 – 116. K. Nishida, *Equimultiple filtrations in Cohen-Macaulay rings*, Preprint.
- [7] O. Zariski and P. Samuel, *Commutative algebra, II*, van Nostrand, New York, 1960.