

CLINICAL AND LABORATORY OBSERVATIONS

Use of cabozantinib to treat *MET*-amplified pediatric colorectal cancer

Sadahiro Fukui¹, Katsutsugu Umeda^{1*}, Hirohito Kubota¹, Atsushi Iwai¹, Ryo Akazawa¹, Kiyotaka Isobe¹, Kuniaki Tanaka¹, Kagehiro Kouzuki¹, Naoko Kawabata¹, Satoshi Saida¹, Itaru Kato¹, Hidefumi Hiramatsu¹, Yoshiro Itatani², Taro Funakoshi³, Souichi Adachi⁴, and Junko Takita¹

Department of ¹Pediatrics, ²Surgery, and ³Clinical Oncology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

⁴Department of Human Health Sciences, Graduate School of Medicine, Kyoto University, Kyoto, Japan

*Corresponding author:

Katsutsugu Umeda, Department of Pediatrics, Graduate School of Medicine, Kyoto University, 54 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto, 606-8507, Japan
Phone: +81-75-751-3290; Fax: +81-75-752-2361

Email address: umeume@kuhp.kyoto-u.ac.jp

Word count: Abstract, 96 words; main text, 1,277 words

Number of tables and figures: 3 figures, 1 table

Running title: Cabozantinib for *MET*-amplified pediatric CRC

Keywords: colorectal cancer, pediatric, cancer gene panel testing, *MET* inhibitor, cabozantinib

ABBREVIATIONS:

CA, cancer antigen; CEA, carcinoembryonic antigen; CRC, colorectal cancer; CT, computed tomography; EGFR, epidermal growth factor receptor; FOLFIRI, fluorouracil, leucovorin, and irinotecan; FOLFOX6, fluorouracil, leucovorin, and oxaliplatin; modified; TCGA, The Cancer Genome Atlas.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest associated with this manuscript.

ABSTRACT

Pediatric colorectal cancer (CRC) is extremely rare, with little information about genetic profiles compared with adult CRC. Here, a 13-year-old male with advanced CRC underwent cancer gene panel testing, which detected four genetic abnormalities (*MET* amplification in addition to *TP53*, *SMAD4*, and *CTNNA1* mutations) that might be associated with a poor prognosis. Based on high-level *MET* amplification, he received a multi-kinase inhibitor, cabozantinib, after failure of first- and second-line chemotherapy, resulting in transient disease stabilization. Tailored targeted therapy based on molecular profiling can be an effective treatment strategy for rare cancers such as pediatric CRC.

INTRODUCTION

Colorectal cancer (CRC) is very rare in the pediatric population, accounting for only 1% of all pediatric malignancies.¹ The predominance of high-grade tumors and advanced stage suggest that the biology of pediatric CRC is different from that of adult CRC.² In contrast to CRC associated with inherited cancer syndromes (e.g., familial adenomatous polyposis and Lynch syndrome) the genetics of pediatric non-inherited CRC remain to be elucidated. Whole-exome sequencing demonstrates that adolescent and young adult CRC patients have a higher frequency of damaging mutations than adult CRCs, while very few data are available for pediatric CRCs due to lack of patients.

In general, pediatric CRCs are treated in accordance with adult CRC treatment algorithms;¹ however, treatment of refractory cases, especially the efficacy of molecular target therapy, remains unknown.

Here, we report a pediatric case of advanced CRC who, based on the results of cancer gene panel testing, received the multi-kinase inhibitor cabozantinib after failure of first- and second-line chemotherapy.

CASE REPORT

A 13-year-old male presented with recurrent abdominal pain and bloody stool. He did not have an identifiable family history of inherited cancer syndromes. Abdominal computed

tomography (CT) revealed moderate ascites and global wall thickening in the splenic flexure (Fig. 1A), and colonoscopy revealed a circumferential mass in the descending colon (Fig. 1B). Serum cancer antigen (CA)19-9 and CA125 were elevated to 161.8 U/mL (normal upper limit, 37.0 U/mL) and 335.3 U/mL (normal upper limit, 35 U/mL), respectively, but carcinoembryonic antigen (CEA) was within the normal range (3.6 ng/mL; normal upper limit, 5.0 ng/mL). There were no physical findings suggestive of inherited cancer syndromes. He underwent left hemicolectomy, during which broad peritoneal dissemination was observed (Fig. 1C). He was diagnosed histogenetically as *RAS/BRAF* wild type, microsatellite stable signet ring cell carcinoma (Fig. 1D). First-line (fluorouracil, leucovorin, and irinotecan; FOLFIRI) and second-line (fluorouracil, leucovorin, and oxaliplatin; modified FOLFOX6) systemic chemotherapy in combination with the anti-epidermal growth factor receptor (EGFR) antibody panitumumab maintained stable disease for 10 months. However, the tumor became refractory to chemotherapy, resulting in progressive peritoneal dissemination and intestinal obstruction, with a gradual increase in CEA (10.8 ng/mL), CA19-9 (111.2 U/mL), and CA125 (194.1 U/mL) levels (Fig. 2).

Due to the advanced stage (pT4aN0M1c, stage IV), the patient underwent cancer gene panel testing (Foundation One®) during first-line chemotherapy, which revealed that the tumor harbored pathogenic mutations in *CTNNA1*, *SMAD4*, and *TP53*, as well as a *MET* amplification (106 copies/cell). No secondary germline findings linked to CRC were found

(Table). Due to the high-level *MET* amplification, the patient received cabozantinib (60 mg/day), after approval for the use of unapproved drugs was received from the Patient Safety Unit at Kyoto University Hospital. Three weeks after cabozantinib treatment, his symptoms (abdominal pain, nausea, and vomiting) improved transiently, which allowed him to spend time at home and maintain a certain quality of life with the aid of analgesics and sedatives. Cabozantinib treatment had no severe adverse effects. However, 1 month after initiation of cabozantinib, his disease exacerbated and he succumbed 13 months after the initial diagnosis.

DISCUSSION

Due to aggressive histology and tumor behavior, the present case underwent cancer gene panel testing, which detected pathogenic mutations in *TP53*, *SMAD4*, and *CTNNA1*, as well as *MET* amplification.

TP53 mutation is the most common genetic alteration found in human cancers. In CRC, *TP53* mutations are associated with advanced stage, and mutations with loss of transcriptional activity are associated with a poor prognosis.³ The *TP53* R196P mutation was found recurrently in CRC patients from the COSMIC database (<http://cancer.sanger.ac.uk/cosmic>).⁴ This mutation has 1% of the transcriptional activity of the wild type, and has a dominant-negative effect.⁵ Therefore, the *TP53* R196P mutation is considered pathogenic and associated with a poor prognosis. *SMAD4* mutations are common

in a variety of cancers. Loss of *SMAD4* expression and loss of heterozygosity at the *SMAD4* locus are associated with a poorer prognosis for CRC.^{6,7} In particular, the proportion of *SMAD4* mutations is higher in signet ring cell carcinoma than in other histological types. *SMAD4* R361H, recurrently found in adult CRC patients,⁴ drives tumor progression by down-regulating TGF- β signaling.⁸ *CTNNA1* is a protein related to cell adhesion, and epithelial mesenchymal transition caused by loss of *CTNNA1* function is closely related to cancer invasion and metastasis. A report showing that downregulation of *CTNNA1* expression is associated with a poor prognosis for CRC⁹ suggests that *CTNNA1* S349fs, which is likely a loss-of-function mutation, may contribute to tumor aggressiveness. *MET* amplification, found in about 1.7% of all CRCs, is a well-known poor prognostic factor.¹⁰ While less common in primary CRC, up to 22.6% of adult CRCs that are refractory to anti-EGFR drugs harbor *MET* amplification.¹¹ To explore the characteristics of the mutational profile of this case, we compared the mutation patterns of the four genes (*TP53*, *SMAD4*, *CTNNA1*, and *MET*) with mutation and copy number data from 212 patients/samples in The Cancer Genome Atlas (TCGA) cohort, which were obtained from cBioPortal (<https://www.cbioportal.org/>).^{12–14} We found no adult CRC cases harboring all four mutations concurrently (Fig 3A), which may reflect the aggressive clinical behavior of the case reported herein.

The present case, who experienced disease progression after multiple courses of panitumumab-containing chemotherapy, was thought to be resistant to other EGFR

1 inhibitors. Furthermore, the objective response rate of other salvage therapies, such as multi-
2 kinase inhibitor regorafenib or trifluridine and tipiracil, is extremely low for such refractory
3 cases.¹⁵ Cabozantinib, an oral multi-kinase inhibitor with activity against MET, RET, AXL,
4 VEGFR2, FLT3, and c-KIT, was approved recently for unresectable or metastatic renal cell
5 carcinoma, and for refractory hepatocellular carcinoma.¹⁶ To date, several MET inhibitors
6 with different kinase selectivity have been developed and used as monotherapy, or in
7 combination with other drugs, in a variety of clinical trials for advanced solid tumors.
8 Regarding CRC, MET inhibitors have excellent anti-tumor effects in tissues refractory to
9 EGFR inhibitors, and subgroup analysis revealed that they are effective in patients with a
10 high *MET* copy number.^{17,18} In a phase II study, approximately 10% of patients who were
11 resistant to EGFR inhibitors and had a high *MET* copy number showed an objective response
12 to combined treatment with the MET inhibitor tivantinib and the anti-human EGFR antibody
13 cetuximab.¹⁹ Taken together, we considered *MET* amplification as a possible actionable
14 molecular target in this case, and the patient received cabozantinib as third-line therapy,
15 which had a transient therapeutic effect. The *MET* copy number in the present case was very
16 high when compared with that of 157 tumors (including five CRCs) with *MET* amplification
17 (a genomic copy number of five or higher) in the International Cancer Genome Consortium
18 cohort from the COSMIC database (Fig. 3B).⁴ Therefore, the limited therapeutic effect of
19 cabozantinib in the present case was probably due to *MET* hyper-amplification-associated

MET inhibitor resistance, as previously described.²⁰ Other possible explanations include involvement of other genetic mutations (*TP53*, *SMAD4*, and *CTNNA1*) in resistance to treatment, and poor absorption of cabozantinib due to decreased gastrointestinal function resulting from ileus or peritoneal dissemination. If cabozantinib had been started earlier, or used in combination with other chemotherapies and molecular targeting drugs, it may have had a greater effect in terms of inhibiting tumor growth.

In conclusion, we experienced a rare case of pediatric CRC that had four genetic abnormalities associated with a poor prognosis. To clarify the pathogenesis and to improve the prognosis of pediatric CRC, it will be necessary to perform comprehensive genomic analysis immediately after histological confirmation of the disease.

FUNDING

This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

ETHICS STATEMENT

The patient's legal guardians provided written informed consent to publish photographs under the guidelines of the Kyoto University Graduate School and Faculty of Medicine Ethics Committee.

REFERENCES

1. Poles GC, Clark DE, Mayo SW, et al: Colorectal carcinoma in pediatric patients: A comparison with adult tumors, treatment and outcomes from the National Cancer Database. *J Pediatr Surg* 51:1061-6, 2016
2. Hill DA, Furman WL, Billups CA, et al: Colorectal carcinoma in childhood and adolescence: a clinicopathologic review. *J Clin Oncol* 25:5808-14, 2007
3. Nakayama M, Hong CP, Oshima H, et al: Loss of wild-type p53 promotes mutant p53-driven metastasis through acquisition of survival and tumor-initiating properties. *Nat Commun* 11:2333, 2020
4. Tate JG, Bamford S, Jubb HC, et al: COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res* 47:D941-d947, 2019
5. Monti P, Perfumo C, Bisio A, et al: Dominant-negative features of mutant TP53 in germline carriers have limited impact on cancer outcomes. *Mol Cancer Res* 9:271-9, 2011
6. Isaksson-Mettävainio M, Palmqvist R, Forssell J, et al: SMAD4/DPC4 expression and prognosis in human colorectal cancer. *Anticancer Res* 26:507-10, 2006
7. Itatani Y, Kawada K, Fujishita T, et al: Loss of SMAD4 from colorectal cancer cells promotes CCL15 expression to recruit CCR1+ myeloid cells and facilitate liver metastasis. *Gastroenterology* 145:1064-1075.e11, 2013
8. Lim SK, Hoffmann FM: Smad4 cooperates with lymphoid enhancer-binding factor 1/T cell-specific factor to increase c-myc expression in the absence of TGF-beta signaling. *Proc Natl Acad Sci U S A* 103:18580-5, 2006
9. Raftopoulos I, Davaris P, Karatzas G, et al: Level of alpha-catenin expression in colorectal cancer correlates with invasiveness, metastatic potential, and survival. *J Surg Oncol* 68:92-9, 1998
10. Zhang M, Li G, Sun X, et al: MET amplification, expression, and exon 14 mutations in colorectal adenocarcinoma. *Hum Pathol* 77:108-115, 2018
11. Raghav K, Morris V, Tang C, et al: MET amplification in metastatic colorectal cancer: an acquired response to EGFR inhibition, not a de novo phenomenon. *Oncotarget* 7:54627-54631, 2016
12. Cerami E, Gao J, Dogrusoz U, et al: The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov* 2:401-4, 2012
13. Gao J, Aksoy BA, Dogrusoz U, et al: Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 6:p11, 2013
14. The Cancer Genome Atlas Network. Comprehensive molecular characterization of human colon and rectal cancer. *Nature* 487:330-7, 2012
15. Xie YH, Chen YX, Fang JY: Comprehensive review of targeted therapy for colorectal cancer. *Signal Transduct Target Ther* 5:22, 2020.

- 1
2 16. Tomita Y, Tatsugami K, Nakaigawa N, et al: Cabozantinib in advanced renal cell
3 carcinoma: A phase II, open-label, single-arm study of Japanese patients. *Int J Urol* 27:952-
4 959, 2020
- 5 17. Bardelli A, Corso S, Bertotti A, et al: Amplification of the MET receptor drives
6 resistance to anti-EGFR therapies in colorectal cancer. *Cancer Discov* 3:658-73, 2013
- 7 18. Eng C, Bessudo A, Hart LL, et al: A randomized, placebo-controlled, phase 1/2
8 study of tivantinib (ARQ 197) in combination with irinotecan and cetuximab in patients with
9 metastatic colorectal cancer with wild-type KRAS who have received first-line systemic
10 therapy. *Int J Cancer* 139:177-86, 2016
- 11 19. Rimassa L, Bozzarelli S, Pietrantonio F, et al: Phase II Study of Tivantinib and
12 Cetuximab in Patients With KRAS Wild-type Metastatic Colorectal Cancer With Acquired
13 Resistance to EGFR Inhibitors and Emergence of MET Overexpression: Lesson Learned for
14 Future Trials With EGFR/MET Dual Inhibition. *Clin Colorectal Cancer* 18:125-132 e2, 2019
- 15 20. Oddo D, Siravegna G, Gloghini A, et al: Emergence of MET hyper-amplification at
16 progression to MET and BRAF inhibition in colorectal cancer. *Br J Cancer* 117:347-352,
17 2017
18

FIGURE LEGENDS

FIGURE 1. (A) Abdominal computed tomography (CT) at onset. The arrow indicates intestinal wall thickening at the splenic curvature. (B) Colonoscopy findings at onset. (C, D) Macroscopic appearance (C) and hematoxylin and eosin staining (D) of the resected specimen (magnification, $\times 400$).

FIGURE 2. Clinical course of the patient. He received fluorouracil, leucovorin, and irinotecan; (FOLFIRI) (a), FOLFIRI plus panitumumab (b), and fluorouracil, leucovorin, and oxaliplatin (modified FOLFOX6) plus panitumumab (c) for 10 months.

FIGURE 3. (A) OncoPrint showing the frequency and co-occurrence of alterations in four genes (*TP53*, *SMAD4*, *CTNNA1*, and *MET*) in our case and in 212 colorectal cancer patients from The Cancer Genome Atlas (TCGA). TCGA data were downloaded from cBioPortal (<https://www.cbioportal.org/>). The variant types are indicated in the color legend at the bottom of the OncoPrint. (B) Genomic copy number of *MET* in this case compared with that in the COSMIC database (<https://cancer.sanger.ac.uk/cosmic>). The red dot represents this case relative to $n=5$ other colorectal cancers (green) and $n=152$ other cancers with *MET* amplification (five or more copies) in the International Cancer Genome Consortium cohort (black).

Figure 1




Figure 2




Figure 3




Table. Sequencing results from FoundationOne CDx

Biomarker findings			
Microsatellite status	stable		
Tumor mutational burden	1 Muts/Mb		
Genomic findings			
Short variants			
Gene	Alteration	Variant allele frequency	COSMIC ID (Frequency of occurrence in colorectal cancer) [#]
SMAD4	R361H	0.3563	COSV61684056 (106)
TP53	R196P	0.3344	COSV52678297 (4)
CTNNA1	S349fs*10	0.318	None
KRAS	Wildtype	N.A	N.A
NRAS	Wildtype	N.A	N.A
Copy number variation			
Gene	Alteration	Copy number	
MET	Amplification	106	

Abbreviation: N.A, not applicable

[#] Based on COSMIC database (v94)