

A generalization of a curvature flow of graphs on $\mathbf{R}$

北海道大学・理学研究科 三上 敏夫 (Toshio Mikami)

Department of Mathematics

Hokkaido University

1. Introduction. Gauss curvature flow is known as a mathematical model of the wearing process of a convex stone rolling on a beach and has been studied by many authors (see [3, 5, 6] and the reference therein). In [5] we proposed and studied the discrete stochastic approximations of nonconvex functions which evolve by a convexified Gauss curvature and the PDE which appears as the continuum limit of discrete stochastic processes (see [6] for the similar results on the convexified Gauss curvature flow of closed hypersurfaces). In this paper we study a class of PDE which gives a generalization of a curvature flow of graphs on $\mathbf{R}$.

We briefly describe [5] to discuss the results in this paper more precisely. Alexandrov-Bakelman's generalized curvature played a crucial role in [5].

Definition 1 (see e.g. [1, section 9.6]). Let $R \in L^1(\mathbf{R}^n : [0, \infty), dx)$ and $u \in C(\mathbf{R}^n)$. For $A \in B(\mathbf{R}^n)(:=\text{Borel } \sigma\text{-field of } \mathbf{R}^n)$, put

$$w(R, u, A) := \int_{\cup_{x \in A} \partial u(x)} R(y) dy, \quad (1)$$

where $\partial u(x) := \{p \in \mathbf{R}^n | u(y) - u(x) \geq \langle p, y - x \rangle \text{ for all } y \in \mathbf{R}^n\}$ and $\langle \cdot, \cdot \rangle$ denotes the inner product in $\mathbf{R}^n$.

(It is known that $w(R, u, \cdot) : B(\mathbf{R}^n) \mapsto [0, \infty)$ is completely additive.)

For $R \in L^1(\mathbf{R}^n : [0, \infty), dx)$, we showed the existence and the uniqueness of a solution $u \in C([0, \infty) \times \mathbf{R}^n)$ to the following equation (see [5, Theorem 1]): for any $\varphi \in C_o(\mathbf{R}^n)$ and any $t \geq 0$,

$$\int_{\mathbf{R}^n} \varphi(x)(u(t, x) - u(0, x))dx = \int_0^t ds \int_{\mathbf{R}^n} \varphi(x)w(R, u(s, \cdot), dx). \quad (2)$$

In [5, Theorem 2], we proved that a continuous solution u to (2) sweeps in time $t > 0$ a region with volume given by $t \cdot w(R, u(0, \cdot), \mathbf{R}^n)$, and that, for a continuous solution u to (2) with a convex $u(0, \cdot)$, $x \mapsto u(t, \cdot)$ is convex for all $t > 0$.

We also showed that a continuous solution u to (2) is a viscosity solution of the following PDE (see [5, Theorem 3]):

$$\begin{aligned} \partial_t u(t, x) &= \chi(u, Du(t, x), t, x)R(Du(t, x)) \\ &\quad \times \max(\text{Det}(D^2u(t, x)), 0) \quad ((0, \infty) \times \mathbf{R}^n), \end{aligned} \quad (3)$$

where $Du(t, x) := (\partial u(t, x)/\partial x_i)_{i=1}^n$, $D^2u(t, x) := (\partial^2 u(t, x)/\partial x_i \partial x_j)_{i,j=1}^n$ and

$$\chi(u, p, t, x) := \begin{cases} 1 & \text{if } p \in \partial u(t, x), \\ 0 & \text{otherwise.} \end{cases}$$

Here $\partial u(t, x)$ denotes the subdifferential of the function $x \mapsto u(t, x)$. Conversely, we discussed under what conditions a viscosity solution to (3) is a solution to (2).

We briefly discuss what we study in this paper. In (1) we only considered the measure $R(y)dy$ which is absolutely continuous with respect to the Lebesgue measure dy . Otherwise $\omega(R, u, dx)$ is not generally completely additive.

Suppose that $n = 1$. In (1), replace $R(y)dy$ by a continuous Borel probability measure $P(dy)$. Then, using a similar notation, $\omega(P, u, dx)$ turn out to be a measure and (2) has a unique continuous solution u (see Theorem 1 in section 2).

Since $P(dy)$ is not generally absolutely continuous with respect to dy , we can not consider the PDE for u . For $(t, x) \in [0, \infty) \times \mathbf{R}$, put

$$U(t, x) := \int_{-\infty}^x (u(t, y) - u(0, y))dy + \int_0^x u(0, y)dy + tF(a), \quad (4)$$

where

$$F(x) := P((-\infty, x]), \quad a := \inf\{\cup_{x \in \mathbf{R}} \partial u(0, x)\}.$$

Then from (2),

$$U(t, x) - U(0, x) = \int_0^t F(D_+(\hat{D}U(s, x)))ds. \quad (5)$$

Here $\hat{u}$ denotes a convex envelope of u and for φ , $D_+\varphi(t, x)$ denotes the right derivative of $x \mapsto \varphi(t, x)$.

When $DU(0, x)$ is convex, we show that $U(t, x)$ is a unique continuous viscosity solution in $(0, \infty) \times \mathbf{R}$ of the following (see Theorems 2 and 3 in section 2):

$$\partial_t U(t, x) = F(D^2 U(t, x)). \quad (6)$$

Definition 2 (Viscosity solution) (1) Let $\Omega = (0, \infty) \times \mathbf{R}$.

(i). A function $U \in USC(\Omega)$ is called a viscosity subsolution of (6) in Ω if whenever $\varphi \in C^{1,2}(\Omega)$, $(s, y) \in \Omega$, and $U - \varphi$ attains a local maximum at (s, y) , then $\partial_t \varphi(s, y) \leq F(D^2 \varphi(s, y))$.

(ii). A function $U \in LSC(\Omega)$ is called a viscosity supersolution of (6) in Ω if whenever $\varphi \in C^{1,2}(\Omega)$, $(s, y) \in \Omega$, and $U - \varphi$ attains a local minimum at (s, y) , then $\partial_t \varphi(s, y) \geq F(D^2 \varphi(s, y))$.

(iii). A function $U \in C(\Omega)$ is called a viscosity solution of (6) in Ω if it is both a viscosity subsolution and a viscosity supersolution of (6) in Ω .

(2) Let $t_2 > t_1 > 0$, O be an open subset of $\mathbf{R}$ and $Q := (t_1, t_2] \times O$. A function $U \in C(\overline{Q})$ is called a viscosity solution of (6) in Q if (1,i)-(1,ii) with Ω replaced by Q hold (see [4, p. 66]). Here $\overline{Q}$ denotes the closure of Q .

2. Main results. We state assumptions under which we can generalize [5] when $n = 1$.

(A.1). P is a continuous Borel probability measure on $\mathbf{R}$.

(A.2). $h \in C(\mathbf{R})$ and the set $\partial h(\mathbf{R})$ has a positive Lebesgue measure.

(A.3). For any $p \notin \partial h(\mathbf{R})$ and $C \in \mathbf{R}$,

$$\int_{\mathbf{R}} \max(px + C - h(x), 0) dx = \infty.$$

Theorem 1 Suppose that (A.1)-(A.3) hold. Then there exists a unique continuous solution u to (2) with $u(0, \cdot) = h$.

The following assumption implies (A.2)-(A.3).

(A.2)'. h is convex and is not a constant.

Theorem 2 Suppose that (A.1) and (A.2)' hold. Then the function U defined, by (4), from u in Theorem 1 is a continuous viscosity solution of (6) in $(0, \infty) \times \mathbf{R}$.

As a regularity result, we have

Proposition 1 Suppose that (A.1) holds. Then for a continuous viscosity solution v of (6) in $(0, \infty) \times \mathbf{R}$,

$$0 \leq v(t, x) - v(s, x) \leq t - s \quad (0 \leq s < t, x \in \mathbf{R}). \quad (7)$$

In particular, $t \mapsto v(t, x)$ is absolutely continuous for all $x \in \mathbf{R}$.

We state an additional assumption and an asymptotic behavior of a viscosity solution of (6).

(A.4). (i) $v_0 : \mathbf{R} \mapsto \mathbf{R}$ is twice continuously differentiable.

(ii) $\lim_{x \rightarrow -\infty} D^2 v_0(x)$ and $\lim_{x \rightarrow \infty} D^2 v_0(x)$ exist and

$$a := \inf_{x \in \mathbf{R}} D^2 v_0(x) = \lim_{x \rightarrow -\infty} D^2 v_0(x),$$

$$b := \sup_{x \in \mathbf{R}} D^2 v_0(x) = \lim_{x \rightarrow \infty} D^2 v_0(x).$$

Proposition 2 *Suppose that (A.1) and (A.4, i) hold. Then for a continuous viscosity solution v of (6) with $v(0, \cdot) = v_0(\cdot)$ in $(0, \infty) \times \mathbf{R}$, the following holds: for any $t \geq 0$ and $x \in \mathbf{R}$,*

$$F(a)t \leq v(t, x) - v(0, x) \leq F(b)t. \quad (8)$$

Suppose in addition that (A.4, ii) holds. Then for any $T \geq 0$,

$$\lim_{x \rightarrow -\infty} \left(\sup_{0 \leq t \leq T} |v(t, x) - v(0, x) - F(a)t| \right) = 0, \quad (9)$$

$$\lim_{x \rightarrow \infty} \left(\sup_{0 \leq t \leq T} |v(t, x) - v(0, x) - F(b)t| \right) = 0. \quad (10)$$

Since F is nondecreasing, (6) is a degenerate elliptic PDE and we can use the maximum principle for this equation in a bounded domain (see [4, p. 244, Theorem 8.1] and also [2]). From Prop. 2, we immediately obtain

Theorem 3 *Suppose that (A.1) and (A.4) hold. Then the viscosity solution v of (6) with $v(0, \cdot) = v_0(\cdot)$ is unique in $C([0, \infty) \times \mathbf{R})$.*

From Theorems 2 and 3, we also obtain

Corollary 1 *Suppose that (A.1) and (A.2)' hold and that h is continuously differentiable. Then U in Theorem 2 is the unique continuous viscosity solution of (6) with $U(0, x) = \int_0^x h(y)dy$ in $(0, \infty) \times \mathbf{R}$.*

In particular, we have

Corollary 2 *Suppose that (A.1) and (A.4, i) hold and that Dv_0 is convex. Then (6) with $v(0, \cdot) = v_0(\cdot)$ has the unique viscosity solution in $C([0, \infty) \times \mathbf{R})$.*

(Proof of Theorem 1). The proof can be done almost in the same way as in [5, Theorem 1] (In [5, (A.3)], " $x \in \mathbf{R}^d$ " should be " $(x, \hat{h}(x)) \in \mathbf{R}^{d+1}$ "). The only thing we have to prove is the following (i)-(ii):

- (i) For a convex $u \in C(\mathbf{R})$, $w(P, u, dx)$ is completely additive,
- (ii) For $u \in C(\mathbf{R})$ for which $\partial u(\mathbf{R}) \neq \emptyset$, $w(P, u, dx) = w(P, \hat{u}, dx)$.

We first prove (i). The set

$$S(u) := \{p \in \mathbf{R} \mid \{x \in \mathbf{R} \mid p \in \partial u(x)\} \text{ contains more than one point}\}$$

contains at most countably many points. Indeed, if $p \in S(u)$, then for x_p for which $p \in \partial u(x_p)$, the graph of $y = u(x)$ and the straight line $y = p(x - x_p) + u(x_p)$ contains a line segment with a positive length and the interiors of such line segments are disjoint. Hence for each $n, m \geq 1$, on the set $S(u) \cap [D_-u(-n), D_+u(n)]$, such line segments with the length $\geq 1/m$ are finitely many. Here $D_-u(x)$ denotes the left derivative of $x \mapsto u(x)$. Since

a continuous measure does not have a point mass, we obtain (i) (see [1, p. 118]).

Next we prove (ii). If $u(x) = \hat{u}(x)$, then $\partial u(x) = \partial \hat{u}(x)$. If $u(x) \neq \hat{u}(x)$, then $\partial u(x) = \emptyset$ and $\partial \hat{u}(x) = D\hat{u}(x) \in S(\hat{u})$. Since $P(S(\hat{u})) = 0$ as we explained above, we obtain (ii). $\square$

(Proof of Theorem 2). (A.2)' implies that $x \mapsto DU(t, x)$ is convex for all $t \geq 0$ (see [5, Theorem 2]). We first prove that U is a viscosity subsolution to (6) in $(0, \infty) \times \mathbf{R}$. Suppose that $\varphi \in C^{1,2}((0, \infty) \times \mathbf{R})$, $(s, y) \in (0, \infty) \times \mathbf{R}$, and $U - \varphi$ attains a local maximum at (s, y) . Then for x and $y \in \mathbf{R}$ for which $x - y$ is positive,

$$\begin{aligned} \partial_t \varphi(s, y) &\leq F\left(\frac{U(s, x) - U(s, y) - DU(s, y)(x - y)}{(x - y)^2/2}\right) \\ &\leq F\left(\frac{\varphi(s, x) - \varphi(s, y) - D\varphi(s, y)(x - y)}{(x - y)^2/2}\right) \\ &\rightarrow F(D^2\varphi(s, y)) \quad (x \downarrow y). \end{aligned} \quad (11)$$

Indeed, from (5), for t and $s \geq 0$ for which $s - t$ is positive and is sufficiently small,

$$\varphi(s, y) - \varphi(t, y) \leq U(s, y) - U(t, y) = \int_t^s F(D_+(DU(\alpha, y)))d\alpha,$$

$$\begin{aligned} &U(\alpha, x) - U(\alpha, y) - DU(\alpha, y)(x - y) \\ &= \int_y^x (DU(\alpha, z) - DU(\alpha, y))dz \\ &\geq \int_y^x D_+(DU(\alpha, y))(z - y)dz = D_+(DU(\alpha, y))(x - y)^2/2. \end{aligned}$$

Since U and $DU \in C([0, \infty) \times \mathbf{R})$ from Theorem 1, we obtain the first inequality in (11). Since $DU(s, y) = D\varphi(s, y)$ and F is nondecreasing, the second inequality of (11) holds.

Next we prove that U is a viscosity supersolution to (6) in $(0, \infty) \times \mathbf{R}$. Suppose that $\varphi \in C^{1,2}((0, \infty) \times \mathbf{R})$, $(s, y) \in (0, \infty) \times \mathbf{R}$, and $U - \varphi$ attains a local minimum at (s, y) . Then in the same way as in (11), for x and $y \in \mathbf{R}$ for which $y - x$ is positive,

$$\begin{aligned} \partial_t \varphi(s, y) &\geq F\left(\frac{U(s, x) - U(s, y) - DU(s, y)(x - y)}{(x - y)^2/2}\right) \\ &\geq F\left(\frac{\varphi(s, x) - \varphi(s, y) - D\varphi(s, y)(x - y)}{(x - y)^2/2}\right) \\ &\rightarrow F(D^2\varphi(s, y)) \quad (x \uparrow y). \square \end{aligned}$$

(Proof of Proposition 1) Without loss of generality, we can put $s = 0$. We first prove the first inequality of (7). Suppose that there exists $(t_0, x_0) \in (0, \infty) \times \mathbf{R}$ such that

$$v(t_0, x_0) - v(0, x_0) < 0. \quad (12)$$

Put

$$\begin{aligned} C_0 &:= \min\{v(t, x) - v(0, x) \mid 0 \leq t \leq t_0, |x - x_0| \leq 1\} (< 0), \quad (13) \\ \underline{v}(t, x) &:= v(0, x) + C_0(x - x_0)^2. \end{aligned}$$

Then it is easy to see that $\underline{v}$ is a viscosity subsolution of (6) in $(0, t_0] \times (x_0 - 1, x_0 + 1)$ since $F \geq 0$. By the maximum principle (see [4, p.244, Th. 8.1]),

$$\begin{aligned} &\min\{v(t, x) - \underline{v}(t, x) \mid 0 \leq t \leq t_0, |x - x_0| \leq 1\} \\ &= \min\{v(t, x) - \underline{v}(t, x) \mid 0 \leq t \leq t_0, |x - x_0| = 1 \text{ or } t = 0, |x - x_0| \leq 1\} \\ &\geq 0, \end{aligned} \quad (14)$$

from (13). This contradicts (12).

Next we prove the second inequality of (7). Suppose that there exists $(\bar{t}_0, \bar{x}_0) \in (0, \infty) \times \mathbf{R}$ such that

$$v(\bar{t}_0, \bar{x}_0) - v(0, \bar{x}_0) - \bar{t}_0 > 0. \quad (15)$$

Put

$$\begin{aligned} \bar{C}_0 &:= \max\{v(t, x) - v(0, x) - t \mid 0 \leq t \leq \bar{t}_0, |x - \bar{x}_0| \leq 1\} (> 0), \\ \bar{v}(t, x) &:= v(0, x) + t + \bar{C}_0(x - \bar{x}_0)^2. \end{aligned} \quad (16)$$

Then it is easy to see that $\bar{v}$ is a viscosity supersolution of (6) in $(0, \bar{t}_0] \times (\bar{x}_0 - 1, \bar{x}_0 + 1)$ since $F \leq 1$. By the maximum principle,

$$\begin{aligned} &\max\{v(t, x) - \bar{v}(t, x) \mid 0 \leq t \leq \bar{t}_0, |x - \bar{x}_0| \leq 1\} \\ &= \max\{v(t, x) - \bar{v}(t, x) \mid 0 \leq t \leq \bar{t}_0, |x - \bar{x}_0| = 1 \text{ or } t = 0, |x - \bar{x}_0| \leq 1\} \\ &\leq 0, \end{aligned} \quad (17)$$

from (16). This contradicts (15). $\square$

(Proof of Proposition 2) First of all, we prove the first inequality in (8). Suppose that there exists $(t_0, x_0) \in (0, \infty) \times \mathbf{R}$ for which $F(a)t_0 > v(t_0, x_0) - v(0, x_0)$. Take $\varepsilon_0 > 0$ so that

$$v(t_0, x_0) - (v(0, x_0) + F(a)t_0 - \varepsilon_0 t_0) < 0. \quad (18)$$

For $n \geq 1$, put

$$\begin{aligned} C_n &:= \min\{v(t, x) - (v(0, x) + F(a)t) \mid 0 \leq t \leq t_0, |x - x_0| \leq n\}, \\ \psi_n(t, x) &:= v(t, x) - \left(v(0, x) + F(a)t - \varepsilon_0 t + \frac{C_n(x - x_0)^2}{n^2}\right). \end{aligned} \quad (19)$$

Then from (18)-(19),

$$\begin{aligned} & \min\{\psi_n(t, x) | 0 \leq t \leq t_0, |x - x_0| \leq n\} \\ &= \min\{\psi_n(t, x) | 0 < t \leq t_0, |x - x_0| < n\}. \end{aligned} \quad (20)$$

Take $(t_n, x_n) \in (0, t_0] \times (x_0 - n, x_0 + n)$ which attains the minimum in (20). Since v is a viscosity supersolution of (6) and since $|C_n| \leq t_0$ from Prop.1,

$$F(a) - \varepsilon_0 \geq F\left(D^2v(0, x_n) + \frac{2C_n}{n^2}\right) \geq F\left(a + \frac{2C_n}{n^2}\right) \rightarrow F(a), \quad (21)$$

as $n \rightarrow \infty$, which is a contradiction.

Next we prove the second inequality in (8). Suppose that there exists $(\bar{t}_0, \bar{x}_0) \in (0, \infty) \times \mathbf{R}$ for which $F(b)\bar{t}_0 < v(\bar{t}_0, \bar{x}_0) - v(0, \bar{x}_0)$. Take $\bar{\varepsilon}_0 > 0$ so that

$$v(\bar{t}_0, \bar{x}_0) - (v(0, \bar{x}_0) + F(b)\bar{t}_0 + \bar{\varepsilon}_0\bar{t}_0) > 0. \quad (22)$$

For $n \geq 1$, put

$$\begin{aligned} \bar{C}_n &:= \max\{v(t, x) - (v(0, x) + F(b)t) | 0 \leq t \leq \bar{t}_0, |x - \bar{x}_0| \leq n\}, \\ \bar{\psi}_n(t, x) &:= v(t, x) - \left(v(0, x) + F(b)t + \bar{\varepsilon}_0 t + \frac{\bar{C}_n(x - \bar{x}_0)^2}{n^2}\right). \end{aligned} \quad (23)$$

Then from (22)-(23),

$$\begin{aligned} & \max\{\bar{\psi}_n(t, x) | 0 \leq t \leq \bar{t}_0, |x - \bar{x}_0| \leq n\} \\ &= \max\{\bar{\psi}_n(t, x) | 0 < t \leq \bar{t}_0, |x - \bar{x}_0| < n\}. \end{aligned} \quad (24)$$

Take $(\bar{t}_n, \bar{x}_n) \in (0, \bar{t}_0] \times (\bar{x}_0 - n, \bar{x}_0 + n)$ which attains the maximum in (24). Since v is a viscosity subsolution of (6) and since $|\bar{C}_n| \leq \bar{t}_0$ from Prop.1,

$$F(b) + \bar{\varepsilon}_0 \leq F\left(D^2v(0, \bar{x}_n) + \frac{2\bar{C}_n}{n^2}\right) \leq F\left(b + \frac{2\bar{C}_n}{n^2}\right) \rightarrow F(b), \quad (25)$$

as $n \rightarrow \infty$, which is a contradiction.

Next we prove (9)-(10). From (8), we only have to prove the following:

$$\limsup_{x \rightarrow -\infty} \left(\sup_{0 \leq t \leq T} \{v(t, x) - v(0, x) - F(a)t\} \right) \leq 0, \quad (26)$$

$$\liminf_{x \rightarrow \infty} \left(\inf_{0 \leq t \leq T} \{v(t, x) - v(0, x) - F(b)t\} \right) \geq 0. \quad (27)$$

We first prove (26). Suppose that (26) does not hold. Then there exists $\varepsilon_1 > 0$ so that

$$\limsup_{x \rightarrow -\infty} \left(\sup_{0 \leq t \leq T} \{v(t, x) - v(0, x) - F(a)t - \varepsilon_1 t\} \right) > 0. \quad (28)$$

In particular, there exists $(s_n, y_n) \in (0, T] \times (-\infty, -n^2)$ for which

$$v(s_n, y_n) - v(0, y_n) - F(a)s_n - \varepsilon_1 s_n > 0 \quad (n \geq 1). \quad (29)$$

For $n \geq 1$, put

$$\begin{aligned} \gamma_n &:= \max\{v(t, x) - (v(0, x) + F(a)t) \mid 0 \leq t \leq T, |x - y_n| \leq n\}, \\ \phi_n(t, x) &:= v(t, x) - \left(v(0, x) + F(a)t + \varepsilon_1 t + \frac{\gamma_n(x - y_n)^2}{n^2} \right). \end{aligned} \quad (30)$$

Then from (29)-(30).

$$\begin{aligned} &\max\{\phi_n(t, x) \mid 0 \leq t \leq T, |x - y_n| \leq n\} \\ &= \max\{\phi_n(t, x) \mid 0 < t \leq T, |x - y_n| < n\}. \end{aligned} \quad (31)$$

Take $(r_n, z_n) \in (0, T] \times (y_n - n, y_n + n)$ which attains the maximum in (31). Since v is a viscosity subsolution of (6), $z_n < -n^2 + n \rightarrow -\infty$ as $n \rightarrow \infty$ and $|\gamma_n| \leq T$ from Prop. 1,

$$F(a) + \varepsilon_1 \leq F\left(D^2v(0, z_n) + \frac{2\gamma_n}{n^2}\right) \rightarrow F(a) \quad (n \rightarrow \infty), \quad (32)$$

which is a contradiction.

Next we prove (27). Suppose that (27) does not hold. Then there exists $\bar{\varepsilon}_1 > 0$ so that

$$\liminf_{x \rightarrow \infty} \left(\inf_{0 \leq t \leq T} \{v(t, x) - v(0, x) - F(b)t + \bar{\varepsilon}_1 t\} \right) < 0. \quad (33)$$

In particular, there exists $(\bar{s}_n, \bar{y}_n) \in (0, T] \times (n^2, \infty)$ for which

$$v(\bar{s}_n, \bar{y}_n) - v(0, \bar{y}_n) - F(b)\bar{s}_n + \bar{\varepsilon}_1 \bar{s}_n < 0 \quad (n \geq 1). \quad (34)$$

Put

$$\begin{aligned} \bar{\gamma}_n &:= \min\{v(t, x) - (v(0, x) + F(b)t) | 0 \leq t \leq T, |x - \bar{y}_n| \leq n\}, \\ \bar{\phi}_n(t, x) &:= v(t, x) - \left(v(0, x) + F(b)t - \bar{\varepsilon}_1 t + \frac{\bar{\gamma}_n(x - \bar{y}_n)^2}{n^2}\right). \end{aligned} \quad (35)$$

Then from (34)-(35),

$$\begin{aligned} &\min\{\bar{\phi}_n(t, x) | 0 \leq t \leq T, |x - \bar{y}_n| \leq n\} \\ &= \min\{\bar{\phi}_n(t, x) | 0 < t \leq T, |x - \bar{y}_n| < n\}. \end{aligned} \quad (36)$$

Take $(\bar{r}_n, \bar{z}_n) \in (0, T] \times (\bar{y}_n - n, \bar{y}_n + n)$ which attains the minimum in (36). Since v is a viscosity supersolution of (6), $\bar{z}_n > n^2 - n \rightarrow \infty$ as $n \rightarrow \infty$ and $|\bar{\gamma}_n| \leq T$ from Prop. 1,

$$F(b) - \bar{\varepsilon}_1 \geq F\left(D^2v(0, \bar{z}_n) + \frac{2\bar{\gamma}_n}{n^2}\right) \rightarrow F(b) \quad (n \rightarrow \infty), \quad (37)$$

which is a contradiction. $\square$

(Acknowledgement) This research is supported in part by the Grant-in-Aid for Scientific Research, No.16654031, JSPS.

References

- [1] I. J. BAKELMAN, *Convex Analysis and Nonlinear Geometric Elliptic Equations*, Springer-Verlag, Berlin Heidelberg New York, 1994.
- [2] M. G. CRANDALL, H. ISHII, AND P.-L. LIONS, *User's guide to viscosity solutions of second order partial differential equations*, Bull. Amer. Math. Soc. (N.S.), 27(1992), pp. 1–67.
- [3] W. J. FIREY, *Shapes of worn stones*, Mathematika, 21(1974), pp. 1–11.
- [4] W. H. FLEMING AND H. M. SONER, *Controlled Markov processes and Viscosity solutions*, Springer-Verlag, , Berlin · Heidelberg · New York · Tokyo, 1993.
- [5] H. ISHII AND T. MIKAMI, *Motion of a graph by R-curvature*, Arch. Ration. Mech. Anal. **171** (2004) pp. 1-23. Correction: <http://www.math.sci.hokudai.ac.jp/mikami/ARMAcorrect.pdf>
- [6] H. ISHII AND T. MIKAMI, *Convexified Gauss Curvature Flow of Sets: A Stochastic Approximation*, SIAM J. MATH. ANAL. **36** (2004) 552-579.