

Uniformly locally o-minimal structures of the second kind and their tame topology

海上保安大学校* 藤田雅人

Masato Fujita

Liberal Arts, Japan Coast Guard Academy

概要

This paper is a brief survey on a uniformly locally o-minimal structure of the second kind for non-specialists of model theory. An o-minimal structure enjoys tame topological properties such as monotonicity theorem and definable cell decomposition theorem. A uniformly locally o-minimal structure of the second kind is a new variant of o-minimal structure. A uniformly locally o-minimal structure of the second kind enjoys the local versions of tame topological properties possessed by an o-minimal structure. It enables to develop a tame dimension theory for definable sets.

1 Introduction

This paper is a survey on a uniformly locally o-minimal structure of the second kind for non-specialists of model theory. The definition of a structure given here is slightly different from the original definition in model theory. A reader who has interest in model theory should consult textbooks such as [1, 17, 19].

The notation $\mathbb{N}$ denotes the set of positive integers. In this paper, a *structure* is a pair $\mathcal{M} = (M, \mathfrak{S} = \{\mathfrak{S}_n\}_{n \in \mathbb{N}})$ of a set M and the collection $\mathfrak{S}$ of families $\mathfrak{S}_n$ of subsets of M^n satisfying the following conditions:

- (i) The empty set and M^n are members of $\mathfrak{S}_n$ for all $n \in \mathbb{N}$. The set $\{(x, y) \in$

$M^2 \mid x = y\}$ is also a member of $\mathfrak{S}_2$.

- (ii) The families $\mathfrak{S}_n$ are closed under the boolean algebra for all $n \in \mathbb{N}$.
- (iii) The Cartesian product $S_1 \times S_2$ belongs to $\mathfrak{S}_{m+n}$ if S_1 and S_2 are members of $\mathfrak{S}_m$ and $\mathfrak{S}_n$, respectively.
- (iv) Let $\pi : M^n \rightarrow M^m$ be a coordinate projection and let X be a member of $\mathfrak{S}_n$. Then, the projection image $\pi(X)$ belongs to $\mathfrak{S}_m$.
- (v) Let σ be a permutation of $\{1, \dots, n\}$. We define the map $\bar{\sigma} : M^n \rightarrow M^n$ by $\bar{\sigma}(x_1, \dots, x_n) = (x_{\sigma(1)}, \dots, x_{\sigma(n)})$. We have $\bar{\sigma}(X) \in \mathfrak{S}_n$ if $X \in \mathfrak{S}_n$.

When a structure $\mathcal{M}$ is given, the set M is called the *universe* or the *underlying set* of the structure $\mathcal{M}$. Members in $\mathfrak{S}_n$ are called *definable sets*. Let X and Y be definable sets. A map $f : X \rightarrow Y$ is called *definable* if its graph is a definable set.

We sometimes need to consider the family of structures such that some sets other than those given in (i) are definable. When M is a densely linearly ordered set with the order $<$, a structure $\mathcal{M} = (M, \mathfrak{S})$ is called an *expansion* of a dense linear order if the set $\{(x, y) \mid x < y\}$ is definable. When $(M, \cdot)$ is a group, a structure $\mathcal{M}$ with the universe M is called an *expansion* of a group if the set $\{(x, y, z) \in M^3 \mid x \cdot y = z\}$ is definable. We define an expansion of an ordered group, an expansion of an ordered field and so on in the same manner.

An *o-minimal* structure $\mathcal{M} = (M, \mathfrak{S})$ is an expansion of a dense linear order without endpoints such that

- (vi) any definable subset of M is a finite union of points and open intervals.

An open interval is a subset of M of the form $\{x \in M \mid a < x < b\}$, where $a \in M \cup \{-\infty\}$ and $b \in M \cup \{+\infty\}$. Definable sets and definable maps in an o-minimal structures are well-behaved. For instance, for a unary definable function $f : M \rightarrow M$, the domain of definition M is decomposed into finite points and open intervals such that the restriction of f to the open intervals are monotone and continuous. It is called the monotonicity theorem. The definable cell decomposition theorem for o-minimal structures guarantees that any definable set is a finite union of good-shaped definable sets called ‘cells.’ Readers who are interested in o-minimal structures should consult van den Dries’s book [4] and Coste’s book [2]. The paper [5] is also recommended.

Many structures relaxing the condition (vi) are proposed and investigated such

as weakly o-minimal structures [16], locally o-minimal structures [21] and structures having (locally) o-minimal open cores [3, 6]. A locally o-minimal structure is defined by localizing the condition (vi). A *locally o-minimal structure* is an expansion of a dense linear order without endpoints satisfying the following condition:

- (vi)' Let X be a definable subset of M . For any $x \in M$, there exists an open interval I containing the point x such that $X \cap I$ is a finite union of points and open intervals.

Unfortunately, even a localized version of monotonicity theorem is unavailable in a general local o-minimal structure [21, Proposition 2.11]. This is the reason why the author proposed a uniformly locally o-minimal structure of the second kind in [8]. A local monotonicity theorem holds true in a uniformly locally o-minimal structure of the second kind [8, Corollary 3.1]. A definably complete locally o-minimal structure admits local definable cell decomposition if and only if it is a uniformly locally o-minimal structure of the second kind [8, Corollary 4.1]. This paper summarizes the results on uniformly locally o-minimal structures of the second kind including the above theorems. It is a survey paper, and does not give a new insight on uniformly locally o-minimal structures of the second kind.

This paper is organized as follows. We first define a uniformly locally o-minimal structure of the second kind and related structures in Section 2. Topology of definable sets in a uniformly locally o-minimal structure of the second kind is discussed in Section 3. We develop a dimension theory for definable sets in a uniformly locally o-minimal structure of the second kind in Section 4 using the results of Section 3. We conclude this paper with remarks in Section 5.

2 Basic Definitions

We first review the definitions given in [18, 21, 15, 8].

Definition 2.1. We consider an expansion $\mathcal{M} = (M, \mathfrak{S})$ of a dense linear order without endpoints. It is *definably complete* if every definable subset of M has both a supremum and an infimum in $M \cup \{\pm\infty\}$ [18]. A definably complete expansion of an ordered group is divisible and abelian [18, Proposition 2.2].

We review the definition of locally o-minimal structures. The structure $\mathcal{M}$ is *locally o-minimal* if, for every definable subset X of M and for every point $a \in M$, there exists an open interval I containing the point a such that $X \cap I$ is a finite union of points and open intervals. A locally o-minimal structure $\mathcal{M}$ is *strongly locally o-minimal* if, for every point $a \in M$, there exists an open interval I containing the point a such that $X \cap I$ is a finite union of points and open intervals for every definable subset X of M .

A locally o-minimal structure $\mathcal{M}$ is a *uniformly locally o-minimal structure of the first kind* if, for any positive integer n , any definable set $X \subset M^{n+1}$ and $a \in M$, there exists an open interval I containing the point a such that the definable sets $X_y \cap I$ are finite unions of points and open intervals for all $y \in M^n$. Here, X_y denotes the fiber $\{x \in M \mid (x, y) \in X\}$. A uniformly locally o-minimal structure of the first kind is called a uniformly locally o-minimal structure in [15].

A locally o-minimal structure $\mathcal{M}$ is a *uniformly locally o-minimal structure of the second kind* if, for any positive integer n , any definable set $X \subset M^{n+1}$, $a \in M$ and $b \in M^n$, there exist an open interval I containing the point a and an open box B containing b such that the definable sets $X_y \cap I$ are finite unions of points and open intervals for all $y \in B$.

We frequently consider a definably complete uniformly locally o-minimal expansion of the second kind of an ordered group. We simply call it a *DCULOAS structure* in this paper.

A locally o-minimal structure whose universe is the set of reals $\mathbb{R}$ is strongly locally o-minimal [21, Corollary 3.4]. A strongly locally o-minimal structure is always a uniformly locally o-minimal structure of the first kind. But the converse is not true in general. A definably complete uniformly locally o-minimal structure of the first kind which is not strongly o-minimal is found in [8, Example 2.4]. A uniformly locally o-minimal structure of the first kind is a uniformly locally o-minimal structure of the second kind. The converse is not true, neither. A counterexample is [8, Example 2.3]. A locally o-minimal structure is not necessarily a uniformly locally o-minimal structure of the second kind. Its counterexample is [8, Example 2.2].

The following proposition indicates that it is futile to consider a uniformly locally o-minimal expansion of the second kind of an ordered field.

Proposition 2.2 ([8, Proposition 2.1]). *A uniformly locally o-minimal expansion of the second kind of an ordered field is o-minimal.*

3 Tame topology

The following is the local monotonicity theorem for uniformly locally o-minimal structure of the second kind.

Theorem 3.1 (Local monotonicity theorem). *Consider a uniformly locally o-minimal structure of the second kind $\mathcal{M} = (M, \mathfrak{S})$. Let I be an interval and $f : I \rightarrow M$ be a definable function. For any $(a, b) \in M^2$, there exist an open interval J_1 containing the point a , an open interval J_2 containing the point b and a mutually disjoint definable partition*

$$f^{-1}(J_2) \cap J_1 = X_d \cup X_c \cup X_+ \cup X_-$$

satisfying the following conditions:

- (1) *the definable set X_d is discrete and closed;*
- (2) *the definable set X_c is open and f is locally constant on X_c ;*
- (3) *the definable set X_+ is open and f is locally strictly increasing and continuous on X_+ ;*
- (4) *the definable set X_- is open and f is locally strictly decreasing and continuous on X_- .*

Furthermore, if the structure $\mathcal{M}$ is a DCULOAS structure, we can choose $J_1 = I$ and $J_2 = M$.

This theorem is first proved in [15, Proposition 11] only for strongly locally o-minimal structures. For any point $a \in M^n$, there exists an open box B such that the intersection of B with a definable subset of M^n in the given strongly locally o-minimal structure is a definable subset in an o-minimal structure having the same universe [15, Theorem 9]. The above theorem for strongly locally o-minimal structures is a direct corollary of it and the monotonicity theorem for o-minimal structures. Theorem 3.1 is found in [8, Corollary 3.1] and its proof is not so easy. The ‘furthermore’ part follows from [10, Theorem 2.11, Proposition 2.13].

We review the definitions of cells and definable cell decomposition.

Definition 3.2 (Definable cell decomposition). Consider an expansion of dense linear order $\mathcal{M} = (M, \mathfrak{S})$. Let $(i_1, \dots, i_n)$ be a sequence of zeros and ones of length n . $(i_1, \dots, i_n)$ -cells are definable subsets of M^n defined inductively as follows:

- A (0)-cell is a point in M and a (1)-cell is an open interval in M .
- An $(i_1, \dots, i_n, 0)$ -cell is the graph of a continuous definable function defined on an $(i_1, \dots, i_n)$ -cell. An $(i_1, \dots, i_n, 1)$ -cell is a definable set of the form $\{(x, y) \in C \times M \mid f(x) < y < g(x)\}$, where C is an $(i_1, \dots, i_n)$ -cell and f and g are definable continuous functions defined on C with $f < g$.

A *cell* is an $(i_1, \dots, i_n)$ -cell for some sequence $(i_1, \dots, i_n)$ of zeros and ones. An *open cell* is a $(1, 1, \dots, 1)$ -cell.

We inductively define a *definable cell decomposition* of an open box $B \subset M^n$. For $n = 1$, a definable cell decomposition of B is a partition $B = \bigcup_{i=1}^m C_i$ into finite cells. For $n > 1$, a definable cell decomposition of B is a partition $B = \bigcup_{i=1}^m C_i$ into finite cells such that $\pi(B) = \bigcup_{i=1}^m \pi(C_i)$ is also a definable cell decomposition of $\pi(B)$, where $\pi : M^n \rightarrow M^{n-1}$ is the projection forgetting the last coordinate. Given a finite family $\{A_\lambda\}_{\lambda \in \Lambda}$ of definable subsets of B , a *definable cell decomposition of B partitioning $\{A_\lambda\}_{\lambda \in \Lambda}$* is a definable cell decomposition of B such that the definable sets A_λ are unions of cells for all $\lambda \in \Lambda$.

In an o-minimal structure, global definable cell decomposition is available. It means that, for any finite family of definable subsets of M^n , there exists a definable cell decomposition of M^n partitioning the given family [4, Chapter 3, Cell decomposition theorem 2.11]. In a general local o-minimal structure, even local definable cell decomposition is unavailable. The following theorem says that it is available when the structure is a definably complete uniformly locally o-minimal structure of the second kind.

Theorem 3.3 (Local definable cell decomposition theorem). *Consider a strongly locally o-minimal structure or a definably complete uniformly locally o-minimal structure of the second kind $\mathcal{M} = (M, \mathfrak{S})$.*

Let $\{A_\lambda\}_{\lambda \in \Lambda}$ be a finite family of definable subsets of M^n . For any point $a \in M^n$,

there exist an open box B containing the point a and a definable cell decomposition of B partitioning the finite family $\{B \cap A_\lambda \mid \lambda \in \Lambda \text{ and } B \cap A_\lambda \neq \emptyset\}$.

This theorem is first proved in [15, Proposition 13] only for strongly locally o-minimal structures. It is also a direct corollary of [15, Theorem 9] and the definable cell decomposition theorem for o-minimal structures. The case in which the structure is a definably complete uniformly locally o-minimal structure of the second kind is [8, Theorem 4.2].

Definition 3.4. A locally o-minimal structure *admits local definable cell decomposition* if the assertion in Theorem 3.3 hold true for all positive integers n .

When the structure is definably complete, we can get the following important corollary:

Corollary 3.5 ([8, Corollary 4.1]). *A definably complete locally o-minimal structure admits local definable cell decomposition if and only if it is a uniformly locally o-minimal structure of the second kind.*

The local definable cell decomposition theorem (Theorem 3.3) cannot be extended to the global one. We want to decompose a definable set into finite good-shaped definable sets, which may not be as good-shaped as cells. One candidate is a quasi-special submanifold defined as follows:

Definition 3.6 (Quasi-special submanifolds). Consider an expansion of a densely linearly order without endpoints $\mathcal{M} = (M, \mathfrak{S} = \{\mathfrak{S}_n\}_{n \in \mathbb{N}})$. Let X be a definable subset of M^n and $\pi : M^n \rightarrow M^d$ be a coordinate projection. The definable set X is a π -*quasi-special submanifold* or simply a *quasi-special submanifold* if, $\pi(X)$ is a definable open set and, for every point $x \in \pi(X)$, there exists an open box U in M^d containing the point x satisfying the following condition: For any $y \in X \cap \pi^{-1}(x)$, there exist an open box V in M^n and a definable continuous map $\tau : U \rightarrow M^n$ such that $\pi(V) = U$, $\tau(U) = X \cap V$ and the composition $\pi \circ \tau$ is the identity map on U .

Let $\{X_i\}_{i=1}^m$ be a finite family of definable subsets of M^n . A *decomposition of M^n into quasi-special submanifolds partitioning $\{X_i\}_{i=1}^m$* is a finite family of quasi-special submanifolds $\{C_i\}_{i=1}^N$ such that

- $\bigcup_{i=1}^N C_i = M^n$,
- $C_i \cap C_j = \emptyset$ when $i \neq j$ and
- either C_i has an empty intersection with X_j or is contained in X_j for any $1 \leq i \leq m$ and $1 \leq j \leq N$.

A decomposition $\{C_i\}_{i=1}^N$ of M^n into quasi-special submanifolds *satisfies the frontier condition* if the closure of any quasi-special submanifold $\overline{C_i}$ is the union of a subfamily of the decomposition.

The following theorem says that any definable set is a disjoint union of finite quasi-special submanifolds.

Theorem 3.7 ([10, Theorem 4.5]). *Consider a DCULOAS structure $\mathcal{M} = (M, \mathfrak{S})$. Let $\{X_i\}_{i=1}^m$ be a finite family of definable subsets of M^n . There exists a decomposition $\{C_i\}_{i=1}^N$ of M^n into quasi-special submanifolds partitioning $\{X_i\}_{i=1}^m$ and satisfying the frontier condition. Furthermore, the number N of quasi-special submanifolds in the decomposition is not greater than the number uniquely determined only by m and n .*

4 Dimension theory

Assuming that the considered structure admits local definable cell decomposition, we can develop a good dimension theory. We first define the dimension of a definable set as follows.

Definition 4.1 (Dimension of a definable set). Consider an expansion of dense linear order $\mathcal{M} = (M, \mathfrak{S} = \{\mathfrak{S}_n\}_{n \in \mathbb{N}})$. A definable set $X \subset M^n$ is of $\dim(X) \geq m$ if there exists an open box $B \subset M^m$ and a definable continuous injective map $f : B \rightarrow X$ which is homeomorphic onto its image. A definable set $X \subset M^n$ is of $\dim(X) = m$ if it is of $\dim(X) \geq m$ and it is not of $\dim(X) \geq m + 1$. The empty set is defined to be of dimension $-\infty$.

The following corollary gives equivalent definitions of dimension.

Corollary 4.2 ([8, Corollary 5.3]). *Consider a locally o-minimal structure $\mathcal{M} = (M, \mathfrak{S} = \{\mathfrak{S}_n\}_{n \in \mathbb{N}})$ which admits local definable cell decomposition. The following*

conditions are equivalent for any definable subset $X \subset M^n$:

- $\dim(X) \geq m$;
- the definable set X contains an $(i_1, \dots, i_n)$ -cell with $\sum_{j=1}^n i_j \geq m$, and
- there exist a coordinate projection $\pi : M^n \rightarrow M^m$ and a point $a \in M^n$ such that the definable set $\pi(B \cap X)$ has a nonempty interior for any open box B containing the point a .

The dimension defined above possesses the good features which we naturally expect as in Theorem 4.3 and Theorem 4.5.

Theorem 4.3 ([8, Lemma 5.1, Corollary 5.4, Theorem 5.6]). *Consider a locally o-minimal structure $\mathcal{M} = (M, \mathfrak{S} = \{\mathfrak{S}_n\}_{n \in \mathbb{N}})$ which admits local definable cell decomposition. The following assertions hold true:*

- (1) *Let $X \subset Y$ be definable sets. Then, the inequality $\dim(X) \leq \dim(Y)$ holds true.*
- (2) *Let σ be a permutation of the set $\{1, \dots, n\}$. The definable map $\bar{\sigma} : M^n \rightarrow M^n$ is defined by $\bar{\sigma}(x_1, \dots, x_n) = (x_{\sigma(1)}, \dots, x_{\sigma(n)})$. Then, we have $\dim(X) = \dim(\bar{\sigma}(X))$ for any definable subset X of M^n .*
- (3) *Let X and Y be definable sets. We have $\dim(X \times Y) = \dim(X) + \dim(Y)$.*
- (4) *Let X and Y be definable subsets of M^n . We have*

$$\dim(X \cup Y) = \max\{\dim(X), \dim(Y)\}.$$

- (5) *Let X be a definable set. The notation ∂X denotes the frontier of X defined by $\partial X = \overline{X} \setminus X$. We have $\dim(\partial X) < \dim X$.*

In the course of the proof of Theorem 4.5, we demonstrate the following strong definable Baire property, which is a definable variant of the famous Baire property.

Proposition 4.4 (Strong definable Baire property, [9, Theorem 4.3]). *Consider a DCULOAS structure $\mathcal{M} = (M, \mathfrak{S} = \{\mathfrak{S}_n\}_{n \in \mathbb{N}})$. Take $c \in M$. Let $\{X\langle r \rangle\}_{r > c}$ be a parameterized increasing family of definable sets of M^n ; that is, there exists a definable subset $\mathcal{X}$ of M^{n+1} such that $X\langle r \rangle$ coincides with the fiber $\mathcal{X}_r$ for any $r > c$ and we have $X\langle r \rangle \subset X\langle s \rangle$ if $r < s$. Set $X = \bigcup_{r > c} X\langle r \rangle$. The definable set $X\langle r \rangle$ has a nonempty interior for some $r > c$ if X has a nonempty interior.*

Structures satisfying a weaker definable Baire property are discussed in [7, 14]

Theorem 4.5 ([9, Theorem 1.1, Corollary 1.2], [10, Theorem 3.14]). *Consider a DCULOAS structure $\mathcal{M} = (M, \mathfrak{S} = \{\mathfrak{S}_n\}_{n \in \mathbb{N}})$. The following assertions hold true:*

- (1) *Let $f : X \rightarrow M^n$ be a definable map. We have $\dim(f(X)) \leq \dim X$.*
- (2) *Let $f : X \rightarrow M^n$ be a definable map. The notation $\mathcal{D}(f)$ denotes the set of points at which the map f is discontinuous. We have $\dim \mathcal{D}(f) < \dim X$.*
- (3) *(Addition Property) Let $\varphi : X \rightarrow Y$ be a definable surjective map whose fibers are equi-dimensional; that is, the dimensions of the fibers $\varphi^{-1}(y)$ are constant. We have $\dim X = \dim Y + \dim \varphi^{-1}(y)$ for all $y \in Y$.*

5 Remarks

We conclude this paper with several remarks. The most restrictive structure considered in this paper is a DCULOAS structure. However, by [10], all the assertions except the local definable cell decomposition theorem (Theorem 3.3) are satisfied in any definably complete locally o-minimal structure such that

- (*) the image of a discrete definable set under a coordinate projection is again discrete.

A DCULOAS structure satisfies the condition (*). Shoutens proposed a locally o-minimal structure called a *model of DCTC* [20]. A model of DCTC is a definably complete locally o-minimal structure with the property (*). A locally o-minimal expansion of an ordered field falls into a model of DCTC.

A natural unsolved question is as follows:

Conjecture. Does any definably complete locally o-minimal structure satisfy the property (*)?

The strongly locally o-minimal structure in [15, Example 12] is definably complete nor satisfies the property (*), neither.

Since a DCULOAS structure has tame topological properties, definable functions are also expected to have tame properties. Definable equi-continuity is defined and investigated in [11]. A variant of the Arzela-Ascoli theorem is demonstrated in the

same paper.

Consider two structures $\mathcal{M} = (M, \mathfrak{S} = \{\mathfrak{S}_n\}_{n \in \mathbb{N}})$ and $\mathcal{M}' = (M, \mathfrak{S}' = \{\mathfrak{S}'_n\}_{n \in \mathbb{N}})$ having the same universe M . When $\mathfrak{S}$ is a subset of $\mathfrak{S}'$, $\mathcal{M}$ is called a *reduct* of $\mathcal{M}'$, and $\mathcal{M}'$ is called an *expansion* of $\mathcal{M}$. For a given structure $\mathcal{M}$, the reduct generated by the open sets definable in $\mathcal{M}$ is called the *open core* of $\mathcal{M}$. A sufficient condition for a structure having an o-minimal open core is discussed in [3]. Definably complete expansions of ordered fields having locally o-minimal open cores are treated in [6]. The author gave a sufficient condition for a structure having uniformly locally o-minimal open core of the first/second kind in [12].

A locally o-minimal structure whose universe is the set of reals $\mathbb{R}$ is strongly o-minimal. It enjoys more tame condition called almost o-minimality.

Definition 5.1. An expansion of densely linearly ordered set without endpoints is *almost o-minimal* if any bounded definable set in M is a finite union of points and open intervals. Here, M is the universe of the expansion.

The notion of almost o-minimality was formulated in [13].

参考文献

- [1] C. C. Chang and H. J. Keisler, Model theory 3rd eds., Studies in logic and the foundation of mathematics, vol. 73, North-Holland Press, Amsterdam, 1990.
- [2] M. Coste, An introduction to o-minimal geometry, Dip. Mat. Univ. Pisa, Dottorato di Ricerca in Matematica, Istituti Editoriali e Poligrafici Internazionali, Pisa, 2000.
- [3] A. Dolich, C. Miller and C. Steinhorn, Structure having o-minimal open core, Trans. Amer. Math. Soc., 362 (2010), 1371-1411.
- [4] L. van den Dries, Tame topology and o-minimal structures, London Mathematical Society Lecture Note Series, vol. 248. Cambridge University Press, Cambridge, 1998.
- [5] L. van den Dries and C. Miller, *Geometric categories and o-minimal structures*, Duke Math. J. **84** (1996), 497-540.
- [6] A. Fornasiero, Locally o-minimal structures and structures with locally o-minimal open core, Ann. Pure Appl. Logic, 164 (2013), 211-229.

- [7] A. Fornasiero and T. Servi, Definably complete Baire structure, *Fund. Math.*, 209 (2010), 215-241.
- [8] M. Fujita, Uniformly locally o-minimal structures and locally o-minimal structures admitting local definable cell decomposition, *Ann. Pure Appl. Logic*, 171 (2020), 102756.
- [9] M. Fujita, Dimension inequality for a definably complete uniformly locally o-minimal structure of the second kind, *J. Symbolic Logic*, 85 (2020), 1654-1663.
- [10] M. Fujita, Locally o-minimal structures with tame topological properties, preprint, arXiv:2006.13481v3.
- [11] M. Fujita, Functions definable in definably complete uniformly locally o-minimal structure of the second kind, preprint, arXiv:2010.02420v2.
- [12] M. Fujita, Uniformly locally o-minimal open core, preprint, arXiv:2002.03077v2
- [13] M. Fujita, Almost o-minimal structures and $\mathfrak{X}$ -structures, preprint, arXiv:2104.01312v1
- [14] P. Hieronymi, An analogue of the Baire category theorem, *J. Symbolic Logic*, 78 (2013), 207-213.
- [15] T. Kawakami, K. Takeuchi, H. Tanaka and A. Tsuboi, Locally o-minimal structures, *J. Math. Soc. Japan*, 64 (2012), 783-797.
- [16] D. Macpherson, D. Marker and C. Steinhorn, Weakly o-minimal structures and real closed fields, *Trans. Amer. Math. Soc.*, 352 (2000), 5435-5483.
- [17] D. Marker, *Model theory: An introduction*, Graduate texts in mathematics, vol. 217, Springer-Verlag, New York, 2002.
- [18] C. Miller, Expansions of dense linear orders with the intermediate value property, *J. Symbolic Logic*, 66 (2001), 1783-1790.
- [19] B. Poizat, *A course in model theory: An introduction to contemporary mathematical logic*, Universitext, Springer-Verlag, New York, 2000.
- [20] H. Schoutens, O-minimalism, *J. Symbolic Logic*, 79 (2014), 355-409.
- [21] C. Toffalori and K. Vozoris, Notes on local o-minimality, *MLQ Math. Log. Q.*, 55 (2009), 617-632.