

Investigation of Luttinger liquids with DMRG and TEBD methods

Eric Jeckelmann, Alex Cojuhovski, Malcolm Einhellinger, Martin Paech

Institute for Theoretical Physics, Leibniz Universität Hannover, 30167 Hannover,
GERMANY

I will discuss two numerical approaches based on matrix-product states which can be used to investigate the dynamical properties of quasi-one-dimensional strongly correlated quantum many-body systems. Both methods are illustrated with studies of the spectral functions and transport properties of lattice models in a Luttinger liquid phase. The first approach is based on the dynamical density-matrix renormalization group (DMRG) method [1] and allows us to compute frequency- and momentum-resolved spectral functions. systems [2, 3] as well as the Drude weight [4]. I will present a method for determining the properties of infinite systems from the DDMRG data calculated for large but finite systems. The second approach is based on the time-evolving block decimation (TEBD) method and allows us to compute the time evolution of a system out of equilibrium. I will show that steady-state transport properties (e.g. the conductance of Luttinger liquid quantum wires) can be determined from TEBD simulations of finite-size chains.

References

- [1] E. Jeckelmann, Phys. Rev. B **66** (2002), 045114.
- [2] H. Benthien, F. Gebhard, and E. Jeckelmann, Phys. Rev. Lett. **92** (2004), 256401.
- [3] E. Jeckelmann, Progress of Theoretical Physics Supplement **176**, (2008), 143.
- [4] T. Shirakawa and E. Jeckelmann, Phys. Rev. B **79** (2009), 195121.