数理解析研究所講究録
1511 巻 2006 年 92-97



# $S$-Unit Equations and Integer Solutions to Exponential Diophantine Equations

Noriko HIRATA-KOHNO

October 19, 2004

# $S$単数方程式と指数方程式の整数解

日本大学理工学部数学科
平田典子

**Abstract**

In this article, we present some new applications of unit equations and linear forms in logarithms to obtain a simple upper bound for the number of the purely exponential Diophantine equations. The main idea essentially relies on a refined result of a bound for the number of the solutions to $S$-unit equations, due to F. Beukers and H. P. Schlickewei as well as that by J. -H. Evertse, H. P. Schlickewei and W.M. Schmidt [Be-Schl] [E-Schl-Schm]. The tool to obtain a bound for the size of the solutions is the theory of linear forms in $m$-adic logarithms where $m$ denotes a positive integer not necessarily a prime.

Keywords: Diophantine approximation, Unit equation, Linear forms in logarithms, Exponential Diophantine equations.

## 1 Introduction

Let us denote by $\mathbb{Z}$ the set of the rational integers. Let $a, b, c \in \mathbb{Z}$ where $a, b, c \geq 2$ and $(a, b, c) = 1$.

Consider the exponential Diophantine equation

$$a^x + b^y = c^z \tag{1}$$

in unknowns $a, b, c, x, y, z \in \mathbb{Z}$, $x, y, z \geq 1$.

In this case, we see $(a, b, c) = 1 \Longleftrightarrow (a, b) = 1 \Longleftrightarrow (a, c) = 1 \Longleftrightarrow (b, c) = 1$.

Let us recall a conjecture due to Tijdeman (sometimes called Beal's conjecture):

**Conjecture 1.** *(Tijdeman) The equation $a^x+b^y = c^z$ has no solutions in $(a,b,c,x,y,z) \in \mathbb{Z}^6$ with $a,b,c \geq 2, x,y,z \geq 3$.*

The equation in the conjecture concerns 6 unknowns. It is known that the *abc*-conjecture of Masser-Osterlé type implies that there is an effective positive number $H$ which depends only on the $\varepsilon > 0$ in the *abc*-conjecture such that Conjecture 1 is true for $x,y,z \geq H$.

It is also investigated by Darmon-Granville, Darmon-Merel, Kraus, Bennett and others that the number of the solutions $a,b,c$ to (1) is finite if $x,y,z$ are fixed with $\frac{1}{x}+\frac{1}{y}+\frac{1}{z} < 1$.

When we consider again the six numbers as unknowns, a slightly different question is asked;

**Conjecture 2.** *(Fermat-Catalan) If $\frac{1}{x}+\frac{1}{y}+\frac{1}{z} < 1$ then the number of the solutions in $(a,b,c,x,y,z) \in \mathbb{Z}^6$ with $a,b,c \geq 2, x,y,z \geq 2$ is finite.*

For example some solutions to the equation of Conjecture 2 including large ones found by Beukers-Zagier are as follows.

**Example 1.** $2^5 + 7^2 = 3^4$
$7^3 + 13^2 = 2^9$
$2^7 + 17^3 = 71^2$
$3^5 + 11^4 = 122^2$
$17^7 + 76271^3 = 21063928^2$
$1414^3 + 2213459^2 = 65^7$
$9262^3 + 15312283^2 = 113^7$
$43^8 + 96222^3 = 30042907^2$
$33^8 + 1549034^2 = 15613^3$.

# 2 Our problem

Up to now, we assume till the end of the text that the integers $a,b,c$ are fixed. We then consider $x,y,z$ as unknowns only. Precisely, let us fix $a,b,c \in \mathbb{Z}$ with $a,b,c \geq 2, (a,b,c) = 1$ and consider the equation

$$a^x + b^y = c^z \tag{2}$$

in unknowns $x,y,z \in \mathbb{Z}$ with $x,y,z \geq 2$.

In 1993, K. Malher used $p$-adic Thue-Siegel method to show that the solutions $x,y,z$ to (2) are only finitely many. The bound for the number of the solutions should depend on $\omega(abc)$ the number of the primes dividing $abc$. A. O. Gel'fond gave in 1940 a lower bound of linear forms in $p$-adic logarithms and then a bound for the

size of the solutions, namely an effectively calculable constant $C > 0$ depending only on $a, b, c$ such that $\max\{|x|, |y|, |z|\} < C$.

Around 1994, Terai and Jésmanowicz conjectured (*see* for example [Cao-Dong]) that if there exists a solution $(x_0, y_0, z_0)$ then this is the only solution:

**Conjecture 3.** *(Terai and Jésmanowicz) The number of the solutions to the equation (2) is at most* 1.

There are several investigations concerning with Conjecture 3 by N. Terai, Z. Li, or others. They essentially show that there exist particular examples of $a, b, c$ where Conjecture 3 holds. Remark that the identity $2^n + 2^n = 2^{n+1}$ does not give infinitely many solutions. It is also noted that there are trivial identities:

$$2^{n+2} + (2^n - 1)^2 = (2^n + 1)^2 \quad (a = 2 \text{ or } a = 2^{n+1}, b = 2^n - 1, c = 2^n + 1)$$

$$2^1 + 2^n - 1 = 2^n + 1 \quad (a = 2, b = 2^n - 1, c = 2^n + 1).$$

Among the knowns, we quote an example of Conjecture 3 which is made by Terai;

**Example 2.** *(Terai)*
*Suppose that $u$ is even, $a = u^3 - 3u$, $b = 3u^2 - 1$, $b$ is a prime, $c = u^2 + 1$, and that there exsists a prime $l$ such that $l$ divides $u^2 - 3$ with $3|e$ for an integer $e > 0$ satisfying $2^e - 1$ is divisible by $l$. Then the equation (2) has the only solution* $(2, 2, 3)$.

# 3 Our statement

Firstly we state a theorem which is quick to obtain.

**Theorem 1.** *Let $N$ be the number of the solutions to (2). Then we have*

$$N \leq 2^{36}.$$

The advantage of Theorem 1 is the fact that the number $N$ is *independent* of the number $a, b, c$ especially of $\omega(abc)$.
It might be possible to refine the bound in Theorem 1 ; we will prove this by a forthcoming article.

Secondly we show a bound for the size of the solutions:

**Theorem 2.** *Suppose that $c$ is odd and that $c$ has the prime decomposition $c = p_1^{r_1}p_2^{r_2}\cdots p_l^{r_l}$. Suppose that there exists an integer $g \in \mathbb{Z}, g \geq 2$ coprime with $c$ such that*

$$v_{p_i}(a^g - 1) \geq r_i$$

*and*

$$v_{p_i}(b^g - 1) \geq 1$$

*for any prime $p_i|c$. Then we have*

$$\max\{|x|, |y|, |z|\} \leq 2^{288}\sqrt{abc}(\log(abc))^3.$$

# 4 Outline of the proof

Theorem 1 is easily implied by the following theorem due to F. Beukers and H. P. Schlickewei [Be-Schl]. Their result corresponds to a refinement in a low-dimensional case of a theorem by J. -H. Evertse, H. P. Schlickewei and W.M. Schmidt [E-Schl-Schm].

**Theorem 3.** *(Evertse-Schlickewei-Schmidt) Let $n \in \mathbb{Z}, n \geq 1$. Let $K$ be an algebraic closed field with characteristic 0, $\Gamma$ be a finitely generated subgroup of the multiplicative group $(K - \{0\})^n$. Denote by $r < \infty$ the number of the generators of $\Gamma$. Let $a_i \in K - \{0\}$. Consider the equation $a_1X_1 + \cdots + a_nX_n = 1$ in unknowns $X_1, \cdots, X_n$ in $\Gamma$ supposed the subsum satisfying $\Sigma_{i\in I}a_iX_i \neq 0$ for any non-empty proper subset $I$ of $\{1, 2, \cdots, n\}$. Then we have that the number of the solutions $(x_1, \cdots, x_n) \in \Gamma^n$ to the equation $a_1X_1 + \cdots a_nX_n = 1$ is at most*

$$\exp\left((6n)^{3n}(r+1)\right).$$

When $n = 2$, a refinement of the above is as follows:

**Theorem 4.** *(Beukers-Schlickewei) Let $n = 2$. Then we have that the number of the solutions $(x_1, x_2) \in \Gamma^2$ to the equation $a_1X_1 + a_2X_2 = 1$ is at most*

$$2^{9(r+1)}.$$

**Proof of Theorem 1**

It is enough to apply the theorem of Beukers-Schlickewei. Our equation is $a^x + b^y = c^z$, thus

$$\frac{a^x}{c^z} + \frac{b^y}{c^z} = 1.$$

We see that it turns out to consider the equation $X + Y = 1$ with $X, Y$ in "$a, b, c$-units", namely in $\Gamma =< a, b, c >= \{a^k b^l c^m \mid k, l.m \in \mathbb{Z}\}$. Thus just use Beukers-Schlickewei with $r = 3$ to arrive at $2^{36}$.

When $a, b, c$ are distinct primes, then we may use Evertse' bound $3 \cdot 7^{12}$.

If we consider $S = \{p|abc\}$ we do not get independence of $\omega(abc)$ in the statement.

**Proof of Theorem 2**

Let $m$ be an integer $\geq 2$ not necessarily a prime. The concept of linear forms in $m$-adic logarithms is basically introduced by Malher and is revisited by Y. Bugeaud.

Recall the definition of $m$-adic valuation. Let $m = p_1^{r_1} \cdots p_l^{r_l}$ where $p_1 < \cdots < p_l$ are primes, $r_1 \cdots, r_l \in \mathbb{Z}, > 0$. Let $x \in \mathbb{Z}, x \neq 0$. We recall that the $p$-adic valuation is $v_p(x) :=$ the greatest integer $v \geq 0$ such that $p^v | x$. Following this, we define

$$\begin{aligned} v_m(x) &:= \text{the greatest integer } v \geq 0 \text{ such that } m^v | x \\ &= \min_{1 \leq i \leq l} [\frac{v_{p_i}(x)}{r_i}] \end{aligned}$$

where $[\cdot]$ denotes the Gauss' symbol.

For a rational number $\frac{a}{b} \neq 0$, $a, b \in \mathbb{Z}, (a, b) = 1$, we define $v_m(\frac{a}{b}) := v_m(a) - v_m(b)$.

We state a variant of a lemma of Y. Bugeaud by removing some specific conditions. Denote here by $h(\cdot)$ the absolute logarithmic height. Theorem 2 is deduced by using Lemma 1:

**Lemma 1.** *Let* $\Lambda := \alpha_1^{b_1} - \alpha_1^{b_2} \neq 0$ *where* $\alpha_1, \alpha_2 \in \mathbb{Q}, \alpha_1 \neq \pm 1, b_1, b_2 \in \mathbb{Z}, b_1, b_2 > 0$. *Let* $m = p_1^{r_1} \cdots p_l^{r_l}$. *Suppose* $v_{p_i}(\alpha_1) = v_{p_i}(\alpha_2) = 0$ *for any* $p_i | m$. *Suppose further that there exists an integer* $g \in \mathbb{Z}, g > 0$, *coprime with* $m$ *such that*

$$v_{p_i}(\alpha_1^g - 1) \geq r_i,$$

$$v_{p_i}(\alpha_2^g - 1) \geq 1$$

*and moreover*

$$v_2(\alpha_1^g - 1) \geq 2,$$

$$v_2(\alpha_2^g - 1) \geq 2$$

*if* $2 | m$. *Then there exists an effectively computable constant* $C > 0$ *depending on the data with*

$$v_m(\Lambda) \leq \frac{Cm^2}{\max(\log m, 1)^2} \left( \log \left( \frac{|b_1|}{\log A_1} + \frac{|b_2|}{\log A_2} \right) \right)^2 \log A_1 \log A_2$$

*where* $\log A_i \geq \max(h(\alpha_i), \log m) \quad (i = 1, 2)$.

# References


[Be-Schl] F. Beukers & H. P. Schlickewei, *The equation x+y=1 in finitely generated groups*, Acta Arith. 78. 2, 189–199, 1996.

[Cao-Dong] Zhenfu Cao & Xiaolei Dong, *An application of a lower bound for linear forms in two logarithms to the Terai-Jésmanowicz conjecture*, Acta Arith. 110, 153–164, 2003.

[E-Schl] J. -H. Evertse & H. P. Schlickewei, *The Absolute Subspace Theorem and linear equations with unknowns from a multiplicative group*, in*Number Theory in Progress, I*, (eds. K. Győry, H. Iwaniec, J. Urbanowicz), Walter de Gruyter, 121–142, 1999.

[E-Schl-Schm] J. -H. Evertse, H. P. Schlickewei & W.M. Schmidt, *Linear equations in variables which lie in a multiplicative group*, Ann. Math. 155, 1–30, 2002.

[Pa-Schf] A. N. Parshin & I. R. Schfarevich (eds.), N. I. Fel'dman & Yu. V. Nesterenko (authors), *Number Theory IV*, Encyclopaedia of Mathematical Sciences vol. 44, 1998.

[Schm1] W. M. Schmidt, Diophantine approximation, Lecture Notes in Math., 785, Springer, 1980.

[Schm2] W. M. Schmidt, Diophantine approximation and Diophantine Equations, Lecture Notes in Math., 1467, Springer, 1991.

[Sho-T] T. N. Shorey & R. Tijdeman, Exponential Diophantine Equations, Cambridge Tracts in Math., Vol. 87, Cambridge Univ. Press, 1986.

[Wa] M. Waldschmidt, *Diophantine Approximation on Linear Algebraic Groups*, Grundlehren der Math. Wissenschften 326, Springer, 2000.

[Wü] G. Wüstholz, A Panorama of Number Theory, Cambridge Univ. Press, 2002.

Noriko HIRATA-KOHNO
Dept. of Mathematics
College of Science and Technology
NIHON University
Kanda, Chiyoda, Tokyo 101-8308
JAPAN