

On the analyticity in time of solutions of initial boundary value problems for semi-linear parabolic differential equations with monotone nonlinearity

By Sunao Ōuchi

In this note we consider the following initial boundary value problems:

$$\begin{aligned} \frac{\partial u}{\partial t} &= \Delta u + f(u) \\ \text{(I.B.V.P.)} \quad u(t, x)|_{\partial\Omega} &= 0 \\ u(0, x) &= a(x), \end{aligned}$$

where Ω is a bounded domain in R^n with smooth boundary $\partial\Omega$.

The purpose of this note is to report that solutions of (I.B.V.P.) are extensible holomorphically in time t to a sector $\Sigma_\theta = \{t \in \mathbb{C}; |\arg t| < \theta\}$ in the complex domain which does not depend on initial values, if the nonlinear term $f(u)$ is a monotone decreasing polynomial.

Let us now introduce definitions to state results.

Definition 1. A polynomial with real coefficients $f(u)$ is said to be monotone or to satisfy condition (M), if $f(0)=0$ and $f'(u) \leq 0$ for $-\infty < u < +\infty$.

Examples. $f(u) = -u^{2p+1}$, $-u-u^3$, $-u^3-u^5$.

Definition 2. A polynomial with real coefficients $f(u)$ is said to be monotone on $R^+ = [0, \infty)$ or satisfy condition (M_+) , if $f(0) = 0$ and $f'(u) \leq 0$ for $0 \leq u < \infty$.

Examples. $f(u) = -u^{2p}$, $-u-u^4$, $-u^2-u^6$.

Theorem 1. Suppose that the nonlinear term $f(u)$ in (I.B.V.P.) satisfies condition (M) and the initial value $a=a(x)$ is real-valued and boundedly continuous in Ω . Then there is a sector $\Sigma_{\theta_0} = \{t; |\arg t| < \theta_0\}$ in the complex domain which is independent of $a(x)$ such that the solution $u(t, x)$ of (I.B.V.P.) is analytically extensible in t to the sector.

Theorem 2. Suppose that the nonlinear term $f(u)$ in (I.B.V.P.) satisfies condition (M_+) and the initial value $a = a(x)$ is a nonnegative and boundedly continuous function. Then there is a sector $\Sigma_{\theta_0} = \{t; |\arg t| < \theta_0\}$ in the complex domain which does not depend on $a(x)$ such that the nonnegative solution $u(t, x)$ of (I.B.V.P.) is analytically extensible in t to the sector.

Details are published in

S. Ōuchi, On the analyticity in time of solutions of initial boundary value problems for semi-linear parabolic differential equations with monotone nonlinearity, J. Fac. Sci. Univ. Tokyo, Vol. 21, No.1 (1974) 19-41.

Department of Mathematics
Faculty of Science
Tokyo Metropolitan University
Fukazawa, Setagaya, Tokyo
158 Japan