

Harmonic functions in a cylinder with the normal derivatives vanishing on the boundary

千葉大・理 宮本 育子 (Ikuko Miyamoto)
千葉大・自然 吉田 英信 (Hidenobu Yoshida)

1 Introduction

Let $\mathbf{R}^n$ ($n \geq 2$) denote the n -dimensional Euclidean space. When we consider the Neumann problem for an infinite cylinder

$$\Gamma_n(D) = \{(X, y) \in \mathbf{R}^n; X \in D, -\infty < y < \infty\}$$

with a bounded domain D of $\mathbf{R}^{n-1}$, the solution of it is not unique, because we can make another solution from a solution by adding harmonic functions in $\Gamma_n(D)$ with the normal derivatives vanishing on the boundary. Hence, to classify general solutions we need to characterize such functions. If $D = (0, \pi)$ and $\Gamma_n(D)$ is the strip

$$H = \{(x, y) \in \mathbf{R}^2; 0 < x < \pi, -\infty < y < \infty\},$$

then by applying a result of Widder [6, Theorem 2] which characterizes a harmonic function in H vanishing continuously on the boundary ∂H of H , we can obtain the following result

Theorem A. *Let $h(x, y)$ be a harmonic function in H such that $\partial h / \partial x$ vanishes continuously on ∂H . Then*

$$h(x, y) = A_0 y + B_0 + \sum_{k=1}^{\infty} (A_k e^{ky} + B_k e^{-ky}) \cos kx,$$

where the series converges for all x and y , and all $A_0, B_0, A_1, B_1, A_2, B_2, \dots$ are constants such that

$$A_k e^{ky} + B_k e^{-ky} = \frac{2}{\pi} \int_0^{\pi} h(x, y) \cos kx dx \quad (k = 1, 2, 3, \dots).$$

Although this theorem is easily proved by using the fact that $\partial h / \partial x$ is a harmonic function which vanishes continuously on ∂H , we can not proceed similarly in the case where $\Gamma_n(D)$ is a cylinder in $\mathbf{R}^n$ ($n \geq 3$). This kind of problem was originally treated by Bouligand [1] in 1914.

Theorem B (Bouligand [1, p.195]). *Let $h(X, y)$ be a harmonic function in $\Gamma_n(D)$ such that the normal derivative of h vanishes continuously on the boundary $\partial\Gamma_n(D)$ of $\Gamma_n(D)$. If $h(X, y)$ tends to zero as $|y| \rightarrow \infty$, then $h(X, y)$ is identically zero in $\Gamma_n(D)$.*

In this paper we shall prove a cylindrical version of Theorem A (Theorem). As Corollaries we shall obtain two results generalizing Theorem B (Corollaries 1 and 2).

2 Preliminaries

Let D be a bounded domain $\mathbf{R}^{n-1}$ ($n \geq 3$) having sufficiently smooth boundary ∂D . For example, D is a $C^{2,\alpha}$ -domain ($0 < \alpha < 1$) in $\mathbf{R}^{n-1}$ surrounded by a finite number of mutually disjoint closed hypersurfaces (see Gilbarg and Trudinger [3, pp. 88-89] for the definition of $C^{2,\alpha}$ -domain). Consider the Neumann problem

$$(2.1) \quad (\Delta_{n-1} + \mu)\varphi(X) = 0$$

for any $X = (x_1, x_2, \dots, x_{n-1}) \in D$,

$$(2.2) \quad \lim_{X \rightarrow X', X \in D} (\nabla_{n-1}\varphi(X), \nu(X')) = 0$$

for any $X' \in \partial D$, where

$$\Delta_{n-1} = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_{n-1}^2}, \quad \nabla_{n-1} = \left(\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_{n-1}} \right)$$

and $\nu(X')$ is the outer unit normal vector at $X' \in \partial D$.

Let $\{\mu_k(D)\}_{k=0}^\infty$ be the non-decreasing sequence of non-negative eigenvalues of this Neumann problem. In this sequence we write $\mu_k(D)$ the same number of times as the dimension of the corresponding eigenspace. When the normalized eigenfunction corresponding to $\mu_k(D)$ is denoted by $\varphi_k(D)(X)$, the set of sequential eigenfunctions corresponding to the same value of $\mu_k(D)$ in the sequence $\{\varphi_k(D)(X)\}_{k=0}^\infty$ makes an orthonormal basis for the eigenspace of the eigenvalue $\mu_k(D)$. It is evident that $\mu_0(D) = 0$ and

$$\varphi_0(D)(X) = |D|^{-1/2} \quad (X \in D), \quad |D| = \int_D dX.$$

In the following we shall denote $\{\mu_k(D)\}_{k=0}^\infty$ and $\{\varphi_k(D)(X)\}_{k=0}^\infty$ by $\{\mu(k)\}_{k=0}^\infty$ and $\{\varphi_k(X)\}_{k=0}^\infty$ without specifying D , respectively. We can also say that for each D there is a sequence $\{k_i\}$ of non-negative integers such that $k_0 = 0, k_1 = 1, \mu(k_i) < \mu(k_{i+1})$,

$$\mu(k_i) = \mu(k_i + 1) = \mu(k_i + 2) = \dots = \mu(k_{i+1} - 1)$$

and $\{\varphi_{k_i}, \varphi_{k_i+1}, \dots, \varphi_{k_{i+1}-1}\}$ is an orthonormal basis for the eigenspace of the eigenvalue $\mu(k_i)$ ($i = 0, 1, 2, 3, \dots$). Since D has sufficiently smooth boundary, we know that

$$\mu(k) \sim A(D, n)k^{2/(n-1)} \quad (k \rightarrow \infty)$$

and

$$\sum_{\mu(k) \leq t} \{\varphi_k(X)\}^2 \sim B(D, n)t^{(n-1)/2} \quad (t \rightarrow \infty)$$

uniformly with respect to $X \in D$, where $A(D, n)$ and $B(D, n)$ are both constants depending on D and n (e.g. see Carleman [2], Minakshisundaram and Pleijel [4], Weyl [5]). Hence there exist two positive constants M_1, M_2 such that

$$M_1 k^{2/(n-1)} \leq \mu(k) \quad (k = 1, 2, \dots)$$

and

$$|\varphi_k(X)| \leq M_2 k^{1/2} \quad (X \in D, k = 1, 2, \dots).$$

3 Statement of our results

The gradient of a functions $f(P)$ defined on $\Gamma_n(D)$ is

$$\nabla_n f(P) = \left(\frac{\partial f}{\partial x_1}(P), \dots, \frac{\partial f}{\partial x_{n-1}}(P), \frac{\partial f}{\partial y}(P) \right) \quad (P = (x_1, x_2, \dots, x_{n-1}, y) \in \Gamma_n(D)).$$

We first remark that both

$$I_k(P) = e^{\sqrt{\mu(k)}y} \varphi_k(X) \quad \text{and} \quad J_k(P) = e^{-\sqrt{\mu(k)}y} \varphi_k(X) \quad (P = (X, y) \in \Gamma_n(D))$$

are two harmonic functions on $\Gamma_n(D)$ satisfying

$$\lim_{P \rightarrow Q, P \in \Gamma_n(D)} (\nabla_n I_k(P), \nu(Q)) = 0 \quad \text{and} \quad \lim_{P \rightarrow Q, P \in \Gamma_n(D)} (\nabla_n J_k(P), \nu(Q)) = 0,$$

respectively, where $\nu(Q)$ is the outer unit normal vector at $Q \in \partial\Gamma_n(D)$.

Theorem. *Let $h(P)$ be a harmonic function on $\Gamma_n(D)$ satisfying*

$$(3.1) \quad \lim_{P \rightarrow Q, P \in \Gamma_n(D)} (\nabla_n h(P), \nu(Q)) = 0$$

for any $Q \in \partial\Gamma_n(D)$. Then

$$h(P) = A_0 y + B_0 + \sum_{k=1}^{\infty} (A_k I_k(P) + B_k J_k(P))$$

for any $P = (X, y) \in \Gamma_n(D)$, where the series converges uniformly and absolutely on any compact subset of the closure $\overline{\Gamma_n(D)}$ of $\Gamma_n(D)$, and both A_k, B_k ($k = 0, 1, 2, \dots$) are two constants such that

$$(3.2) \quad A_k e^{\sqrt{\mu(k)}y} + B_k e^{-\sqrt{\mu(k)}y} = \int_D h(X, y) \varphi_k(X) dX \quad (k = 1, 2, 3, \dots).$$

Corollary 1. Let p and q be two non-negative integers. If $h(P)$ is a harmonic function on $\Gamma_n(D)$ satisfying (3.1) and

$$(3.3) \quad \lim_{y \rightarrow \infty} e^{-\sqrt{\mu(k_{p+1})}y} M_h(y) = 0 \quad \lim_{y \rightarrow -\infty} e^{\sqrt{\mu(k_{q+1})}y} M_h(y) = 0,$$

where

$$M_h(y) = \sup_{X \in D} |h(X, y)| \quad (-\infty < y < \infty),$$

then

$$h(P) = A_0 y + B_0 + \sum_{k=1}^{k_{p+1}-1} A_k I_k(P) + \sum_{k=1}^{k_{q+1}-1} B_k J_k(P)$$

for any $P = (X, y) \in \Gamma_n(D)$, where A_k ($k = 0, 1, 2, \dots, k_{p+1} - 1$) and B_k ($k = 0, 1, 2, \dots, k_{q+1} - 1$) are constants.

Corollary 2. Let $h(P)$ be a harmonic function on $\Gamma_n(D)$ satisfying (3.1) and

$$M_h(y) = o(e^{\sqrt{\mu(1)}|y|}) \quad (|y| \rightarrow \infty).$$

Then

$$h(P) = A_0 y + B_0$$

for any $P = (X, y) \in \Gamma_n(D)$, where A_0 and B_0 are two constants.

4 Proofs of Theorem and Corollaries 1, 2

Let $f(X, y)$ be a function on $\Gamma_n(D)$. The function $c_k(f, y)$ of y ($-\infty < y < \infty$) defined by

$$c_k(f, y) = \int_D f(X, y) \varphi_k(X) dX$$

is simply denoted by $c_k(y)$ without specifying f , in the following.

Lemma 1. Let $h(P)$ be a harmonic function on $\Gamma_n(D)$ satisfying (3.1), then

$$(4.1) \quad c_0(y) = A_0 y + B_0$$

$$(4.2) \quad c_k(y) = A_k e^{\sqrt{\mu(k)}y} + B_k e^{-\sqrt{\mu(k)}y} \quad (k = 1, 2, 3, \dots)$$

with two constants A_k, B_k ($k \geq 0$) and

$$(4.3) \quad c_k(y) = \frac{\{e^{\sqrt{\mu(k)}(y-y_2)} - e^{\sqrt{\mu(k)}(y_2-y)}\} c_k(y_1) + \{e^{\sqrt{\mu(k)}(y_1-y)} - e^{\sqrt{\mu(k)}(y-y_1)}\} c_k(y_2)}{e^{\sqrt{\mu(k)}(y_1-y_2)} - e^{\sqrt{\mu(k)}(y_2-y_1)}}$$

for any y_1 and y_2 , $-\infty < y_1 < y_2 < \infty$, ($k = 1, 2, 3, \dots$).

Proof. First of all, we remark that $h \in C^2(\overline{\Gamma_n(D)})$ (Gilbarg and Trudinger [3, p.124]). Since

$$\int_D (\Delta_{n-1} h(X, y)) \varphi_k(X) dX = \int_D h(X, y) (\Delta_{n-1} \varphi_k(X)) dX \quad (-\infty < y < \infty),$$

from Green's identity, (2.2) and (3.1), we have

$$\begin{aligned} \frac{\partial^2 c_k(y)}{\partial y^2} &= \int_D \frac{\partial^2 h(X, y)}{\partial y^2} \varphi_k(X) dX = - \int_D \Delta_{n-1} h(X, y) \varphi_k(X) dX. \\ &= - \int_D h(X, y) (\Delta_{n-1} \varphi_k(X)) dX = \mu(k) \int_D h(X, y) \varphi_k(X) dX = \mu(k) c_k(y) \end{aligned}$$

from (2.1) ($k = 0, 1, 2, \dots$). With two constants A_k and B_k ($k = 0, 1, 2, \dots$) these give

$$c_0(y) = A_0 y + B_0$$

and

$$c_k(y) = A_k e^{\sqrt{\mu(k)}y} + B_k e^{-\sqrt{\mu(k)}y} \quad (k = 1, 2, \dots),$$

which are (4.1) and (4.2). When we solve A_k and B_k from

$$c_k(y_i) = A_k e^{\sqrt{\mu(k)}y_i} + B_k e^{-\sqrt{\mu(k)}y_i} \quad (i = 1, 2),$$

we immediately obtain (4.3).

Remark. From (4.2) we have

$$\lim_{y \rightarrow \infty} c_k(y) e^{-\sqrt{\mu(k)}y} = A_k, \quad \text{and} \quad \lim_{y \rightarrow -\infty} c_k(y) e^{\sqrt{\mu(k)}y} = B_k \quad (k = 1, 2, 3, \dots).$$

Lemma 2. Let $h(P)$ be a harmonic function on $\Gamma_n(D)$ satisfying (3.1). Let y be any number and y_1, y_2 be two any numbers satisfying $-\infty < y_1 < y - 1$, $y + 1 < y_2 < \infty$. For two non-negative integers p and q ,

$$\sum_{k=k_{p+q+1}}^{\infty} |c_k(y)| |\varphi_k(X)| \leq L(p) M_h(y_1) + L(q) M_h(y_2),$$

where

$$L(j) = M_2^2 |D| \sum_{k=k_{j+1}}^{\infty} k \exp(-\sqrt{M_1} k^{1/(n-1)}).$$

Proof. From Lemma 1, we see that

$$\begin{aligned} c_k(y) &= \exp\{-\sqrt{\mu(k)}(y - y_1)\} \frac{1 - \exp\{2\sqrt{\mu(k)}(y - y_2)\}}{1 - \exp\{2\sqrt{\mu(k)}(y_1 - y_2)\}} c_k(y_1) \\ &+ \exp\{\sqrt{\mu(k)}(y - y_2)\} \frac{1 - \exp\{2\sqrt{\mu(k)}(y_1 - y)\}}{1 - \exp\{2\sqrt{\mu(k)}(y_1 - y_2)\}} c_k(y_2). \end{aligned}$$

Hence

$$(4.4) \quad \sum_{k=k_{p+q+1}}^{\infty} |c_k(y)| |\varphi_k(X)| \leq I_1 + I_2,$$

where

$$I_1 = \sum_{k=k_{p+1}}^{\infty} \exp\{-\sqrt{\mu(k)}(y - y_1)\} |c_k(y_1)| |\varphi_k(X)|$$

and

$$I_2 = \sum_{k=k_{q+1}}^{\infty} \exp\{-\sqrt{\mu(k)}(y_2 - y)\} |c_k(y_2)| |\varphi_k(X)|.$$

For I_1 , we have

$$\begin{aligned} (4.5) \quad I_1 &\leq M_2^2 |D| M_h(y_1) \sum_{k=k_{p+1}}^{\infty} k \exp(-\sqrt{\mu(k)}) \\ &\leq M_2^2 |D| M_h(y_1) \sum_{k=k_{p+1}}^{\infty} k \exp(-\sqrt{M_1} k^{1/(n-1)}), \end{aligned}$$

because $y - y_1 > 1$.

For I_2 , we also have

$$(4.6) \quad I_2 \leq M_2^2 |D| M_h(y_2) \sum_{k=k_{q+1}}^{\infty} k \exp(-\sqrt{M_1} k^{1/(n-1)}).$$

Finally (4.4), (4.5) and (4.6) give the conclusion of this Lemma.

Proof of Theorem. Take any compact set T , $T \subset \overline{\Gamma_n(D)}$ and two numbers y_1, y_2 satisfying

$$\max\{y; (X, y) \in T\} + 1 < y_2, \quad \min\{y; (X, y) \in T\} - 1 > y_1.$$

Let (X, y) be any point in T . Since $c_k(y)$ is the Fourier coefficient of the function $h(X, y)$ of X with respect to the orthonormal sequence $\{\varphi_k(X)\}_{k=0}^{\infty}$, we have

$$h(X, y) = \sum_{k=0}^{\infty} c_k(y) \varphi_k(X)$$

in which the series converges uniformly and absolutely on T by Lemma 2. Further (4.1) and (4.2) in Lemma 1 give (3.2). The proof of Theorem is complete.

Proofs of Corollaries 1 and 2. From (3.3) and Remark, it follows that $A_k = 0$ for any k , $k \geq k_{p+1}$ and $B_k = 0$ for any k , $k \geq k_{q+1}$. Hence Theorem immediately gives the conclusion of Corollary 1. By putting $p = q = 0$ in Corollary 1, we obtain Corollary 2 at once.

参考文献

- [1] M.G.Bouligand, *Sur les fonctions de Green et de Neumann du cylindre*, Bull. Soc. Math. France 42 (1914), 168-242.
- [2] T.Carleman, *Propriétés asymptotiques des fonctions fondamentales des membranes vibrantes*, C. R. Skand. Math. Kongress 1934, 34-44.
- [3] D.Gilbarg and N.S.Trudinger, *Elliptic partial differential equations of second order*, Springer, 1977.
- [4] S.Minakshisundaram and A.Pleijel, *Some properties of the eigenfunctions of the Laplace-operator on Riemannian manifolds*, Canad. J. Math. 1 (1949), 242-256.
- [5] H.Weyl, *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen*, Math. Ann. 71 (1911), 441-471.
- [6] D.V.Widder, *Functions harmonic in a strip*, Proc. Amer. Math. Soc. 12 (1961), 67-72.