

1 **Title page**

2 **Enhanced echo intensity and a higher extracellular water-to-intracellular water ratio are helpful**
3 **clinical signs for detecting muscle degeneration in patients with knee osteoarthritis**

4

5 **The names of the authors:**

6 Masashi Taniguchi (ORCID: 0000-0003-0548-0210)¹, Yoshihiro Fukumoto (ORCID: 0000-0003-0516-
7 770X)^{1,2}, Masahide Yagi (ORCID: 0000-0002-1825-3570)¹, Momoko Yamagata (ORCID: 0000-0001-
8 8192-5651)^{1,3}, Masashi Kobayashi ⁴, Yosuke Yamada (ORCID: 0000-0002-4284-6317)⁵, Misaka
9 Kimura (ORCID: 0000-0001-8110-8299)⁶, Noriaki Ichihashi (ORCID: 0000-0003-2508-2172)¹

10

11 **Affiliations:**

12 ¹ Human Health Sciences, Graduate School of Medicine, Kyoto University

13 ² Department of Physical Medicine and Rehabilitation, Kansai Medical University

14 ³ Graduate School of Human Development and Environment, Kobe university

15 ⁴ Department of Orthopaedic Surgery, Kobayashi Hospital

16 ⁵ National Institute of Health and Nutrition, National Institutes of Biomedical Innovation, Health and
17 Nutrition

18 ⁶ Graduate School of Nursing, Doshisha Women's College of Liberal Arts

19

20 Address correspondence to Masashi Taniguchi, RPT, PhD,

21 Human Health Sciences, Graduate School of Medicine, Kyoto University

22 53-Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan.

23 Email: taniguchi.masashi.7a@kyoto-u.ac.jp

24 Tel.: +81-75-751-3964 Fax: +81-: +81-75-751-3964

25 ORCID Number: 0000-0003-0548-0210

26

27

28 **Abstract**

29 **Introduction/objectives:** Enhanced muscle echo intensity (EI) with ultrasound imaging and a higher

30 extracellular water-to-intracellular water (ECW/ICW) ratio with segmental-bioelectrical impedance

31 spectroscopy (S-BIS) represent muscle quality loss. This study aimed to clarify quadriceps muscle

32 degeneration characteristics, focusing on muscle quality changes in patients with knee osteoarthritis (OA).

33 **Method:** Forty-one women with knee OA (mean age, 71.4 ± 6.0 years) and 27 healthy women (mean age,

34 75.6 ± 4.9 years) participated. Ultrasonography was used to evaluate the muscle thickness (MT) and the

35 EI of each quadriceps compartment. The ECW/ICW ratio was obtained by S-BIS. MT, EI, and ECW/ICW

36 ratio differences between the two groups were tested using univariate analysis of variance, adjusting for

age and body mass index. Logistic regression analyses were performed with the group as the dependent variable, and the MT and EI of the vastus medialis (VM) and the ECW/ICW ratio as independent variables.

Results: Patients with knee OA had a significant decrease in VM MT, enhanced VM, and vastus intermedius EIs and a higher ECW/ICW ratio compared with healthy participants. Logistic regression analysis showed that the VM EI (odds ratio [OR], 1.19; 95% confidence interval [CI], 1.06–1.35) and the ECW/ICW ratio were independently associated with knee OA (OR, 1.19; 95% CI, 1.00–1.42).

Conclusions: VM EI and the ECW/ICW ratio, rather than VM MT, characterised quadriceps muscle degeneration in patients with knee OA. Therefore, enhanced EI and a higher ECW/ICW ratio are helpful clinical signs for detecting muscle degeneration in patients with knee OA.

Key Points

- Echo intensity (EI) of the vastus medialis and the extracellular-to-intracellular water (ECW/ICW) ratio significantly increased in patients with knee osteoarthritis (OA).
- Enhanced EI and a higher ECW/ICW ratio are useful clinical signs for detecting muscle degeneration in patients with knee OA.

Keywords: knee osteoarthritis, quadriceps muscle degeneration, echo intensity, extracellular-to-intracellular water ratio, muscle quality.

55 **Introduction**

56 Quadriceps muscle degeneration, which includes loss of muscle mass and quality, is a known risk factor
57 for knee osteoarthritis (OA) development and the decline of functional abilities [1-3]. A recent meta-
58 analysis [4] indicated that muscle fat infiltration (i.e., poor muscle quality) was higher in patients with
59 knee OA than in healthy participants. A more recent study [5] also showed that poor muscle quality, but
60 not muscle mass loss, is associated with functional disabilities. Therefore, muscle quality assessment
61 may be fundamental for elucidating the association between muscle dysfunction and functional
62 disabilities in knee OA.

63 Evaluating morphological and composition changes of the quadriceps femoris is important
64 for estimating knee OA pathogenesis [6,7]. However, the use of imaging modalities, such as magnetic
65 resonance imaging and dual-energy X-ray absorptiometry, are often limited in research fields. By
66 contrast, muscle thickness (MT) and muscle echo intensity (EI) can be non-invasively and easily
67 measured via ultrasound images in a clinical setting and are widely used for muscle quantity and quality
68 indices [8-10]. Previous studies have reported that an EI increase occurred at an earlier age and OA
69 stage than did an MT decrease [9,11,12]. Specifically, the vastus medialis (VM) muscle EI in patients
70 with knee OA significantly increased compared with healthy participants [12], suggesting that muscle
71 quality assessment using the VM EI was useful for detecting muscle degeneration in patients with knee
72 OA.

73 The relative increase of the extracellular water (ECW) compartment within the skeletal
74 muscle is also an important determinant of muscle quality [13]. Segmental-bioelectrical impedance
75 spectroscopy (S-BIS) is a convenient, affordable, and non-invasive tool for evaluating muscle quality.
76 Skeletal muscle tissue contains a large amount of water, and S-BIS can distinguish intracellular water
77 (ICW) and ECW from total body water [14,15]. Generally, ICW represents muscle cell mass, and ECW
78 represents non-contractile tissue, including interstitial fluid in the extracellular space [16]. A higher
79 ECW/ICW ratio indicates a relative increase of non-contractile tissue to muscle mass (i.e., muscle
80 quality loss) [17]. Although a high ECW/ICW ratio in patients with knee OA was associated with
81 physical dysfunction [5], it remains unclear whether the ECW/ICW ratio in patients with knee OA is
82 higher than that in healthy participants.

83 Furthermore, a previous study that simultaneously investigated EI and the ECW/ICW ratio
84 suggested that these indicators represent different qualitative changes within the skeletal muscle [18].
85 Ultrasound images cannot distinguish a liquid component from muscle composition. Therefore, EI
86 assessment is assumed to reflect mainly adipose and fibrous tissue. Conversely, the ECW/ICW ratio
87 indicates an expansion of the extracellular space based on increased water content [18]. To our
88 knowledge, no studies have simultaneously investigated EI and the ECW/ICW ratio in patients with
89 knee OA. Specifically, it is unknown if an enhanced EI and a higher ECW/ICW ratio can be used to
90 characterise knee OA muscle degeneration, which would indicate a loss of muscle quality.

91 This study aimed to clarify the characteristics of quadriceps muscle degeneration by focusing
92 on muscle quality changes in patients with knee OA. We hypothesized that (1) the VM EI and the
93 ECW/ICW ratio in patients with knee OA are increased compared to that of healthy participants; and
94 (2) these indicators, which suggest a loss of muscle quality rather than muscle quantity, could be used
95 to characterise quadriceps muscle degeneration in patients with knee OA.

96

97

98 **Materials and Methods**

99 *Study participants and selection*

100 Forty-one female patients with symptomatic medial knee OA were recruited from one community
101 orthopaedic clinic. Twenty-seven healthy control women were also recruited from local communities in
102 Kyoto and neighbouring cities. All study procedures were approved by the Ethics Committee of the
103 Kyoto University Graduate School of Medicine and were conducted following the Declaration of
104 Helsinki principles. All participants were informed regarding the purpose and procedures of this study
105 and gave written informed consent.

106 All patients with knee OA were diagnosed based on the American College of Rheumatology criteria for
107 osteoarthritis of the knee [19] and classified using the Kellgren-Lawrence (KL) grading system ≥ 2 for
108 unilateral or bilateral knees [20]. The inclusion criteria for knee OA were symptomatic and radiographic

109 knee OA, the ability to live independently, and the ability to walk with or without assistive devices. The
110 exclusion criteria for all patients were a surgical history for the back or both limbs, rheumatoid arthritis,
111 and cardiovascular or neurological disorders. In the case of bilateral knee OA, the more severe side was
112 selected for data analysis. If the participant had equal OA severity for both knees, the more painful side
113 was selected. Of 41 patients with knee OA, the KL grades were as follows: grade 2, n = 19; grade 3, n
114 = 13; grade 4, n = 9.

115 The inclusion criteria in the healthy control group were no self-reported history of knee pain,
116 the ability to live independently, and the ability to walk with or without assistive devices. The exclusion
117 criteria for healthy control participants were the same as that used for the knee OA group, and the right
118 knee was selected for data analysis. Based on a previous study that reported the VM EI between knee
119 OA and healthy control groups [12], an effect size of 0.98 was applied for the sample size calculation
120 with an α level of 0.05 and a β level of 0.20. The sample size was calculated using G*Power software
121 (version 3.1.9.7, Universität Kiel, Germany); 18 participants were needed in each group.

122

123 *Self-reported knee function and symptoms*

124 The Knee Society's Knee Scoring System (KSS) 2011 was used to evaluate knee function and
125 symptoms. The KSS 2011 is a self-administered measurement tool for knee conditions, and the validity
126 of KSS has been confirmed in the Japanese population [21]. There are four KSS subcategories:

127 symptoms, satisfaction, expectations, and functional activities. This study focused on the functional
128 activities and symptoms categories. The functional activities category was chosen to evaluate the degree
129 of physical dysfunction during daily activities. This category has four components: walking and
130 standing, standard activities, advanced activities, and discretionary activities. The symptom category
131 has three components: the degree of knee pain during walking, the degree of knee pain when travelling
132 up or down stairs, and knee stiffness. The maximum possible functional activities and symptom scores
133 are 100 and 25 points, respectively, and lower scores represent worse functional activity abilities and
134 symptoms.

135

136 *Muscle thickness and echo intensity measurements using ultrasound images*

137 Transverse ultrasound images were acquired using a B-mode ultrasound imaging device with an 8 MHz
138 linear-array probe (LOGIQ e; GE Healthcare UK Ltd., Chalfont, UK). The participants rested for more
139 than 3 min in the relaxed supine position, and then ultrasound image measurements were performed.
140 All measurements were taken by the same investigator using the same equipment settings (58-dB gain
141 and 69-dB dynamic range). The dynamic focus depth was set to the middle of the muscle of interest.
142 The measurement sites of each muscle followed a previous study [12], and the MTs of the rectus femoris
143 (RF), vastus intermedius (VI), vastus lateralis (VL), and VM were measured. The MT was measured on
144 the ultrasound images using an electronic calliper. EI was measured by converting the image pixels to

an 8-bit grey-scale using image processing software (ImageJ-WinJP; LISIT, Japan). The average EI in the region of interest was calculated on a 256-point scale from 0 to 255, with the high EI values indicating more augmented fat and connective tissue within the muscle [22]. All EI analyses were conducted by another investigator who did not know the group attributes. The reliability of the MT and EI measurements were confirmed by estimating the intra-class correlation coefficients (ICC) for the between-day test-retest reliability. The ICC values of MT and EI ranged from 0.94 to 0.85 and 0.88 to 0.81, respectively. Ultrasound measurements were performed as previously described [18].

Quantification of the ECW/ICW ratio in the upper thigh

The S-BIS measurement procedure was conducted following the ultrasound measurements to avoid body water re-distribution. The ECW/ICW ratio was assessed with a multi-frequency S-BIS device (Fig. 1). The S-BIS device (SFB7, ImpediMed Inc., Australia) measures bioelectrical impedances using a logarithmic distribution of 256 frequencies (ranging from 4 kHz to 1000 kHz) with disposable tab-type electrodes (Red Dot TM; 3M Inc., Japan). The S-BIS measurements were performed for three consecutive repetitions, and the average bioelectrical impedance value was used for analysis. The acquisition, storage, and data processing were conducted using SFB7 Bioimp software (ImpediMed Inc., Australia). The resistance of zero (R_0) and infinity (R_∞) were estimated and obtained by fitting the spectrum of impedance data to the Cole-Cole model. The R_0 and R_∞ indicate the ECW compartment

163 (i.e., $R_0 = R_{ECW}$) and the total body water compartment (TBW; i.e., $R_\infty = R_{TBW}$). The resistance of the
 164 ICW compartment (R_{ICW}) was calculated as $1/[(1/R_{TBW}) - (1/R_{ECW})]$. The ECW and ICW in the thigh
 165 were calculated by applying the estimation algorithm as follows: $ECW = \rho_{ECW} \times L^2/R_{ECW}$ and $ICW =$
 166 $\rho_{ICW} \times L^2/R_{ICW}$, where ρ represents the specific resistivity ($\rho_{ECW} = 47 \Omega\text{cm}$) and intracellular resistivity
 167 ($\rho_{ICW} = 273.9 \Omega\text{m}$). L indicates the segment length (cm), which corresponds to the distance between the
 168 anterior superior iliac spine and the proximal end of the patella. Then, the ECW/ICW ratio was
 169 converted to ECW against ICW. The ICC value for the test-retest reliability in between-day was 0.99 in
 170 the ECW/ICW ratio. The detailed description of S-BIS measurements is indicated elsewhere [18].

171

172 *Statistical analysis*

173 All values are presented as means and standard deviations (SDs). After normal distributions were
 174 confirmed with the Kolmogorov-Smirnov test, unpaired t-tests were applied to the participant
 175 characteristics and KSS scores. Univariate analysis of variance (ANOVA) was performed to investigate
 176 the differences in MT, EI, and the ECW/ICW ratio between the groups, and the adjusted mean difference
 177 between the groups was also estimated, with adjustments for age and BMI. Furthermore, we conducted
 178 logistic regression analyses to characterise quadriceps muscle degeneration in patients with knee OA.
 179 Logistic regression analysis was performed with the knee OA group (reference, healthy control group)
 180 as the dependent variable and MT, VM EI, and the ECW/ICW ratio as independent variables, after

181 adjusting for age and BMI. To identify a cut-off for muscle degeneration characterisation in knee OA,
182 receiver operating characteristic (ROC) curve analysis using the Youden index was performed on the
183 significant variables selected by logistic regression analysis. ROC curve analysis also calculated the
184 area under the curve (AUC), sensitivity, and specificity.

185 All statistical tests were conducted with SPSS software version 25.0 (SPSS Japan Inc., Tokyo,
186 Japan). Statistical significance was set at $p < 0.05$.

187

188

189 **Results**

190 Table 1 presents the characteristics of the patients with knee OA and healthy control participant. The
191 knee OA group was significantly younger than the healthy control group, but body mass and BMI in the
192 knee OA group were significantly higher. The KSS function and symptom scores in the knee OA group
193 were significantly lower than those in the healthy control group.

194 Univariate ANOVAs with adjustment for age and BMI showed significant group differences
195 in the VM MT, EIs of VI and VM, and the ECW/ICW ratio (Table 2). The VM MT was significantly
196 smaller in the knee OA group than in the healthy control group (adjusted mean difference, -0.44 cm;
197 95 % confidence interval [CI], -0.65 to -0.23). Additionally, the EIs of VI and VM in the knee OA group
198 were significantly higher than those in the healthy control group, and the adjusted mean difference in

199 the VM EI was remarkably higher in the knee OA group compared to the healthy control group (22.24
200 arbitrary units [a.u.]; 95% CI, 15.19 to 29.29). Figure 2 shows representative ultrasound images of the
201 VM in healthy controls and patients with knee OA. Moreover, the ECW/ICW ratio in the knee OA
202 group significantly increased compared with the healthy control group (0.10 a.u.; 95% CI, 0.05 to 0.15).

203 Logistic regression analysis showed that the VM EI (odds ratio [OR], 1.19; 95% CI, 1.06 to
204 1.35) and the ECW/ICW ratio were independently associated with knee OA (OR, 1.19; 95% CI, 1.00 to
205 1.42), but the VM MT was not (Table 3). ROC curve analyses determined that the optimal cut-off points
206 for characterising muscle degeneration in knee OA were 80.4 a.u. for VM EI and 0.45 a.u. for the
207 ECW/ICW ratio (Fig. 3). The ROC model on VM EI had high accuracy, with an AUC of 0.90, sensitivity
208 of 0.81, and specificity of 0.89.

209

210

211 **Discussion**

212 This was the first study to clarify the characteristics of quadriceps muscle degeneration in patients with
213 knee OA, focusing on muscle quality changes. In agreement with a previous report [12], the VM MT in
214 patients with knee OA significantly decreased, and EIs of the VI and VM in patients with knee OA
215 increased compared to healthy participants. The novel finding in this study was that the ECW/ICW ratio
216 in patients with knee OA was higher than that in healthy participants, consistent with our hypotheses.

217 Moreover, the logistic regression analysis indicated that both enhanced EI of the VM and a higher
218 ECW/ICW ratio could distinguish between OA and healthy knees, also supporting our hypotheses.
219 These findings suggest that quadriceps muscle degeneration in patients with knee OA is characterised
220 by muscle quality loss rather than muscle quantity.

221 Previous studies investigating muscle size and fat content using magnetic resonance imaging
222 indicated that muscle atrophy and fat infiltration of VM in patients with knee OA were associated with
223 cartilage loss and worsening symptoms [1,23]. This study measured MT and EI using ultrasound and
224 found an MT decrease and an EI increase in the VM of patients with knee OA. Since the VM in patients
225 with knee OA exhibits selective atrophy of type 2 fibres and fatty degeneration [24], these morphologic
226 and composition changes can be indirectly assessed by measuring MT and EI using ultrasound.
227 Although this was known, we found that the ECW/ICW ratio measured using S-BIS was significantly
228 higher in patients with knee OA than in healthy participants, which is new. Increases in the EI and the
229 ECW/ICW ratio indicate lower muscle quality, reflecting the relative increase of non-contractile tissue
230 to muscle mass. Interestingly, Noehren et al. [25] confirmed the expansion of the extracellular matrix
231 in muscle cells by biopsy of patients with knee OA with lower quadriceps function. The ECW
232 compartment on bioelectrical impedance theoretically reflects interstitial fluid in the extracellular space.
233 Therefore, the ECW/ICW ratio could be associated with the expansion of the extracellular matrix,
234 suggesting that assessing the ECW/ICW ratio can effectively evaluate quadriceps muscle degeneration

235 in patients with knee OA. Although the mechanism for an increasing ECW/ICW ratio is unknown, some
236 reports suggest that high inflammation within the muscle in knee OA may be involved in muscle
237 degeneration [26-28].

238 Notably, our finding indicated that quadriceps muscle degeneration in patients with knee OA
239 was characterised by an EI and ECW/ICW ratio increase rather than an MT decrease. Kumar et al. [7]
240 also suggested that the quadriceps intramuscular fat fraction, rather than muscle size, was associated
241 with the structural and symptomatic severity of knee OA. These suggestions support the need to focus
242 on both the accumulation of adipose and fibrous tissues and the expansion of extracellular water
243 contents within the skeletal muscle, rather than only muscle size, to assess muscle degeneration in
244 patients with knee OA. Therefore, simultaneously assessing the EI and the ECW/ICW ratio was helpful
245 for a more accurate characterisation of muscle quality loss in knee OA. Additionally, the ROC curve
246 analysis results showed that the discrimination accuracy for characterising muscle degeneration in knee
247 OA was high in the VM EI. A possible reason for this difference is that the EI measured using ultrasound
248 can assess individual muscles, such as VM, whereas the ECW/ICW ratio measured using S-BIS cannot
249 divide the quadriceps muscle group into four individual muscles. Given the poor muscle quality in the
250 VM of patients with knee OA, the ECW/ICW ratio (which cannot evaluate individual quadriceps
251 muscles) may underestimate specific muscle degeneration. The ECW/ICW ratio was a useful biomarker
252 for poor functional disabilities [5]. However, our findings suggested that the EI assessment of VM was

253 more sensitive for detecting muscle degeneration in patients with knee OA.

254 As a clinical suggestion, assessments of muscle quality using ultrasound images and/or S-
255 BIS are recommended in primary care to better understand the exact function of the quadriceps
256 regardless of knee pain. Although EI assessment using ultrasound is useful for detecting a decline of
257 individual muscle quality, ECW/ICW ratio assessment using bioelectrical impedance equipment can
258 also be substituted. Previous studies suggested that these muscle quality indicators changed during 8–
259 12 weeks of training intervention and detraining [29-31], and thus regular monitoring of 8–12 weeks on
260 muscle degeneration is required for physicians. In the future, given that decline of muscle quality
261 occurred at an earlier age and OA stage than did loss of muscle mass [9,11,12], muscle quality indicators
262 could be a predictor of knee OA development.

263 The present study had several limitations. First, the cross-sectional design of our study could
264 not determine a causal relationship between quadriceps muscle degeneration and the presence of knee
265 OA. Since a previous study already reported that higher intramuscular fat content is associated with
266 knee OA progression [23], future studies should evaluate whether muscle quality loss (i.e., enhanced EI
267 and a higher ECW/ICW ratio) results in knee OA development and progression. Second, the participants
268 of both groups were only older women, and there were also significant differences in age and body mass
269 between the two groups. Although the prevalence of knee OA is higher in older and obese women
270 [32,33], our findings might not be generalizable to older men because there is also a difference between

271 the sexes on quadriceps muscle function in patients with knee OA.

272 In conclusion, patients with knee OA had significantly decreased MT, an enhanced VM EI,
273 and an increased ECW/ICW ratio compared with healthy participants. Furthermore, an increase in the
274 VM EI and the ECW/ICW ratio characterised quadriceps muscle degeneration in patients with knee OA.
275 Therefore, enhanced EI and a higher ECW/ICW ratio are helpful clinical signs for detecting muscle
276 degeneration in patients with knee OA.

277

278

279 **Declarations**

280 ***Funding***

281 This study was supported by the JSPS KAKENHI Grant-in-Aid for Scientific Research (18H03164 and
282 20K19376). The grant was used for data collection, data analysis, and writing the manuscript in this
283 study.

284

285 ***Conflicts of Interest***

286 On behalf of all authors, the corresponding author states that there is no conflict of interest.

287

288 ***Availability of data and material***

289 Data are not available due to ethical restrictions.

290

291 ***Code availability***

292 Not applicable

293

294 ***Authors' contributions***

295 All authors have made substantial contributions to (1) the conception and design of the study; (2)

296 revising it critically for important intellectual content; and (3) final approval of the version to be

297 submitted. The specific contributions of each author are as follows:

298 (1) Analysis and interpretation of the data: MT, YF, MY, YY, and NI.

299 (2) Drafting of the article: MT, YF, MY, and NI.

300

301 ***Ethics approval***

302 All study procedures were approved by the Ethics Committee of the Kyoto University Graduate School

303 of Medicine and were conducted in accordance with the principles of the Declaration of Helsinki.

304

305 ***Consent to participate and consent for publication***

306 Written informed consent was obtained from all participants.

307

308 **Acknowledgements**

309 We would like to thank Editage (www.editage.jp) for English language editing.

310

311

312 **References**

313 1. Wang Y, Wluka AE, Berry PA, Siew T, Teichtahl AJ, Urquhart DM, Lloyd DG, Jones G, Cicuttini

314 FM (2012) Increase in vastus medialis cross-sectional area is associated with reduced pain, cartilage

315 loss, and joint replacement risk in knee osteoarthritis. *Arthritis Rheum* 64 (12):3917-3925.

316 <https://doi.org/10.1002/art.34681>

317 2. Maly MR, Calder KM, Macintyre NJ, Beattie KA (2013) Relationship of intermuscular fat volume

318 in the thigh with knee extensor strength and physical performance in women at risk of or with knee

319 osteoarthritis. *Arthritis Care Res (Hoboken)* 65 (1):44-52. <https://doi.org/10.1002/acr.21868>

320 3. Ruhdorfer A, Wirth W, Eckstein F (2015) Relationship between isometric thigh muscle strength and

321 minimum clinically important differences in knee function in osteoarthritis: data from the

322 osteoarthritis initiative. *Arthritis Care Res (Hoboken)* 67 (4):509-518.

323 <https://doi.org/10.1002/acr.22488>

324 4. Pedroso MG, de Almeida AC, Aily JB, de Noronha M, Mattiello SM (2019) Fatty infiltration in the

thigh muscles in knee osteoarthritis: a systematic review and meta-analysis. *Rheumatol Int* 39 (4):627-635. <https://doi.org/10.1007/s00296-019-04271-2>

5. Taniguchi M, Ikezoe T, Kamitani T, Tsuboyama T, Ito H, Matsuda S, Tabara Y, Matsuda F, Ichihashi N, Nagahama Study Group (2021) Extracellular-to-intracellular water ratios are associated with functional disability levels in patients with knee osteoarthritis: results from the Nagahama Study. *Clin Rheumatol*. Online ahead of print. <https://doi.org/10.1007/s10067-021-05591-0>

6. Scott D, Blizzard L, Fell J, Jones G (2012) Prospective study of self-reported pain, radiographic osteoarthritis, sarcopenia progression, and falls risk in community-dwelling older adults. *Arthritis Care Res (Hoboken)* 64 (1):30-37. <https://doi.org/10.1002/acr.20545>

7. Kumar D, Karampinos DC, Macleod TD, Lin W, Nardo L, Li X, Link TM, Majumdar S, Souza RB (2014) Quadriceps intramuscular fat fraction rather than muscle size is associated with knee osteoarthritis. *Osteoarthritis Cartilage* 22 (2):226-234. <https://doi.org/10.1016/j.joca.2013.12.005>

8. Fukumoto Y, Ikezoe T, Yamada Y, Tsukagoshi R, Nakamura M, Mori N, Kimura M, Ichihashi N (2012) Skeletal muscle quality assessed from echo intensity is associated with muscle strength of middle-aged and elderly persons. *Eur J Appl Physiol* 112 (4):1519-1525. <https://doi.org/10.1007/s00421-011-2099-5>

9. Fukumoto Y, Ikezoe T, Tateuchi H, Tsukagoshi R, Akiyama H, So K, Kuroda Y, Yoneyama T, Ichihashi N (2012) Muscle mass and composition of the hip, thigh and abdominal muscles in women

- 343 with and without hip osteoarthritis. *Ultrasound Med Biol* 38 (9):1540-1545.
 344 <https://doi.org/10.1016/j.ultrasmedbio.2012.04.016>
- 345 10. Chopp-Hurley JN, Wiebenga EG, Bulbrook BD, Keir PJ, Maly MR (2020) Evaluating the
 346 relationship between quadriceps muscle quality captured using ultrasound with clinical severity in
 347 women with knee osteoarthritis. *Clin Biomech (Bristol, Avon)* 80:105165.
 348 <https://doi.org/10.1016/j.clinbiomech.2020.105165>
- 349 11. Fukumoto Y, Ikezoe T, Yamada Y, Tsukagoshi R, Nakamura M, Takagi Y, Kimura M, Ichihashi N
 350 (2015) Age-related ultrasound changes in muscle quantity and quality in women. *Ultrasound Med*
 351 *Biol* 41 (11):3013-3017. <https://doi.org/10.1016/j.ultrasmedbio.2015.06.017>
- 352 12. Taniguchi M, Fukumoto Y, Kobayashi M, Kawasaki T, Maegawa S, Ibuki S, Ichihashi N (2015)
 353 Quantity and quality of the lower extremity muscles in women with knee osteoarthritis. *Ultrasound*
 354 *Med Biol* 41 (10):2567-2574. <https://doi.org/10.1016/j.ultrasmedbio.2015.05.014>
- 355 13. Yamada Y, Yoshida T, Yokoyama K, Watanabe Y, Miyake M, Yamagata E, Yamada M, Kimura M,
 356 Kyoto-Kameoka S (2016) The extracellular to intracellular water ratio in upper legs is negatively
 357 associated with skeletal muscle strength and gait speed in older people. *J Gerontol A Biol Sci Med*
 358 *Sci*:293-298. <https://doi.org/10.1093/gerona/glw125>
- 359 14. Segal KR, Burastero S, Chun A, Coronel P, Pierson RN, Jr., Wang J (1991) Estimation of
 360 extracellular and total body water by multiple-frequency bioelectrical-impedance measurement. *The*

Am J Clin Nutr 54 (1):26-29. <https://doi.org/10.1093/ajcn/54.1.26>

15. Gudivaka R, Schoeller DA, Kushner RF, Bolt MJ (1999) Single- and multifrequency models for bioelectrical impedance analysis of body water compartments. *Journal of applied physiology* 87 (3):1087-1096. <https://doi.org/10.1152/jappl.1999.87.3.1087>

16. Mingrone G, Bertuzzi A, Capristo E, Greco AV, Manco M, Pietrobelli A, Salinari S, Heymsfield SB (2001) Unreliable use of standard muscle hydration value in obesity. *Am J Physiol Endocrinol Metab* 280 (2):E365-371. <https://doi.org/10.1152/ajpendo.2001.280.2.E365>

17. Yamada Y, Buehring B, Krueger D, Anderson RM, Schoeller DA, Binkley N (2017) Electrical properties assessed by bioelectrical impedance spectroscopy as biomarkers of age-related loss of skeletal muscle quantity and quality. *J Gerontol A Biol Sci Med Sci* 72 (9):1180-1186. <https://doi.org/10.1093/gerona/glw225>

18. Taniguchi M, Yamada Y, Fukumoto Y, Sawano S, Minami S, Ikezoe T, Watanabe Y, Kimura M, Ichihashi N (2017) Increase in echo intensity and extracellular-to-intracellular water ratio is independently associated with muscle weakness in elderly women. *Eur J Appl Physiol* 117 (10):2001-2007. <https://doi.org/10.1007/s00421-017-3686-x>

19. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, Christy W, Cooke TD, Greenwald R, Hochberg M, et al. (1986) Development of criteria for the classification and reporting of osteoarthritis. Classification of osteoarthritis of the knee. *Diagnostic and Therapeutic Criteria*

Committee of the American Rheumatism Association. *Arthritis Rheum* 29 (8):1039-1049.
<https://doi.org/10.1002/art.1780290816>

20. Kellgren JH, Lawrence JS (1957) Radiological assessment of osteo-arthritis. *Ann Rheum Dis* 16 (4):494-502. <https://doi.org/10.1136/ard.16.4.494>

21. Taniguchi N, Matsuda S, Kawaguchi T, Tabara Y, Ikezoe T, Tsuboyama T, Ichihashi N, Nakayama T, Matsuda F, Ito H (2015) The KSS 2011 Reflects Symptoms, Physical Activities, and Radiographic Grades in a Japanese Population. *Clin Orthop Relat Res* 473 (1):70-75.
<https://doi.org/10.1007/s11999-014-3650-6>

22. Pillen S, Tak RO, Zwarts MJ, Lammens MM, Verrijp KN, Arts IM, van der Laak JA, Hoogerbrugge PM, van Engelen BG, Verrips A (2009) Skeletal muscle ultrasound: correlation between fibrous tissue and echo intensity. *Ultrasound Med Biol* 35 (3):443-446.
<https://doi.org/10.1016/j.ultrasmedbio.2008.09.016>

23. Raynauld JP, Pelletier JP, Roubille C, Dorais M, Abram F, Li W, Wang Y, Fairley J, Cicuttini FM, Martel-Pelletier J (2015) Magnetic resonance imaging-assessed vastus medialis muscle fat content and risk for knee osteoarthritis progression: relevance from a clinical trial. *Arthritis Care Res (Hoboken)* 67 (10):1406-1415. <https://doi.org/10.1002/acr.22590>

24. Fink B, Egl M, Singer J, Fuerst M, Bubenheim M, Neuen-Jacob E (2007) Morphologic changes in the vastus medialis muscle in patients with osteoarthritis of the knee. *Arthritis Rheum* 56 (11):3626-

3633. <https://doi.org/10.1002/art.22960>

25. Noehren B, Kosmac K, Walton RG, Murach KA, Lyles MF, Loeser RF, Peterson CA, Messier SP (2018) Alterations in quadriceps muscle cellular and molecular properties in adults with moderate knee osteoarthritis. *Osteoarthritis Cartilage* 26 (10):1359-1368. <https://doi.org/10.1016/j.joca.2018.05.011>

26. Levinger I, Levinger P, Trenerry MK, Feller JA, Bartlett JR, Bergman N, McKenna MJ, Cameron-Smith D (2011) Increased inflammatory cytokine expression in the vastus lateralis of patients with knee osteoarthritis. *Arthritis Rheum* 63 (5):1343-1348. <https://doi.org/10.1002/art.30287>

27. Ikemoto-Uezumi M, Matsui Y, Hasegawa M, Fujita R, Kanayama Y, Uezumi A, Watanabe T, Harada A, Poole AR, Hashimoto N (2017) Disuse atrophy accompanied by intramuscular ectopic adipogenesis in vastus medialis muscle of advanced osteoarthritis patients. *Am J Pathol* 187 (12):2674-2685. <https://doi.org/10.1016/j.ajpath.2017.08.009>

28. Dalle S, Koppo K (2020) Is inflammatory signaling involved in disease-related muscle wasting? Evidence from osteoarthritis, chronic obstructive pulmonary disease and type II diabetes. *Exp Gerontol* 137:110964. <https://doi.org/10.1016/j.exger.2020.110964>

29. Fukumoto Y, Tateuchi H, Ikezoe T, Tsukagoshi R, Akiyama H, So K, Kuroda Y, Ichihashi N (2014) Effects of high-velocity resistance training on muscle function, muscle properties, and physical performance in individuals with hip osteoarthritis: a randomized controlled trial. *Clin Rehabil* 28

415 (1):48-58. <https://doi.org/10.1177/0269215513492161>

416 30. Dos Santos L, Cyrino ES, Antunes M, Santos DA, Sardinha LB (2016) Changes in phase angle and
417 body composition induced by resistance training in older women. *Eur J Clin Nutr* 70 (12):1408-
418 1413. <https://doi.org/10.1038/ejcn.2016.124>

419 31. Ikezoe T, Kobayashi T, Nakamura M, Ichihashi N (2020) Effects of low-load, higher-repetition
420 versus high-load, lower-repetition resistance training not performed to failure on muscle strength,
421 mass, and echo intensity in healthy young men: a time-course study. *J Strength Cond Res* 34
422 (12):3439-3445. <https://doi.org/10.1519/JSC.0000000000002278>

423 32. Niu J, Zhang YQ, Torner J, Nevitt M, Lewis CE, Aliabadi P, Sack B, Clancy M, Sharma L, Felson
424 DT (2009) Is obesity a risk factor for progressive radiographic knee osteoarthritis? *Arthritis Rheum*
425 61 (3):329-335. <https://doi.org/10.1002/art.24337>

426 33. Muraki S, Oka H, Akune T, Mabuchi A, En-yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H,
427 Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N (2009) Prevalence of
428 radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese
429 population-based cohorts: the ROAD study. *Osteoarthritis Cartilage* 17 (9):1137-1143.
430 <https://doi.org/10.1016/j.joca.2009.04.005>

431

432

433 **Table 1.** Knee OA and healthy control participant characteristics

	Healthy control	Knee OA		
	n = 27	n = 41	p-value	95% CI
Age, y	75.6 (4.9)	71.4 (6.0)	0.004	[-6.97, -1.42]
Height, cm	151.7 (5.6)	153.7 (5.6)	0.144	[-0.72, 4.83]
Body mass, kg	48.9 (5.1)	57.2 (9.3)	<0.001	[4.35, 12.17]
Body mass index, kg/m ²	21.3 (2.1)	24.2 (3.7)	<0.001	[1.37, 4.47]
KSS function score, /100	92.0 (11.3)	67.1 (15.4)	<0.001	[-31.87, -18.11]
KSS symptom score, /25	24.0 (1.5)	15.7 (5.1)	<0.001	[-10.35, 6.31]

434

435 Variables are presented as means (SDs).

436

437 Abbreviations: OA, osteoarthritis; CI, confidence interval; KSS, Knee Society Score.

438

439

440 **Table 2.** Muscle property differences between the knee OA and healthy control groups

	Unadjusted mean (SD)		Adjusted mean (SE)		p-value	Adjusted mean difference [95% CI]
	Healthy control	Knee OA	Healthy control	Knee OA		Knee OA against healthy control
Muscle thickness, cm						
Rectus femoris	1.82 (0.35)	1.80 (0.28)	1.88 (0.06)	1.76 (0.05)	0.153	-0.12 [-0.29, 0.05]
Vastus intermedius	1.52 (0.42)	1.58 (0.47)	1.63 (0.09)	1.51 (0.07)	0.312	-0.12 [-0.36, 0.12]
Vastus lateralis	1.65 (0.26)	1.72 (0.36)	1.74 (0.06)	1.66 (0.05)	0.324	-0.08 [-0.25, 0.08]
Vastus medialis	1.95 (0.36)	1.67 (0.42)	2.05 (0.08)	1.61 (0.06)	<0.001	-0.44 [-0.65, -0.23]
Echo intensity, a.u.						
Rectus femoris	92.5 (10.3)	92.7 (13.3)	91.6 (2.5)	93.3 (1.9)	0.611	1.69 [-4.93, 8.32]
Vastus intermedius	56.5 (13.6)	60.0 (14.5)	52.9 (2.8)	62.4 (2.2)	0.014	9.48 [2.01, 16.95]

Vastus lateralis	83.7 (9.7)	89.2 (15.0)	85.2 (2.6)	88.2 (2.0)	0.387	3.02 [-3.91, 9.94]
Vastus medialis	70.1 (8.9)	91.4 (14.3)	69.5 (2.6)	91.8 (2.1)	<0.001	22.24 [15.19, 29.29]
ECW/ICW ratio, a.u.	0.41 (0.10)	0.47 (0.10)	0.39 (0.02)	0.49 (0.01)	<0.001	0.10 [0.05, 0.15]

441

442 Univariate ANOVAs with adjustment for age and BMI were conducted to investigate the muscle property differences between groups. The adjusted mean differences
443 were calculated and presented as the knee OA group values against the healthy control group.

444

445 Abbreviations: OA, osteoarthritis; SD, standard deviation; SE, standard error; CI, confidence interval; ECW/ICW; extracellular-to-intracellular water; a.u., arbitrary
446 unit.

447

448 **Table 3.** Logistic regression and ROC analyses for clarifying muscle degeneration characteristics in knee OA patients

	Logistic regression analysis		ROC analyses				
	OR [95% CI]	p-value	AUC	p-value	Cut-off value	Sensitivity	Specificity
Muscle thickness in vastus medialis	0.47 [0.04, 5.56]	0.547	-				
Echo intensity in vastus medialis	1.19 [1.06, 1.35]	0.004	0.90	<0.001	80.4	0.81	0.89
ECW/ICW ratio	1.19 [1.00, 1.42]	0.047	0.69	0.008	0.45	0.56	0.74

449

450 Logistic regression analyses were conducted with the group (reference, healthy control group = 0; knee OA group = 1) as the dependent variable and the MT, VM EI,

451 and the ECW/ICW ratio as independent variables (after adjusting for age and BMI). The ROC analyses were performed to obtain the AUC, cut-off value, sensitivity,

452 and specificity for characterising muscle degeneration in knee OA.

453

454 Abbreviations: OR, odds ratio; CI, confidence interval; ROC, receiver-operating characteristic; AUC, area under the curve; ECW/ICW; extracellular-to-intracellular
455 water.

Figure Caption

Fig. 1 The setting for the measurement of segmental-bioelectrical impedance spectroscopy in the thigh

To evaluate the bioelectrical impedance of the thigh, two sensing electrodes were placed on the anterior superior iliac spine and the lateral knee articular condyles, and injection electrodes were placed on the dorsal surfaces of the hand and forefoot.

Fig. 2 Representative ultrasound images of the vastus medialis (VM) in (a) healthy control and (b) knee


OA patient

Legend: The arrows represent the thickness of the VM. The entire VM in patients with knee osteoarthritis is white, implying that the mean echo intensity increases.

Fig. 3 Receiver operating characteristic (ROC) curve analyses of echo intensity in the vastus medialis and

extracellular water-to-intracellular water (ECW/ICW) ratio for clarifying muscle degeneration characteristics in patients with knee osteoarthritis.

Legend: The bullet in ROC curve represents a cut-off point.



471

472 **Fig. 1**




Fig. 2





Fig. 3