

Infinite sequences of non-Weierstrass numerical semigroup with odd conductor ¹

神奈川工科大学・基礎・教養教育センター 米田 二良

Jiryō Komeda

Center for Basic Education and Integrated Learning
Kanagawa Institute of Technology

Abstract

We construct infinite sequences of non-Weierstrass almost symmetric numerical semigroups with a fixed odd conductor through dividing by three.

1 Introduction

Let $\mathbb{N}_0$ be the additive monoid of non-negative integers. A submonoid H of $\mathbb{N}_0$ is called a *numerical semigroup* if the complement $\mathbb{N}_0 \setminus H$ is finite. The cardinality of $\mathbb{N}_0 \setminus H$ is called the *genus* of H , which is denoted by $g(H)$. In this article H always stands for a numerical semigroup. We set

$$c(H) = \min\{c \in \mathbb{N}_0 \mid c + \mathbb{N}_0 \subseteq H\},$$

which is called the *conductor* of H . It is known that $c(H) \leq 2g(H)$. A numerical semigroup H is said to be *symmetric* if $c(H) = 2g(H)$. This semigroup has the following symmetric property: For $\gamma \in \mathbb{N}_0$ we have $\gamma \notin H$ if and only if $2g(H) - 1 - \gamma \in H$. A numerical semigroup H is said to be *quasi-symmetric* if $c(H) = 2g(H) - 1$. We set

$$PF(H) = \{\gamma \in \mathbb{N}_0 \setminus H \mid \gamma + h \in H, \text{ all } h \in H > 0\},$$

whose elements are called *pseudo-Frobenius numbers* of H . We have $c(H) - 1 \in PF(H)$. We set $\iota(H) = \#PF(H)$, which is called the *type* of H .

Remark 1.1 We have $c(H) + \iota(H) \leq 2g(H) + 1$. (For example, see [6].)

A numerical semigroup H is said to be *almost symmetric* if the equality $c(H) + \iota(H) = 2g(H) + 1$ holds.

Remark 1.2 i) H is symmetric if and only if $\iota(H) = 1$. In this case H is almost symmetric.
ii) If H is quasi-symmetric, then $\iota(H) = 2$. The converse does not hold. In this case H is also almost symmetric.
iii) If $c(H) = 2g(H) - 2$, then $\iota(H) = 2$ or 3 .

¹This paper is an extended abstract and the details will appear elsewhere.
This work was supported by JSPS KAKENHI Grant Number18K03228.

We set $PF^*(H) = PF(H) \setminus \{c(H) - 1\}$.

Remark 1.3 ([6]) *If H is almost symmetric, then it has the following symmetric property: The map sending γ to $c(H) - 1 - \gamma$ induces a bijection on $PF^*(H)$. The converse is true.*

A curve means a projective non-singular irreducible algebraic curve over an algebraically closed field k of characteristic 0. For a pointed curve (C, P) we denote by $H(P)$

$$\{\alpha \in \mathbb{N}_0 \mid \exists \text{ a rational function } f \text{ on } C \text{ such that } (f)_\infty = \alpha P\}.$$

Then $H(P)$ is a numerical semigroup of genus $g(C)$ where $g(C)$ is the genus of C . A numerical semigroup H is said to be *Weierstrass* if there exists a pointed curve (C, P) with $H(P) = H$.

For any integer $t \geq 2$ we set

$$d_t(H) = \{h' \in \mathbb{N}_0 \mid th' \in H\},$$

which is a numerical semigroup. In this article we are interested in the case $t = 3$. Our main result is the following:

Theorem 1.4 *For any $u \geq 1$ there exist infinite sequences*

$$H_0 \xleftarrow{d_3} H_1 \xleftarrow{d_3} H_2 \xleftarrow{d_3} \dots \xleftarrow{d_3} H_{i-1} \xleftarrow{d_3} H_i \xleftarrow{d_3} \dots$$

of non-Weierstrass numerical semigroups H_i with $c(H_i) = 2g(H_i) - (2u - 1)$ and $\iota(H_i) = 2u$, hence, H_i is almost symmetric for any i .

2 Non-Weierstrass almost symmetric numerical semigroups

In this section we find numerical semigroups in the starting points of the infinite sequences in Theorem 1.4. For a numerical semigroup H the least positive integer in H is denoted by $m(H)$, which is called the *multiplicity* of H . First we state the key lemmas for constructing the numerical semigroups.

Lemma 2.1 *Let u be an integer with $u \geq 1$ and H be a numerical semigroup. Let g be an integer with $g > 4u - 3$, $g \not\equiv u \pmod{3}$ and*

$$g > \max\{m(H) - 2u, u + \frac{3}{2}(c(H) - 1) + \frac{m(H)}{2}, 2u + 2c(H) - 3\}.$$

We set

$$\tilde{H} = 3H \cup \{g + 2u + 3\mathbb{N}_0\} \cup \{2g - 2u - 3r \mid r \in \mathbb{Z} \setminus H\}.$$

Then we have

- i) $\tilde{H}$ is a numerical semigroup and $g(\tilde{H}) = g$.
- ii) $d_3(\tilde{H}) = H$ and $c(\tilde{H}) = 2g(\tilde{H}) - (2u - 1)$.

See [2] for the proof. We give an example which we get by applying the above lemma.

Example 2.1 In Lemma 2.1 let $u = 3$, $H = \langle 2, 3 \rangle$ and $g = 10$. Then $\tilde{H}$ is equal to

$$\begin{aligned} & 3\langle 2, 3 \rangle \cup \{10 + 6 + 3\mathbb{N}_0\} \cup \{20 - 6 - 3r \mid r \in \mathbb{Z} \setminus \langle 2, 3 \rangle\} \\ &= \langle 6, 9 \rangle \cup \{16, 19, 22, \dots\} \cup \{11, 17, 20, 23, \dots\} = \langle 6, 9, 11, 16, 19 \rangle. \end{aligned}$$

In this case, $c(\tilde{H}) = 15 = 2g(\tilde{H}) - 5$.

Lemma 2.2 Let u , H , g and $\tilde{H}$ be as in Lemma 2.1. We set $m = m(H)$. Then we have

$$PF(\tilde{H}) \subseteq \{g + 2u + 3l - 3m \mid 0 \leq l \leq m - 1\} \cup \{2g - 2u + 3m - 3m\}.$$

Hence, $t(\tilde{H}) \leq m + 1$. Moreover, $s_{\max}(\tilde{H}) = 2g - 2u + 3m$.

See [2] for the proof.

Theorem 2.3 Let u , H , g and $\tilde{H}$ be as in Lemma 2.2. Moreover, assume that $2u - 1 \geq m(H)$ and $g \geq 2u + 2c(H) - 1$. Then we have the following:

- i) $t(\tilde{H}) = m(H) + 1$.
- ii) If $2u - 1 = m(H)$, then $c(\tilde{H}) + t(\tilde{H}) = 2g(\tilde{H}) + 1$, hence $\tilde{H}$ is almost symmetric

Using Lemmas 2.1 and 2.2 we can prove the above statement. See [2] for the details of the proof.

To construct the desired non-Weierstrass numerical semigroups we need the known facts.

Remark 2.4 (Oliveira-Stöhr [7]) A numerical semigroup $\tilde{H}$ satisfies that $H = d_3(\tilde{H})$ is non-Weierstrass. If $g(\tilde{H}) \geq 15g(H) + 11$ and $g(\tilde{H}) \not\equiv 1 \pmod{3}$, then $\tilde{H}$ is non-Weierstrass.

Remark 2.5 (Buchweitz [1] and Komeda [4]) Let m be an integer with $m \geq 13$. Then there exists a non-Weierstrass numerical semigroup H with $m(H) = m$.

Remark 2.6 (Komeda [3]) Let $m = 8$ or 12 . Then there exists a non-Weierstrass numerical semigroup H with $m(H) = m$.

Corollary 2.7 Let u be an integer with $u \geq 7$. Then there exists a non-Weierstrass almost symmetric numerical semigroup $\tilde{H}$ with $c(\tilde{H}) = 2g(\tilde{H}) - (2u - 1)$.

Proof. Let m be an odd integer with $m \geq 13$. We set $u = \frac{m+1}{2}$. Then $u \geq 7$. Using Remark 2.5, Lemma 2.1, Remark 2.4 and Theorem 2.3 in this order we get the desired non-Weierstrass numerical semigroups. $\square$

Corollary 2.8 Let u be an integer with $1 \leq u \leq 6$. Then there exists a non-Weierstrass almost symmetric numerical semigroup $\tilde{H}$ with $c(\tilde{H}) = 2g(\tilde{H}) - (2u - 1)$.

Proof. If $u = 2, 3, 4$ (resp. 5, resp. 6), then we take a non-Weierstrass 8-semigroup (resp. 12-semigroup, resp. 14-semigroup H). We construct a non-Weierstrass numerical semigroup $\tilde{H}$ with $t(\tilde{H}) = 2u$ (see Komeda [2] for the details of the proof). In the case $u = 1$, the result is due to Oliveira-Stöhr [7]. $\square$

We give an example in the case $u = 3$, namely, a non-Weierstrass almost symmetric numerical semigroup $\tilde{H}$ with $c(\tilde{H}) = 2g(\tilde{H}) - 5$.

Example 2.2 In Corollary 2.8 let $u = 3$. Let $H = \langle 8, 12, 18, 22, 45, 49 \rangle$, which is a non-Weierstrass numerical semigroup of genus 31 from [3]. Let $g = 15g(H) + 11 = 15 \times 31 + 11 = 476$. We set $\tilde{H} = 3H \cup \{476 + 2 \times 3 + 3\mathbb{N}_0\} \cup \{2 \times 476 - 2 \times 3 - 3r \mid r \in \mathbb{Z} \setminus H\}$. Then $PF(\tilde{H}) = \{482 + 3l - 24 \mid l = 3, \dots, 7\} \cup \{952 - 6 + 24 - 24\}$, i.e., $t(\tilde{H}) = 6$. Hence $\tilde{H}$ is a non-Weierstrass almost symmetric numerical semigroup of genus 476 with $c(\tilde{H}) = 2g(\tilde{H}) - 5$.

3 Proof of Theorem 1.4

To prove Theorem 1.4 we need the following Lemma:

Lemma 3.1 (Komeda [2]) *For any $t \geq 1$ we can construct an infinite sequence of numerical semigroups H_i with $c(H_i) = 2g(H_i) - t$ as follows:*

$$H_0 \xleftarrow{d_3} H_1 \xleftarrow{d_3} H_2 \xleftarrow{d_3} \dots \xleftarrow{d_3} H_{i-1} \xleftarrow{d_3} H_i \xleftarrow{d_3} \dots$$

Here, H_0 is any almost symmetric numerical semigroup with $c(H_0) = 2g(H_0) - t$ and H_i is also almost symmetric for any $i \geq 1$.

We take H_0 in Lemma 3.1 as the non-Weierstrass almost numerical semigroup $\tilde{H}$ in Corollaries 2.7 and 2.8. Using Remark 2.4 and the method of construction of Lemma 3.1 (see Komeda [2]) we get Theorem 1.4.

References

- [1] R.-O. Buchweitz, *On Zariski's criterion for equisingularity and non-smoothable monomial curves*, Preprint 115, University of Hannover, 1980.
- [2] J. Komeda, *Infinite sequences of non-Weierstrass numerical semigroups with a fixed conductor*, In preparation.
- [3] J. Komeda, *Double coverings of curves and non-Weierstrass semigroups*, Communications in Algebra **41** (2013) 312–324.
- [4] J. Komeda, *Non-Weierstrass numerical semigroups*, Semigroup Forum **57** (1988) 157–185.
- [5] J. Komeda and A. Ohbuchi, *On double coverings of a pointed non-singular curve with any Weierstrass semigroup*, Tsukuba J. Math. Soc. **31** (2007) 205–215.

- [6] H. Nari, *Symmetries on almost symmetric numerical semigroups*, Semigroup Forum **86** (2013) 140–154.
- [7] G. Oliveira and K.-O. Stöhr, *Gorenstein curves with quasi-symmetric Weierstrass semigroups*, Geometriae Dedicata **67** (1997) 45–63.
- [8] F. Torres, *Weierstrass points and double coverings of curves with application: Symmetric numerical semigroups which cannot be realized as Weierstrass semigroups*, Manuscripta Math. **83** (1994) 39–58.