

On the concavity of the arithmetic volumes

Hideaki Ikoma

October 23, 2014 (Kinosaki Symposium).

1 Introduction

We pursue the following analogy.

Convex geometry (Bonnesen, Diskant, ...)	Algebraic geometry (Boucksom-Favre -Jonsson, Cutkosky)	Arakelov geometry
convex bodies	nef & big divisors	nef & big —
Euclidean volumes	$\text{vol}(P)$	$\widehat{\text{vol}}(\overline{P})$
mixed volumes	$\deg(P^i \cdot Q^{\dim X - i})$	$\widehat{\deg}(\overline{P}^i \cdot \overline{Q}^{\dim X - i})$
P, Q : homothetic	$P \equiv_{\text{num}} Q$	$\overline{P} \sim_{\mathbb{R}} \overline{Q}$
inradius $s(P, Q) = \sup\{t : P \supset tQ + c, \exists c\}$	$s(P, Q) = \sup\{t : P - tQ \text{ is psef}\}$	$s(\overline{P}, \overline{Q})$
$\vdots$	$\vdots$	$\vdots$

In [7], Yuan showed that the arithmetic volumes also fit into the Brunn-Minkowski inequality, that is, if X is a projective arithmetic variety and $\overline{P}, \overline{Q}$ are pseudo-effective arithmetic ($\mathbb{R}$ -Cartier) $\mathbb{R}$ -divisors on X , then

$$\widehat{\text{vol}}(\overline{P} + \overline{Q})^{\frac{1}{\dim X}} \geq \widehat{\text{vol}}(\overline{P})^{\frac{1}{\dim X}} + \widehat{\text{vol}}(\overline{Q})^{\frac{1}{\dim X}}. \quad (1.1)$$

Our purpose is to obtain equality conditions for this inequality (Theorem 4.5). Let me illustrate the ideas with a toy example.

Toy case Let $A = \text{diag}(a_1, \dots, a_n)$, $B = \text{diag}(b_1, \dots, b_n)$ be diagonal positive-definite matrices. The mixed volumes of A, B are given by

$$V(A^{(k)} \cdot B^{(n-k)}) = \frac{1}{\binom{n}{k}} \sum_{\substack{I \subset \{1, \dots, n\}, \\ \#I=k}} \prod_{i \in I} a_i \cdot \prod_{j \notin I} b_j.$$

The AM-GM inequality says that $\forall k$

$$V(A^{(k)} \cdot B^{(n-k)}) \geq \left(\prod_{\substack{I \subset \{1, \dots, n\}, \\ \#I=k}} \prod_{i \in I} a_i \cdot \prod_{j \notin I} b_j \right)^{\binom{n}{k}^{-1}} = \det(A)^{\frac{k}{n}} \det(B)^{\frac{n-k}{n}} \quad (1.2)$$

and

$$\begin{aligned} \det(A + B) &= \sum_{k=0}^n \binom{n}{k} V(A^{(k)} \cdot B^{(n-k)}) \\ &\geq \sum_{k=0}^n \binom{n}{k} \det(A)^{\frac{k}{n}} \det(B)^{\frac{n-k}{n}} = \left(\det(A)^{\frac{1}{n}} + \det(B)^{\frac{1}{n}} \right)^n. \end{aligned} \quad (1.3)$$

By the equality condition for the AM-GM inequality, we know that equalities in (1.2) $\forall k$ iff $a_1/b_1 = \dots = a_n/b_n$. But we can also go by a very very roundabout way ...

Alexandrov inequality (Corollary 2.6). *Let $C = \text{diag}(c_1, \dots, c_n)$ be another positive definite matrix. Then*

$$V((A + B)^{(n-1)} \cdot C)^{\frac{1}{n-1}} \geq V(A^{(n-1)} \cdot C)^{\frac{1}{n-1}} + V(B^{(n-1)} \cdot C)^{\frac{1}{n-1}}. \quad (1.4)$$

Diskant inequality (Theorem 4.4). *Set $s = s(A, B) = \min\{a_i/b_i\}$. Then*

$$0 \leq \left(V(A^{(n-1)} \cdot B)^{\frac{1}{n-1}} - s \det(B)^{\frac{1}{n-1}} \right)^n \leq V(A^{(n-1)} \cdot B)^{\frac{n}{n-1}} - \det(A) \cdot \det(B)^{\frac{1}{n-1}}. \quad (1.5)$$

Proof. Since $s = \sup\{t \in \mathbb{R} : \det(A - tB) > 0\}$, we have

$$\begin{aligned} \det(A) &= n \int_{t=0}^s V((A - tB)^{(n-1)} \cdot B) dt \\ &\leq n \int_{t=0}^s \left(V(A^{(n-1)} \cdot B)^{\frac{1}{n-1}} - t \det(B)^{\frac{1}{n-1}} \right)^{n-1} dt \end{aligned}$$

by (1.4). We can calculate the last integral. $\square$

If equality in (1.3), then, by (1.5), $s(A, B) = s(B, A)^{-1} = (\det(A)/\det(B))^{\frac{1}{n}}$.

2 Arithmetic $\mathbb{R}$ -divisors

Let me explain some terminology. Let X be a normal projective arithmetic variety, that is, a normal and integral scheme projective and flat over $\text{Spec}(\mathbb{Z})$. We set $d := \dim X - 1$ and denote the rational function field of X by $\text{Rat}(X)$.

Definition 2.1 (Arith. $\mathbb{R}$ -divisors). An *arithmetic $\mathbb{R}$ -divisor* is a pair $\overline{D} = (D, g)$ of an $\mathbb{R}$ -Cartier $\mathbb{R}$ -divisor $D = a_1 D_1 + \cdots + a_l D_l$ and a *D -Green function* $g : (X \setminus \bigcup \text{Supp}(D_i))(\mathbb{C}) \rightarrow \mathbb{R}$, that is, g is continuous, invariant under the complex conjugation, and, $\forall p \in X(\mathbb{C})$,

$$u_p(x) := g(x) + \sum_{i=1}^l a_i \log |f_i(x)|^2 \quad (2.1)$$

extends to a C^0 -function around p , where f_i is a local equation defining D_i around p . We denote the (∞ -dimensional) $\mathbb{R}$ -vector space of all the arith. $\mathbb{R}$ -divisors on X by $\widehat{\text{Div}}(X)$.

Example 2.1. Let $\overline{L} = (L, |\cdot|)$ be a continuous Hermitian line bundle on X , and let s be a non-zero rational section of L . Then $\widehat{\text{div}}(s) := (\text{div}(s), -\log |s|^2)$ is an arith. $\mathbb{R}$ -divisor of C^0 -type.

Example 2.2. A $\phi \in \text{Rat}(X)^\times \otimes_{\mathbb{Z}} \mathbb{R}$ is a formal product $\phi_1^{e_1} \cdots \phi_r^{e_r}$ with $\phi_i \in \text{Rat}(X)^\times$ and $e_i \in \mathbb{R}$. Such ϕ defines an arith. $\mathbb{R}$ -divisor by

$$\widehat{(\phi)} := e_1((\phi_1), -\log |\phi_1|^2) + \cdots + e_r((\phi_r), -\log |\phi_r|^2).$$

Given an arith. $\mathbb{R}$ -divisor $\overline{D}$ on X , we set

$$H^0(D) := \{\phi \in \text{Rat}(X)^\times : (\phi) + D \geq 0\} \cup \{0\}$$

and

$$\widehat{H}^0(\overline{D}) := \{\phi \in H^0(D) : \|\phi\|_{\text{sup}}^g \leq 1\},$$

where $\|\cdot\|_{\text{sup}}^g$ is the sup norm on $H^0(D) \otimes_{\mathbb{Z}} \mathbb{R}$ defined as

$$\|\phi\|_{\text{sup}}^g := \text{ess.sup}_{x \in X(\mathbb{C})} |\phi(x)| \exp \left(\frac{g(x)}{2} \right).$$

An arith. $\mathbb{R}$ -divisor $\overline{D}$ is said to be *effective* if $D \geq 0$ and $g \geq 0$. $\overline{D}$ is effective iff $1 \in \widehat{H}^0(\overline{D})$.

Definition 2.2 (Arith. volumes). The *arith. volume* of $\overline{D}$ is defined as

$$\widehat{\text{vol}}(\overline{D}) = \limsup_{m \rightarrow \infty} \frac{\log \sharp \widehat{H}^0(m\overline{D})}{m^{\dim X} / \dim X!}.$$

Remark 2.1. (1) The function $\overline{D} \rightarrow \widehat{\text{vol}}(\overline{D})$ is positively homogeneous of degree $\dim X$ and continuous (Moriwaki [5]).

(2) $\overline{D}$ is called *big* if $\widehat{\text{vol}}(\overline{D}) > 0$. The cone of all the big arith. $\mathbb{R}$ -divisors is denoted by $\widehat{\text{Big}}(X)$.

(3) $\overline{D}$ is called *pseudo-effective* if $\widehat{\text{vol}}(\overline{A}) > 0$ implies $\widehat{\text{vol}}(\overline{D} + \overline{A}) > 0$.

Let $\overline{D} = (a_1 D_1 + \cdots + a_l D_l, g)$ be an arith. $\mathbb{R}$ -divisor on X . Assume that D_i are all effective and Cartier.

Definition 2.3 (Heights). Given a rational point $x \in X(\overline{\mathbb{Q}})$, we denote the minimal field of definition for x by $K(x)$ and the normalization of $\overline{\{x\}}$ by C_x .

If $(*)$ $x \notin \text{Supp}(D_i)$, $\forall i$, then we define the *height* of x as

$$h_{\overline{D}}(x) := \frac{1}{[K(x) : \mathbb{Q}]} \left(\sum_{i=1}^l a_i \log \# \mathcal{O}_{C_x}(D_i) / \mathcal{O}_{C_x} + \frac{1}{2} \sum_{\sigma: K(x) \rightarrow \mathbb{C}} g(x^\sigma) \right).$$

In general, we can choose a suitable $\phi \in \text{Rat}(X)^\times \otimes_{\mathbb{Z}} \mathbb{R}$ s.t. $\overline{D} + (\widehat{\phi})$ satisfies the condition $(*)$.

(1) $\overline{D}$ is said to be *nef* if D is relatively nef, u_p (2.1) is continuous PSH $\forall p$, and $h_{\overline{D}}(x) \geq 0 \forall x \in X(\overline{\mathbb{Q}})$. The cone of all the nef arith. $\mathbb{R}$ -divisors on X is denoted by $\widehat{\text{Nef}}(X)$.

(2) $\overline{D}$ is said to be *integrable* if $\overline{D}$ can be written as (nef arith. div.) – (nef arith. div.). The $(\infty\text{-dimensional})$ $\mathbb{R}$ -vector space of all the integrable arith. $\mathbb{R}$ -divisors on X is denoted by $\widehat{\text{Int}}(X)$.

Example 2.3. Let $\mathbb{P}_{\mathbb{Z}}^d = \text{Proj}(\mathbb{Z}[X_0, \dots, X_d])$ be the projective space. Let $H := \{X_0 = 0\}$ and let

$$g_{\text{FS}} := \log(1 + |X_1/X_0|^2 + \cdots + |X_d/X_0|^2).$$

Then $\overline{H} = (H, g_{\text{FS}})$ is nef and big (but not arithmetically ample). If we add some $\lambda > 0$, then $(H, g_{\text{FS}} + \lambda)$ is arithmetically ample.

Define the naive height of a rational point $x := (x_0 : \cdots : x_d) \in \mathbb{P}_{\mathbb{Z}}^d(\overline{\mathbb{Q}})$ as

$$h_{\text{naive}}(x) := \frac{1}{[K(x) : \mathbb{Q}]} \sum_{v \in M_{K(x)}} \log \left(\max_i \{|x_i|_v\} \right),$$

which is invariant under the multiplication by $\alpha \in K(x)^\times$ by the product formula. Then we can prove $h_{\text{naive}}(x) = h_{\overline{H}}(x) + O(1)$. (In other words, $h_{\overline{D}} + O(1)$ gives the Weil height associated to D .)

Proposition-Definition 2.2. *There exists a unique, symmetric (in $\overline{D}_0, \dots, \overline{D}_{d-1}$), multilinear, and continuous map*

$$\widehat{\deg} : \overbrace{\widehat{\text{Int}}(X) \times \dots \times \widehat{\text{Int}}(X)}^{d\text{-times}} \times \widehat{\text{Div}}(X) \rightarrow \mathbb{R},$$

$$(\overline{D}_0, \dots, \overline{D}_{d-1}; \overline{D}_d) \mapsto \widehat{\deg}(\overline{D}_0 \cdots \overline{D}_d)$$

having the following properties.

- (1) For every nef arith. $\mathbb{R}$ -divisor $\overline{N}$, $\widehat{\deg}(\overline{N}^{d+1}) = \widehat{\text{vol}}(\overline{N})$.
- (2) If $\overline{D}_0, \dots, \overline{D}_{d-1}$ are nef and $\overline{D}_d$ is pseudo-effective, then $\widehat{\deg}(\overline{D}_0 \cdots \overline{D}_d) \geq 0$.

Remark 2.3. (1) The above map extends the usual arith. intersection numbers of C^∞ -Hermitian line bundles (that is defined by the $*$ -products).

- (2) As in the algebraic case, $\overline{D}$ is pseudo-effective iff, for any normalized blow-up $\varphi : X' \rightarrow X$ and for any nef arith. $\mathbb{R}$ -divisor $\overline{H}$ on X' ,

$$\widehat{\deg}(\overline{H}^d \cdot \varphi^* \overline{D}) \geq 0$$

([4, Theorem 6.4]).

Theorem 2.4 (Faltings, Hriljac, Moriawaki, Yuan-Zhang, ...). *Let $\overline{D}$ be an integrable arith. $\mathbb{R}$ -divisor. Let $\overline{H}_1, \dots, \overline{H}_d$ be nef arith. $\mathbb{R}$ -divisors s.t. $H_{1,\mathbb{Q}}, \dots, H_{d,\mathbb{Q}}$ are all big.*

- (1) If $\deg(D_{\mathbb{Q}} \cdot H_{2,\mathbb{Q}} \cdots H_{d,\mathbb{Q}}) = 0$, then $\widehat{\deg}(\overline{D}^2 \cdot \overline{H}_2 \cdots \overline{H}_d) \leq 0$.
- (2) If $\widehat{\deg}(\overline{D} \cdot \overline{H}_1 \cdots \overline{H}_d) = 0$, then $\widehat{\deg}(\overline{D}^2 \cdot \overline{H}_2 \cdots \overline{H}_d) \leq 0$.

Sketch of proof. (1) By using an arith. Bertini theorem, we can reduce the result to Faltings-Hriljac's theorem (on arith. surfaces).

- (2) Set $t = \deg(D_{\mathbb{Q}} \cdot H_{2,\mathbb{Q}} \cdots H_{d,\mathbb{Q}}) / \deg(H_{1,\mathbb{Q}} \cdot H_{2,\mathbb{Q}} \cdots H_{d,\mathbb{Q}})$ and apply (1) to $\overline{D} - t\overline{H}_1, \overline{H}_2, \dots, \overline{H}_d$.

□

Remark 2.5. Yuan and Zhang [8] have proved that (under suitable conditions) the equality holds in (1) iff $\overline{D}$ comes from $\text{Spec}(H^0(\mathcal{O}_X))$.

Corollary 2.6. *Let $\overline{D}, \overline{E}, \overline{H}_1, \dots, \overline{H}_d$ be nef arith. $\mathbb{R}$ -divisors on X .*

(1) (Teissier-Khovanskii-type) For any i with $1 \leq i \leq d$,

$$\widehat{\deg}(\overline{D}^i \cdot \overline{E}^{(d-i+1)})^2 \geq \widehat{\deg}(\overline{D}^{(i-1)} \cdot \overline{E}^{(d-i+2)}) \cdot \widehat{\deg}(\overline{D}^{(i+1)} \cdot \overline{E}^{(d-i)}).$$

(2) For any k with $1 \leq k \leq d+1$ and for any i with $0 \leq i \leq k$,

$$\begin{aligned} \widehat{\deg}(\overline{D}^i \cdot \overline{E}^{(k-i)} \cdot \overline{H}_k \cdots \overline{H}_d)^k \\ \geq \widehat{\deg}(\overline{D}^k \cdot \overline{H}_k \cdots \overline{H}_d)^i \cdot \widehat{\deg}(\overline{E}^k \cdot \overline{H}_k \cdots \overline{H}_d)^{k-i}. \end{aligned}$$

(3) (Alexandrov-type) For any k with $1 \leq k \leq d+1$,

$$\begin{aligned} \widehat{\deg}((\overline{D} + \overline{E})^k \cdot \overline{H}_k \cdots \overline{H}_d)^{\frac{1}{k}} \\ \geq \widehat{\deg}(\overline{D}^k \cdot \overline{H}_k \cdots \overline{H}_d)^{\frac{1}{k}} + \widehat{\deg}(\overline{E}^k \cdot \overline{H}_k \cdots \overline{H}_d)^{\frac{1}{k}}. \end{aligned}$$

3 Arithmetic positive intersection numbers

An *approximation* of $\overline{D}$ is a pair $(\varphi : X' \rightarrow X, \overline{M})$ having the following properties.

- (1) φ is a projective birational morphism s.t. X' is normal and $X'_{\mathbb{Q}}$ is smooth.
- (2) $\overline{M}$ is a nef arith. $\mathbb{R}$ -divisor on X' s.t. $\varphi^* \overline{D} - \overline{M}$ is pseudo-effective.

We denote the set of all the approximations of $\overline{D}$ by $\widehat{\Theta}(\overline{D})$. If $\overline{D}$ is pseudo-effective, then $\widehat{\Theta}(\overline{D}) \neq \emptyset$.

Definition 3.1. Let $0 \leq n \leq d$. Suppose that $\overline{D}_0, \dots, \overline{D}_n$ are all big and that $\overline{D}_{n+1}, \dots, \overline{D}_d$ are all nef and big. The *arithmetic positive intersection number* of $(\overline{D}_0, \dots, \overline{D}_n; \overline{D}_{n+1}, \dots, \overline{D}_d)$ is defined as

$$\langle \overline{D}_0 \cdots \overline{D}_n \rangle \overline{D}_{n+1} \cdots \overline{D}_d := \sup_{(\varphi, \overline{M}_i) \in \widehat{\Theta}(\overline{D}_i)} \widehat{\deg}(\overline{M}_0 \cdots \overline{M}_n \cdot \varphi^* \overline{D}_{n+1} \cdots \varphi^* \overline{D}_d).$$

Proposition 3.1. (1) The map

$$\begin{aligned} \widehat{\text{Big}}(X)^{\times(n+1)} \times (\widehat{\text{Nef}}(X) \cap \widehat{\text{Big}}(X))^{\times(d-n)} \rightarrow \mathbb{R}, \\ (\overline{D}_0, \dots, \overline{D}_n; \overline{D}_{n+1}, \dots, \overline{D}_d) \mapsto \langle \overline{D}_0 \cdots \overline{D}_n \rangle \overline{D}_{n+1} \cdots \overline{D}_d, \end{aligned}$$

is multi-additive in $\overline{D}_{n+1}, \dots, \overline{D}_d$ and uniquely extends to

$$\begin{aligned} \widehat{\text{Big}}(X)^{\times(n+1)} \times \widehat{\text{Int}}(X)^{\times(d-n)} \rightarrow \mathbb{R}, \\ (\overline{D}_0, \dots, \overline{D}_n; \overline{D}_{n+1}, \dots, \overline{D}_d) \mapsto \langle \overline{D}_0 \cdots \overline{D}_n \rangle \overline{D}_{n+1} \cdots \overline{D}_d. \end{aligned}$$

(2) If $n = d - 1$, then we can further extend the map to

$$\begin{aligned} \widehat{\text{Big}}(X)^{\times d} \times \widehat{\text{Div}}(X) &\rightarrow \mathbb{R}, \\ (\overline{D}_0, \dots, \overline{D}_{d-1}; \overline{D}_d) &\mapsto \langle \overline{D}_0 \cdots \overline{D}_{d-1} \rangle \overline{D}_d. \end{aligned}$$

Theorem 3.2 (Arithmetic Fujita approximation: Yuan [7], Chen [2]). *If $\overline{D}$ is big, then $\widehat{\text{vol}}(\overline{D}) = \langle \overline{D}^{(d+1)} \rangle$.*

By Corollary 2.6 + Theorem 3.2, we have

Proposition 3.3. *Let $\overline{D}, \overline{E}$ be big arith. $\mathbb{R}$ -divisors. For any i with $1 \leq i \leq d - 1$,*

$$\langle \overline{D}^i \cdot \overline{E}^{(d-i+1)} \rangle \geq \widehat{\text{vol}}(\overline{D})^{\frac{i}{d+1}} \cdot \widehat{\text{vol}}(\overline{E})^{\frac{d-i+1}{d+1}}$$

and

$$\langle \overline{D}^d \rangle \overline{E} \geq \langle \overline{D}^d \cdot \overline{E} \rangle \geq \widehat{\text{vol}}(\overline{D})^{\frac{d}{d+1}} \cdot \widehat{\text{vol}}(\overline{E})^{\frac{1}{d+1}}.$$

In particular,

$$\widehat{\text{vol}}(\overline{D} + \overline{E}) \geq \sum_{i=0}^{d+1} \binom{d+1}{i} \langle \overline{D}^i \cdot \overline{E}^{d-i+1} \rangle \geq \left(\widehat{\text{vol}}(\overline{D})^{\frac{1}{d+1}} + \widehat{\text{vol}}(\overline{E})^{\frac{1}{d+1}} \right)^{d+1}.$$

4 Concavity of the arithmetic volumes

Theorem 4.1 (Yuan [6]). *If $\overline{D}, \overline{E}$ are nef arith. $\mathbb{R}$ -divisors, then*

$$\widehat{\text{vol}}(\overline{D} - \overline{E}) \geq \widehat{\text{vol}}(\overline{D}) - (\dim X) \widehat{\text{deg}}(\overline{D}^d \cdot \overline{E}).$$

Corollary 4.2. *The function $\overline{D} \mapsto \widehat{\text{vol}}(\overline{D})$ is differentiable at big arithmetic $\mathbb{R}$ -divisors. If $\overline{D}$ is big and $\overline{E}$ is arbitrary, then*

$$\lim_{t \rightarrow 0} \frac{\widehat{\text{vol}}(\overline{D} + t\overline{E}) - \widehat{\text{vol}}(\overline{D})}{t} = (\dim X) \langle \overline{D}^d \rangle \overline{E}.$$

Suppose that $\overline{D}$ is big. The (positive) height of X is defined as

$$h_D^+(X) := \frac{\widehat{\text{vol}}(\overline{D})}{(\dim X) \text{vol}(D_{\mathbb{Q}})}. \quad (4.1)$$

A sequence (x_n) of rational points on X is said to be *generic* if every subsequence is Zariski dense in X . If (x_n) is generic, then

$$\liminf_{n \rightarrow \infty} h_{\overline{D}}(x_n) \geq h_D^+(X). \quad (4.2)$$

Moreover, if $h_{\overline{D}}(x_n)$ converges to $h_{\overline{D}}^+(X)$ and we move $\overline{D}$ along $\overline{D} + t(0, 2f)$, then the both functions in (4.2) have the same slope at $\overline{D}$. So we can extend the equidistribution theorem (Yuan [6], Berman-Boucksom [1], Chen [3], ...) to the case of big arith. $\mathbb{R}$ -div's.

Corollary 4.3. *Let $f : X(\mathbb{C}) \rightarrow \mathbb{R}$ be a continuous function that is invariant under the complex conjugation, and let (x_n) be a generic sequence of rational points. If $h_{\overline{D}}(x_n)$ converges to $h_{\overline{D}}^+(X)$, then*

$$\lim_{n \rightarrow \infty} \frac{1}{[K(x_n) : \mathbb{Q}]} \sum_{\sigma: K(x_n) \rightarrow \mathbb{C}} f(x_n^\sigma) = \frac{\langle \overline{D}^d \rangle(0, 2f)}{\text{vol}(D_{\mathbb{Q}})}.$$

Theorem 4.4 (Diskant inequality). *If $\overline{D}$ is big and $\overline{P}$ is nef and big, then*

$$0 \leq \left((\langle \overline{D}^d \rangle \overline{P})^{\frac{1}{d}} - s \widehat{\text{vol}}(\overline{P})^{\frac{1}{d}} \right)^{d+1} \leq (\langle \overline{D}^d \rangle \overline{P})^{1+\frac{1}{d}} - \widehat{\text{vol}}(\overline{D}) \cdot \widehat{\text{vol}}(\overline{P})^{\frac{1}{d}},$$

where $s = s(\overline{D}, \overline{P}) = \sup\{t \in \mathbb{R} : \overline{D} - t\overline{P} \text{ is pseudo-effective}\}$.

Theorem 4.5 ([4]). *Let $\overline{D}, \overline{E}$ be nef and big arith. $\mathbb{R}$ -divisors. TFAE.*

- (1) $\widehat{\text{vol}}(\overline{D} + \overline{E})^{\frac{1}{d+1}} = \widehat{\text{vol}}(\overline{D})^{\frac{1}{d+1}} + \widehat{\text{vol}}(\overline{E})^{\frac{1}{d+1}}$.
- (2) For i with $1 \leq i \leq d$, $\widehat{\deg}(\overline{D}^i \cdot \overline{E}^{(d-i+1)}) = \widehat{\text{vol}}(\overline{D})^{\frac{i}{d+1}} \cdot \widehat{\text{vol}}(\overline{E})^{\frac{d-i+1}{d+1}}$.
- (3) $\widehat{\deg}(\overline{D}^d \cdot \overline{E}) = \widehat{\text{vol}}(\overline{D})^{\frac{d}{d+1}} \cdot \widehat{\text{vol}}(\overline{E})^{\frac{1}{d+1}}$.
- (4) $\exists \phi \in \text{Rat}(X)^\times$,

$$\frac{\overline{D}}{\widehat{\text{vol}}(\overline{D})^{\frac{1}{d+1}}} - \frac{\overline{E}}{\widehat{\text{vol}}(\overline{E})^{\frac{1}{d+1}}} = (\widehat{\phi}).$$

Proof of Theorem 4.5. (4) $\Rightarrow$ (1) $\Rightarrow$ (2) $\Rightarrow$ (3) are obvious (by the arith. Teissier-Khovanskii inequalities). The key is (3) $\Rightarrow$ (4).

By the arith. Diskant inequality, we have

$$s = s(\overline{D}, \overline{E}) = \left(\frac{\widehat{\text{vol}}(\overline{D})}{\widehat{\text{vol}}(\overline{E})} \right)^{\frac{1}{d+1}} \quad \text{and} \quad s(\overline{E}, \overline{D}) = s^{-1}.$$

Thus $\overline{D} - s\overline{E}$ and $s\overline{E} - \overline{D}$ are both pseudo-effective. By Moriwaki's Dirichlet theorem, we have (4). $\square$

5 Computation formula

Suppose that $X_{\mathbb{Q}}$ is smooth and fix a volume form ω with $\int_{X(\mathbb{C})} \omega = 1$. Given a big arith. divisor $\overline{D}$, blow-up X along

$$\mathfrak{b}(m\overline{D}) := \text{Image} \left(\left\langle \widehat{H}^0(m\overline{D}) \right\rangle_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_X(-mD) \rightarrow \mathcal{O}_X \right).$$

We obtain $\mu_m : X_m \rightarrow X$ s.t. X_m is normal, the generic fibre $X_{m,\mathbb{Q}}$ is smooth, and $\mathfrak{b}(m\overline{D})\mathcal{O}_{X_m}$ is Cartier. Set

$$F(m\overline{D}) := \mathfrak{b}(m\overline{D})\mathcal{O}_{X_m} \quad \text{and} \quad M(m\overline{D}) := \mu_m^*(m\overline{D}) - F(m\overline{D}).$$

We can endow these divisors with Green functions as follows:

Take an L^2 -ONB $e_1, \dots, e_{r_m}$ for $\left\langle \widehat{H}^0(m\overline{D}) \right\rangle_{\mathbb{C}}$ and let

$$\text{Berg}(m\overline{D})(x) := |e_1(x)|^2 + \dots + |e_{r_m}(x)|^2, \quad x \in X(\mathbb{C}),$$

be the Bergman function.

We can define a continuous Hermitian metric on $\mathcal{O}_{X_m}(F(m\overline{D}))$ by

$$|1_{F(m\overline{D})}|(x) = \sqrt{\text{Berg}(m\overline{D})(\mu_m(x))}, \quad x \in X_m(\mathbb{C}).$$

Then $\overline{F}(m\overline{D}) := (F(m\overline{D}), -\mu_m^* \log \text{Berg}(m\overline{D}))$ is effective and $\overline{M}(m\overline{D}) := \mu_m^*(m\overline{D}) - \overline{F}(m\overline{D})$ is nef.

Suppose that $X_{\mathbb{Q}}$ is smooth. Let $\overline{D}$ be a big arith. divisor.

Theorem 5.1. *Let k be an integer with $1 \leq k \leq d+1$, let $\overline{D}_k, \dots, \overline{D}_n$ be big arith. $\mathbb{R}$ -divisors, and let $\overline{D}_{n+1}, \dots, \overline{D}_d$ be integrable arith. $\mathbb{R}$ -divisors. Then*

$$\langle \overline{D}^k \cdot \overline{D}_k \cdots \overline{D}_n \rangle \overline{D}_{n+1} \cdots \overline{D}_d = \lim_{m \rightarrow \infty} \frac{\langle \overline{D}_k \cdots \overline{D}_n \rangle \overline{M}(m\overline{D})^k \cdot \overline{D}_{n+1} \cdots \overline{D}_d}{m^k}.$$

Corollary 5.2 (Asymptotic orthogonality).

$$\lim_{m \rightarrow \infty} \frac{\widehat{\deg}(\overline{M}(m\overline{D})^d \cdot \overline{F}(m\overline{D}))}{m^{d+1}} = 0.$$

6 Applications

Definition 6.1. An arith. Zariski decomposition of a big arith. $\mathbb{R}$ -divisor $\overline{D}$ is a sum $\overline{D} = \overline{P} + \overline{N}$ s.t. $\overline{P}$ is a nef arith. $\mathbb{R}$ -divisor, $\overline{N}$ is an effective arith. $\mathbb{R}$ -divisor, and $\widehat{\text{vol}}(\overline{P}) = \widehat{\text{vol}}(\overline{D})$.

Remark 6.1. (1) If $\dim X = 2$, then an arith. Zariski decomposition of a big $\overline{D}$ always exists and unique (Moriwaki [5]).

(2) If $\dim X \geq 3$, there exists no arith. Zariski decomposition in general even after any blow-up of X (Moriwaki '11).

Example 6.1. Let $\mathbb{P}_{\mathbb{Z}}^2 = \text{Proj}(\mathbb{Z}[X_0, X_1, X_2])$ and let $z_i := X_i/X_0$ be the affine coordinate. Let $H := \{X_0 = 0\}$ and let

$$g := \max \{ -2, \log |X_1/X_0|^2 + 2, \log |X_2/X_0|^2 + 2 \},$$

which is an H -Green function of PSH-type. Moreover, we can add a “bump” $\rho : \mathbb{P}^2(\mathbb{C}) \rightarrow \mathbb{R}_{\geq 0}$ such that

$$\text{Supp}(\rho) \subseteq \{|z_1| < \exp(-2)\} \times \{|z_2| < \exp(-2)\}.$$

Then $\overline{H} = (H, g + \rho)$ are big and non-nef ($h_{\overline{H}}(1 : 0 : 0) < 0$ or $g + \rho$ is not of PSH-type).

Blow up $\mathbb{P}_{\mathbb{Z}}^2$ with center $(1 : 0 : 0)$, viz. over $\{X_0 \neq 0\}$,

$$\varphi : \text{Proj}(\mathbb{Z}[z_1, z_2][Y_1, Y_2]/(z_2 Y_1 - z_1 Y_2)) \rightarrow \{X_0 \neq 0\}.$$

Then $\varphi^* \overline{H}$ admits an arith. Zariski decomposition. Let E be the exceptional divisor and let $w_{ij} := Y_j/Y_i$. Then the positive part is given by

$$\overline{P} = \left(\varphi^* H - \frac{1}{2} E, \max \{ \log |z_i w_{i1}|, \log |z_i w_{i2}|, \log |z_i w_{i1}|^2 + 2, \log |z_i w_{i2}|^2 + 2 \} \right),$$

the negative part is

$$\overline{N} = \left(\frac{1}{2} E, \max \{ 0, -2 - \max \{ \log |z_i w_{i1}|, \log |z_i w_{i2}| \} \} + \varphi^* \rho \right) \geq 0,$$

and $\widehat{\text{vol}}(\overline{H}) = \widehat{\text{vol}}(\overline{P}) = 5/4$.

Corollary 6.2. *Let $\overline{P}, \overline{Q}$ be nef and big arith. $\mathbb{R}$ -divisors. If $\widehat{\text{vol}}(\overline{P}) = \widehat{\text{vol}}(\overline{Q})$ and $\overline{P} \geq \overline{Q}$, then $\overline{P} = \overline{Q}$.*

Proof.

$$2\widehat{\text{vol}}(\overline{P})^{\frac{1}{d+1}} = \widehat{\text{vol}}(\overline{P})^{\frac{1}{d+1}} + \widehat{\text{vol}}(\overline{Q})^{\frac{1}{d+1}} \leq \widehat{\text{vol}}(\overline{P} + \overline{Q})^{\frac{1}{d+1}} \leq \widehat{\text{vol}}(2\overline{P})^{\frac{1}{d+1}}.$$

Thus, by Theorem 4.5, $\exists \phi \in \text{Rat}(X)^{\times} \otimes_{\mathbb{Z}} \mathbb{R}$ s.t. $\overline{P} - \overline{Q} = (\widehat{\phi}) \geq 0$.

$$(\widehat{\phi}) \geq 0 \quad \Leftrightarrow \quad (\widehat{\phi}) = 0 \quad (\Leftrightarrow \quad \phi \in H^0(\mathcal{O}_X^*) \otimes_{\mathbb{Z}} \mathbb{R}).$$

□

Corollary 6.3. *An arith. Zariski decomposition of a big arith. $\mathbb{R}$ -divisor is (if it exists) unique.*

Acknowledgement I thank the organizers for giving me this opportunity. I thank Professors Moriwaki and Kawaguchi for communications. This research is supported by Research Fellow of Japan Society for the Promotion of Science.

References

- [1] Robert Berman and Sébastien Boucksom. Growth of balls of holomorphic sections and energy at equilibrium. *Inventiones Mathematicae*, 181(2):337–394, 2010.
- [2] Huayi Chen. Arithmetic Fujita approximation. *Annales Scientifiques de l'École Normale Supérieure. Quatrième Série*, 43(4):555–578, 2010.
- [3] Huayi Chen. Differentiability of the arithmetic volume function. *J. Lond. Math. Soc. (2)*, 84(2):365–384, 2011.
- [4] Hideaki Ikoma. On the concavity of the arithmetic volumes. to appear in International Mathematics Research Notices (available at <http://arxiv.org/abs/1310.8424>), 2013.
- [5] Atsushi Moriwaki. Zariski decompositions on arithmetic surfaces. *Publ. Res. Inst. Math. Sci.*, 48(4):799–898, 2012.
- [6] Xinyi Yuan. Big line bundles over arithmetic varieties. *Invent. Math.*, 173(3):603–649, 2008.
- [7] Xinyi Yuan. On volumes of arithmetic line bundles. *Compositio Mathematica*, 145(6):1447–1464, 2009.
- [8] Xinyi Yuan and Shou-Wu Zhang. The arithmetic Hodge index theorem for adelic line bundles I: number fields. preprint available at <http://front.math.ucdavis.edu/1304.3538>, 2013.

Graduate School of Mathematical Sciences,
The University of Tokyo,
Tokyo, 153-8914, Japan
ikoma@ms.u-tokyo.ac.jp