

Oscillatory expression of Hes1 regulates cell proliferation and neuronal differentiation in the embryonic brain

Shohei Ochi^{1,2}, Yui Imaizumi^{1,3}, Hiromi Shimojo^{1*}, Hitoshi Miyachi¹, and Ryoichiro Kageyama^{1,2,4,5*}

¹Institute for Frontier Life and Medical Sciences, Kyoto University, Shogoin-Kawahara, Sakyo-ku, Kyoto 606-8507, Japan.

²Kyoto University Graduate School of Medicine, Kyoto 606-8501, Japan.

³Graduate School of Pharmaceutical Sciences, The University of Tokyo, Tokyo, Japan.

⁴Kyoto University Graduate School of Biostudies, Kyoto 606-8501, Japan.

⁵Institute for Integrated Cell-Material Sciences, Kyoto University, Kyoto 606-8507, Japan.

*Correspondence to:

Ryoichiro Kageyama, Institute for Frontier Life and Medical Sciences, Kyoto University, Shogoin-Kawahara, Sakyo-ku, Kyoto 606-8507, Japan.

E-mail: rkageyam@infront.kyoto-u.ac.jp

Hiromi Shimojo, Institute for Frontier Life and Medical Sciences, Kyoto University, Shogoin-Kawahara, Sakyo-ku, Kyoto 606-8507, Japan.

E-mail: hshimojo@infront.kyoto-u.ac.jp

Running title: Roles of Hes1 oscillations

Keywords: Hes1; neural development; neural progenitor cell; oscillation

Summary statement: Dampened Hes1 oscillations reduce the proliferation of neural progenitor cells (NPCs) and accelerate neuronal differentiation, suggesting the functional significance of robust Hes1 oscillations in efficient proliferation and maintenance of NPCs.

Abstract

The expression of the transcriptional repressor *Hes1* oscillates in many cell types, including neural progenitor cells (NPCs), but the significance of *Hes1* oscillations in development is not fully understood. To examine the effect of altered oscillatory dynamics of *Hes1*, we generated two types of *Hes1* knock-in mice, a shortened (type-1) and an elongated (type-2) *Hes1* gene, and examined their phenotypes focusing on neural development. While both mutations affected *Hes1* oscillations, the type-1 mutation dampened *Hes1* oscillations more severely, resulting in much lower amplitudes. The average levels of *Hes1* expression in type-1-mutant NPCs were also lower than in wild-type NPCs but similar to or slightly higher than those in *Hes1* heterozygous mutant mice, which exhibit no apparent defects. While type-2-mutant mice were apparently normal, type-1-mutant mice displayed smaller brains than wild-type mice and up-regulated proneural gene expression. Furthermore, proliferation of NPCs decreased and cell death increased in type-1-mutant embryos. When *Hes3* and *Hes5* were additionally deleted, neuronal differentiation was also accelerated, leading to microcephaly. Thus, robust *Hes1* oscillations are required for maintenance and proliferation of NPCs and the normal timing of neurogenesis, thereby regulating brain morphogenesis.

Introduction

Many gene activities are oscillatory, and oscillatory expression regulates cellular activities (Levine et al., 2013; Purvis and Lahav, 2013; Isomura and Kageyama, 2014; Johnson and Toettcher, 2019). For example, the transcription factor NF- κ B exhibits nuclear-cytoplasmic shuttling upon activation of this pathway and induces downstream gene expression differently depending on the shuttling frequencies (Hoffmann et al., 2002; Nelson et al., 2004; Ashall et al., 2009). Similarly, phosphorylated ERK (pERK) levels are pulsatile upon activation, and pulsatile but not sustained induction of pERK is important for cell proliferation rates (Albeck et al., 2013; Aoki et al., 2013). Another example is the somite segmentation clock gene *Hes7*, which exhibits oscillatory expression with ~2-hour periodicity in the mouse presomitic mesoderm; each cycle of *Hes7* oscillation leads to formation of a bilateral pair of somites (Bessho et al., 2001; Pourquié, 2011; Oates et al., 2012). *Hes7* represses its own expression by directly binding to the *Hes7* promoter, resulting in its oscillatory expression (Bessho et al.,

2003). When *Hes7* expression is absent or sustained, all somites are severely fused (Bessho et al., 2001; Takashima et al., 2011), while an increase in the frequency of *Hes7* oscillations accelerates somite formation (Harima et al., 2013), indicating that *Hes7* oscillations regulate the pace of somitogenesis. However, the significance of oscillatory expression in the development of other tissues still remains to be analyzed.

The transcriptional repressor *Hes1* is expressed in an oscillatory manner by negative feedback, similar to *Hes7* oscillations, in many cell types including neural progenitor cells (NPCs) (Jouve et al., 2000; Hirata et al., 2002; Shimojo et al., 2008; Bonev et al., 2012; Imayoshi et al., 2013; Chen et al., 2017). In the absence of *Hes1* and its related genes, NPCs are not properly maintained and prematurely differentiate into neurons without sufficient cell proliferation, resulting in microcephaly or anencephaly (Ishibashi et al., 1995; Hatakeyama et al., 2004; Sueda et al., 2019). However, sustained overexpression of *Hes1* also inhibits proliferation of NPCs and enhances their quiescence (Baek et al., 2006), raising the possibility that oscillatory *Hes1* expression is important for proliferation of NPCs. *Hes1* oscillations periodically repress the expression of proneural genes such as *Ascl1* and *Neurog2*, thereby driving their oscillatory expression in NPCs (Shimojo et al., 2008; Imayoshi et al., 2013). By contrast, in differentiating neurons, *Hes1* expression disappears, while *Ascl1* or *Neurog2* expression becomes sustained, indicating that the expression dynamics of proneural genes are different between NPCs and differentiating neurons (Shimojo et al., 2008; Imayoshi et al., 2013). Previous study demonstrated that *Ascl1* has dual opposing functions (Castro et al., 2011), and optogenetic analysis showed that *Ascl1* induces neuronal differentiation when its expression is sustained, but activates the proliferation of NPCs when its expression is oscillatory (Imayoshi et al., 2013). These results suggest that the oscillatory expression of proneural factors, which normally depends on *Hes1* oscillations, is important for efficient proliferation of NPCs (Imayoshi and Kageyama, 2014).

Despite the above findings, the significance of *Hes1* oscillations in development is still obscure. Although sustained overexpression of *Hes1* inhibits proliferation of NPCs, it is not clear which is more responsible for the inhibition of NPC proliferation, sustained (non-oscillatory) or high levels of *Hes1* expression. To address this issue, it is important to examine whether dampened *Hes1* oscillations, in the absence of increased expression levels, affect developmental processes. Mathematical modeling suggested that oscillatory expression is controlled by negative feedback with a delayed timing, which depends on transcriptional delays, the time required for production of mature mRNAs, and that changing such delays alter the oscillatory

dynamics in the absence of increased expression levels (Lewis, 2003; Monk, 2003). We previously found that an important part of transcriptional delays is intronic delays, the time required for transcription and splicing of intronic sequences (Takashima et al., 2011). Deletion of all three introns from the *Hes7* locus shortens the intronic delays, and under such a condition, *Hes7* oscillation dampens without increasing its expression levels (Takashima et al., 2011). Similarly, removal of all intronic sequences from the Delta-like1 (*Dll1*) gene produced a shorter *Dll1* gene and decreased its transcriptional delay. By contrast, insertion of *Dll1* cDNA into the first *Dll1* exon while maintaining all exon and intron sequences produced a longer *Dll1* gene and increased its transcriptional delay (Shimojo et al., 2016), suggesting that the gene length also affects the transcriptional delay.

Therefore, to change the transcriptional delays, we decided to generate two types of *Hes1* mutant mice: a shorter version of *Hes1* with all three introns removed (type-1 mutant), and a longer version of *Hes1*, in which *Hes1* cDNA was inserted between the 5'UTR and the coding sequence (type-2 mutant). We then examined *Hes1* expression dynamics and developmental defects in these mutant mice, focusing on neural development.

Results

Generation of *Hes1* type-1 and type-2 mutant mice

To examine the significance of *Hes1* oscillation in development, we tried to alter the *Hes1* expression dynamics. To this end, we first performed numerical simulations, using the same equations and parameter values to those of *Hes7* oscillations, because *Hes1* has similar oscillation dynamics to *Hes7* (Hirata et al., 2004; Takashima et al., 2011). This simulation suggested that altering the transcriptional delay (T_m , see Materials and methods) would change the dynamics of *Hes1* oscillations (Fig. S1). When the transcriptional delay was shortened compared to that of the wild type (Fig. S1A), oscillations were accelerated (faster oscillations) but dampened (Fig. S1B,C). On the other hand, when the transcriptional delay was lengthened, oscillations with a longer period continued (slower oscillations, Fig. S1D). We previously showed that the intronic delay and the gene length affect the transcriptional delay (Takashima et al., 2011; Harima et al., 2013; Shimojo et al., 2016). Therefore, we generated two types of *Hes1* mutants, a shorter version of *Hes1* with all three introns removed (type-1 mutant, Fig. 1A) and a longer version of *Hes1* in which the coding sequence with a stop codon

of *Hes1* cDNA was inserted between the 5'UTR and the coding sequence in the first exon (type-2 mutant, Fig. 1B).

To determine whether type-1 and type-2 mutations change the transcriptional delay of *Hes1* expression compared to the wild-type control, we generated three reporters, the wild-type *Hes1* gene (control), the intronless *Hes1* gene (type-1 mutant), and the *Hes1* gene with *Hes1* cDNA knocked-in into the first exon (type-2 mutant), each of which contained *Luc2* cDNA fused in-frame at the 5' end of the coding region (Fig. S2A). We introduced each reporter into C2C12 myoblast cells and neural stem (NS) cells and activated *Hes1* promoter with serum treatment to measure the expression kinetics of *Hes1* reporters. Expression of the type-1 mutant reporter occurred ~13.5 min faster than the control in C2C12 cells and 3-12 min faster in NS cells (Fig. S2B,C). By contrast, expression of the type-2 mutant reporter was ~5 min slower in C2C12 cells than the control but did not exhibit a significant difference in NPCs compared to the control (Fig. S2B,C), implying that the transcriptional delay was shorter in the *Hes1* type-1 mutant but longer or the same in the type-2 mutant cells compared to the control.

To generate *Hes1* type-1 mutant mice, a targeting vector containing *Hes1* cDNA and a neomycine expression cassette (pPGK-Neo-pA) was prepared (Fig. 1A). To generate *Hes1* type-2 mutant mice, *Hes1* cDNA and pPGK-Neo-pA were inserted between the 5'UTR and the coding region of the first exon (Fig. 1B). Each targeting vector was introduced into mouse embryonic stem (ES) cells and homologous recombinants were obtained. These recombinants were used to make chimeric mice, which were then crossed with mice ubiquitously expressing flippase (FLP) to remove the neomycine expression cassette (Fig. 1A,B). Heterozygous mutant mice derived from the recombinant ES cells were born normally, and from these mice, we generated homozygous mice (*Hes1* type-1 and type-2 mutant mice). Both *Hes1* type-1 and type-2 mutant mice were born at the Mendelian ratio and grew to fertile adults. We also confirmed that the expression continued in the downstream region of *Hes1* type-2 mutant NPCs (Fig. 1C).

Hes1 expression dynamics in the type-1 and type-2 mutant NPCs

To examine the *Hes1* expression dynamics of *Hes1* type-1 and type-2 mutants, we analyzed time-lapse imaging of the *Hes1* promoter-driven destabilized luciferase reporter (pHes1-Ub-luc, Fig. 2A), which monitors the endogenous *Hes1* expression (Fig. S3) (Masamizu et al., 2006). As previously described, the *Hes1* reporter exhibited oscillatory expression with a period of 173.5 ± 4.4 min in NPCs derived from wild-type mice (Fig. 2B,E,H,K,O and Fig. S4A). *Hes1* expression also oscillated in both *Hes1*

type-1 and type-2 mutant NPCs (Fig. 2C,D,F,G,I,J,L,M and Fig. S4B,C). However, in *Hes1* type-1 mutant NPCs, the amplitude of the oscillation was much smaller (about 37% of the WT, Fig. 2N), and the period was slightly shorter (159.9 ± 2.6 min, Fig. 2O). In *Hes1* type-2 mutant NPCs, the oscillation amplitude was smaller (about 62% of the wild-type, Fig. 2N), but the period was slightly longer (187.0 ± 4.3 min, Fig. 2O). These results indicated that *Hes1* oscillations were severely dampened in type-1 mutant NPCs, while they were less affected in type-2 mutant NPCs. We previously found that *Hes1* oscillations in NPC cultures are very similar to those in brain slices (Shimojo et al., 2008), suggesting that *Hes1* oscillations are also affected in the intact brain of both types of mutants. In the type-2 mutant embryos, we did not find any significant defects in the developing nervous system (Fig. S5 and data not shown), but the type-1 mutant embryos displayed defects in neural development. Thus, we decided to examine the type-1 mutant mice in more detail.

Analyses of anti-*Hes1* immunohistochemical staining in the developing nervous system demonstrated that *Hes1* protein expression was variable and displayed a “salt-and-pepper” pattern in wild-type NPCs (Fig. 3A-C), which might reflect oscillatory expression. In individual cells, there was a wide range of *Hes1* protein levels in the wild-type cortex (Fig. 3B,C). By contrast, in *Hes1* type-1 homozygous mutant NPCs, *Hes1* protein levels were less variable (Fig. 3A-C) and average levels were slightly reduced (Fig. 3D). This reduction in the average level was also confirmed by western blot analysis (Fig. 3E). Thus, *Hes1* type-1 mutant NPCs, in which the transcriptional delay of *Hes1* expression was shortened, exhibited dampened oscillations without increasing the expression levels.

In situ hybridization analysis indicated that *Hes1* mRNA was slightly down-regulated in *Hes1* type-1 homozygous mutant embryos, compared to the wild type (Fig. 3F). Quantification analysis revealed that *Hes1* mRNA levels were ~40% lower at E12.5 in the *Hes1* type-1 mutant brain than in the wild type (Fig. 3H). By contrast, *Hes3* and *Hes5* mRNAs were slightly up-regulated in the *Hes1* type-1 homozygous mutant brain at E12.5, compared to the wild type (Fig. 3G,H), suggesting that *Hes3* and *Hes5* up-regulation may compensate for *Hes1* down-regulation in the *Hes1* type-1 mutant brain at E12.5. However, at E14.5, while the Notch ligand gene *Dll1* was slightly up-regulated, *Hes1*, *Hes3* and *Hes5* mRNA levels were not significantly different in *Hes1* type-1 mutant compared to wild-type embryos (Fig. 3H), suggesting that *Hes* gene expression became normalized at later stages of development.

Defects of neural development in *Hes1* type-1 homozygous mutant embryos

To explore the possible effects of altered *Hes1* expression dynamics on development, we first examined the neural development of *Hes1* type-1 mutant mice. *Hes1* type-1 mutant mice had smaller whole bodies than wild type at E12.5 but recovered afterwards (Fig. 4A,B). The brains of *Hes1* type-1 mutant mice were also smaller than those of the wild type at E12.5 and E16.5 (Fig. 4A,C). Immunohistological and in situ hybridization analyses showed that the proportions of the Nestin⁺ and Pax6⁺ cells in the ventricular zone, Tbr2⁺ intermediate progenitor cells, and Tuj1⁺ neurons were not significantly different between *Hes1* type-1 homozygous mutant and wild-type embryos (Fig. 4D-I). While expression of the proneural gene *Neurog2* was not significantly different, the proneural gene *Ascl1* was slightly up-regulated in *Hes1* type-1 homozygous mutant embryos compared to the wild type at E12.5 (Fig. 4J-L). There were no significant differences in the proportions of Cux1⁺ cells in cortical layers 2-4, Ctip2⁺ cells in layer 5, or Tbr1⁺ cells in layer 6 between *Hes1* type-1 homozygous mutant and wild-type embryos (Fig. 4M-O). Furthermore, differentiation of inhibitory neurons (GAD65⁺, GABA⁺) was not affected in the lateral and medial ganglionic eminence (Fig. 4P,Q). We also compared the proliferation and neuronal differentiation of NPC cultures prepared from E12.5 wild-type and *Hes1* type-1 homozygous mutant embryos but did not find any significant differences (Fig. S6). Together, these results indicated that despite a smaller brain size and up-regulation of *Ascl1*, neural development proceeds almost normally in *Hes1* type-1 homozygous mutant embryos.

We next examined whether cell proliferation and death are affected in *Hes1* type-1 homozygous mutant embryos. The number of phospho-histone H3 (pH3)-positive mitotic cells in the apical (ventricular) but not basal region of *Hes1* type-1 homozygous mutant brain was slightly lower than in the wild type at E12.5, suggesting that proliferation of NPCs decreased in the mutants (Fig. 5A-D). Furthermore, the number of cleaved caspase3⁺ apoptotic cells was slightly higher in the cortical layers (Fig. 5A,E) but not significantly different in the ganglionic eminence (Fig. 5F) of *Hes1* type-1 homozygous mutant brain at E14.5 than in the wild type. These results suggested that decreased cell proliferation and increased apoptosis led to smaller brains of *Hes1* type-1 homozygous mutant embryos.

In *Hes1* type-1 homozygous mutant mice, both Hes1 protein and *Hes1* mRNA levels were lower at E10.5-E12.5 (Fig. 3D,E,H), and therefore the defects observed in these mutant mice might be due to lower levels of Hes1 expression. To test this possibility, we also examined *Hes1* (+/-) mice, which had ~60% and ~30% lower levels of *Hes1* mRNA (Fig. S7C) and Hes1 protein (Fig. S7F), respectively. These mice expressed a lower level of *Hes1* mRNA and a similar level of Hes1 protein compared to

Hes1 type-1 homozygous mutant mice (see Fig. 3D,E,H). The analysis of *Hes1* expression dynamics monitored with pHes1-Ub-luc revealed a similar period but a significant reduction in the amplitude of *Hes1* oscillations (Fig. S7A-C,E). However, this reduction in the amplitude is rather proportional to that in the mean expression value, and the ratio of the amplitude to the mean of *Hes1* oscillations in *Hes1* (+/-) NPCs was similar to that in *Hes1* type-2 homozygous mutant but higher than in *Hes1* type-1 homozygous mutant NPCs (Fig. S7D). In *Hes1* (+/-) mice, we did not find any significant defects in body and brain sizes, proliferation of NPCs, apoptosis, or neuronal differentiation (Fig. S8A,B,E and data not shown). These results suggest that the defects observed in *Hes1* type-1 homozygous mutant mice were due to dampened *Hes1* oscillations rather than to lower levels of *Hes1* expression.

Defects of neural development in *Hes1* type-1-mutant;*Hes5*-null and embryos

The defects observed in the *Hes1* type-1 homozygous mutant brains were rather mild, suggesting that *Hes1*-related genes such as *Hes5*, whose levels are also fluctuating (Imayoshi et al., 2013; Manning et al., 2019), may compensate for the *Hes1* abnormality. Therefore, we next examined *Hes1* type-1-mutant;*Hes5*-null embryos. While neural development is slightly delayed in *Hes5*-null embryos compared to the wild type (Bansod et al., 2017), the size and the final morphologies of *Hes5*-null mice are normal (Ohtsuka et al., 1999); therefore, we used *Hes5*-null mice as controls. The brains of *Hes1* type-1-mutant;*Hes5*-null embryos were significantly smaller than those of *Hes5*-null embryos at E12.5 (Fig. 6A,B). They also tended to be smaller than *Hes5*-null embryos at later stages (Fig. 6A). Immunohistochemical analyses showed that there were more Tuj1⁺ neurons at E14.5 (Fig. 6C,D), while the number of Tbr2⁺ intermediate progenitors increased at E12.5 in *Hes1* type-1-mutant;*Hes5*-null embryos, compared to *Hes5*-null embryos (Fig. 6E,F). Furthermore, there were more Ctip2⁺ layer 5 neurons at E 14.5 in *Hes1* type-1-mutant;*Hes5*-null embryos compared to *Hes5*-null embryos (Fig. 6G,H), although at postnatal day (P)0 there were no longer any significant differences (Fig. 6I,J). In addition, there were more Cux1⁺ layers 2-4 neurons in *Hes1* type-1-mutant;*Hes5*-null embryos than in *Hes5*-null embryos at P0 (Fig. 6I,J). Together, these results suggested that neurogenesis was accelerated in *Hes1* type-1-mutant;*Hes5*-null embryos compared to *Hes5*-null embryos.

We also examined *Hes1*(+/-);*Hes5*(-/-) mice. There were no significant differences in body and brain sizes (Fig. S8C,D), neuronal differentiation (Fig. S8F-I), proliferation (Fig. S8J-M), or apoptosis (Fig. S8N) between *Hes1*(+/-);*Hes5*(-/-) and *Hes5*(-/-) mice, suggesting that neural development proceeds almost normally in

Hes1(+/-);*Hes5*(-/-) mice, and that the defects observed in *Hes1* type-1-mutant;*Hes5*-null embryos are due to dampened *Hes1* oscillations and not to decreased *Hes1* expression levels.

Defects of neural development in *Hes1* type-1-mutant;*Hes3*;*Hes5*-double-null embryos
Because the defects observed in *Hes1* type-1-mutant;*Hes5*-null embryos could be still compensated by *Hes3*, we next examined *Hes1* type-1-mutant in the *Hes3*(-/-);*Hes5*(-/-) background. We previously showed that *Hes3*(-/-);*Hes5*(-/-) mice were mostly normal (Hatakeyama et al., 2004) and therefore used them as controls. The size of the nervous system, including the telencephalon and spinal cord, was significantly smaller in *Hes1* type-1-mutant;*Hes3*;*Hes5*-double-null embryos than in the controls at E12.5 (Fig. 7A-D). Furthermore, neurogenesis was increased in the ventral telencephalon of *Hes1* type-1-mutant;*Hes3*;*Hes5*-double-null embryos compared to the controls (Fig. 7C,E), while apoptosis was not significantly affected (Fig. 7F,G). Thus, *Hes1* type-1-mutation led to microcephaly, indicating that robust *Hes1* oscillations are required for maintenance and proliferation of NPCs and the normal brain morphogenesis.

Discussion

Requirement of robust *Hes1* oscillations for efficient proliferation of NPCs

In this study, to examine the significance of *Hes1* oscillations in NPCs, we made two types of *Hes1*-mutant mice that altered *Hes1* expression dynamics. In *Hes1* type-1 homozygous mutant embryos, *Hes1* oscillations were dampened, and the maintenance and proliferation of NPCs were reduced. The average levels of *Hes1* expression in these mutant NPCs were about 60-70% of those in wild-type NPCs but slightly higher (mRNA level) than or similar (protein level) to those in *Hes1* heterozygous mutant mice, which exhibit no apparent defects. Together, these findings suggest that the defects observed in *Hes1* type-1 mutant mice are due to dampened *Hes1* oscillations rather than to decreased levels of *Hes1*. It was previously shown that *Hes1* oscillations periodically repress the expression of the proneural gene *Ascl1*, driving *Ascl1* oscillations, and that optogenetically induced *Ascl1* oscillations promote proliferation of NPCs (Imayoshi et al., 2013). Furthermore, it was shown that sustained, high levels of *Hes1*, which suppress *Ascl1* expression, promote quiescence in NPCs (Sueda et al., 2019). These results indicated that robust *Hes1* oscillations and resultant proneural gene oscillations are important for efficient proliferation of NPCs. The defects in *Hes1* type-1

homozygous mutant embryos were rather mild, because *Hes1*-related genes such as *Hes3* and *Hes5* compensate for the *Hes1* abnormality. Indeed, when *Hes3* and *Hes5* were additionally deleted, neurogenesis was enhanced in *Hes1* type-1 homozygous mutant embryos. These findings suggest that dampened *Hes1* oscillations affect proneural gene oscillations, which may affect the proliferation and differentiation of NPCs.

In *Hes1* type-2 homozygous mutant mice, *Hes1* oscillations were dampened but less altered than in *Hes1* type-1 homozygous mutant mice, and *Hes1* type-2 homozygous mutant mice were mostly normal. Thus, developmental processes may be somewhat resistant to dampened *Hes1* oscillations. This is partly due to compensation by *Hes1*-related genes such as *Hes3* and *Hes5*, and it remains to be analyzed whether *Hes1* type-2 homozygous mutant embryos exhibit any abnormality in the *Hes3*- and/or *Hes5*-null background.

Deleting introns leads to dampened *Hes1* oscillations

According to mathematical modeling, negative feedback with a delayed timing is required for oscillatory expression, and negative feedback at shorter than optimal delays dampens the oscillations (Fig. S1; Takashima et al., 2011). It has been shown that such delays depend on intronic delays, the time required for transcription and splicing of intronic sequences (Swinburne et al., 2008a; 2008b). Indeed, it was previously shown that deletion of all three introns from the *Hes7* gene accelerates the timing of negative feedback (faster *Hes7* mRNA formation), leading to rather steady (non-oscillatory) *Hes7* expression and severe somite fusion (Takashima et al., 2011). In *Hes1* type-1 homozygous mutant embryos, deletion of all three introns led to dampened *Hes1* oscillations, but *Hes1* expression still oscillated unstably with shorter periodicity and smaller amplitudes. In the developing nervous system, gene expression oscillates out of phase between neighboring cells (Shimojo et al., 2016), and it has been mathematically suggested that whereas in-phase oscillations observed in the presomitic mesoderm are robust and stable, out-of-phase oscillations are unstable (Yoshioka-Kobayashi et al., 2020). Thus, unstable *Hes1* oscillations in NPCs may be susceptible to interfering noises, and in *Hes1* type-1 homozygous mutant embryos, noise produced by cell movements and/or cell division may still induce fluctuations in *Hes1* expression. Another issue is that *Hes1* expression decreased in *Hes1* type-1 mutant NPCs. This decrease could be partly because introns are important for both efficient transport and translation of mRNA. Further analyses are required to understand how the dynamics of in-phase and out-of-phase oscillations are controlled.

As mathematically modeled, the *Hes1* type-2 mutation delays the timing of negative feedback (slower *Hes1* mRNA formation) and leads to *Hes1* oscillations with longer periodicity. However, while modeling suggests that slower negative feedback generates oscillations with higher amplitudes, the actual amplitudes of *Hes1* oscillations in *Hes1* type-2 homozygous mutant embryos were smaller than those in wild-type embryos. This might be partly due to nonsense-mediated mRNA decay (He and Jacobson, 2015), which is often observed when the stop codon is present in the first exon, but the exact mechanism remains to be analyzed.

The mechanism by which Hes1 and proneural gene oscillations regulate cell proliferation and differentiation

Our data suggest that robust Hes1 and proneural gene oscillations are necessary for efficient proliferation of NPCs. This notion agrees well with the previous data demonstrating that dampened *Hes1* oscillations induced by accelerated or delayed Dll1 expression resulted in inhibition of NPC proliferation and acceleration of neurogenesis (Shimojo et al., 2016). Similarly, it was recently shown that Hes1 oscillations drive oscillatory expression of the muscle determination factor MyoD in activated muscle progenitors, and that Hes1 and MyoD oscillations may be important for proliferation and differentiation of muscle progenitors (Lahmann et al., 2019). Furthermore, it was shown that high and sustained levels of Hes1 repress the expression of cell cycle regulators and lead to G1 arrest in cultured fibroblasts, NPCs, and hematopoietic stem cells (Beak et al. 2006; Yu et al., 2006; Sang et al., 2008), suggesting that high and sustained Hes1 is a general feature of maintaining quiescence in many cell types. However, the exact mechanism by which oscillations activate cell proliferation remains to be analyzed. Genes up-regulated by oscillatory and down-regulated by sustained levels of Hes1, or *vice versa*, should be determined to understand whether genes, particularly those involved in cell cycle progression, are differentially controlled by the two modes of *Hes1* expression.

Another important issue is how the timing of neuronal differentiation is regulated. When *Hes1* oscillations were dampened in *Hes1* type-1-mutant;*Hes5*-null and *Hes1* type-1-mutant;*Hes3*;*Hes5*-double-null embryos, neurogenesis was enhanced compared to the controls, suggesting that *Hes1* oscillations play an important role in the normal timing of neurogenesis. Dampened *Hes1* oscillations may also dampen proneural gene oscillations, and rather sustained proneural gene expression might lead to enhanced neurogenesis. It was previously shown that Dll1 oscillations, which are controlled by Hes1 and proneural gene oscillations, are important for the normal timing

of neurogenesis (Shimojo et al., 2016). In *Hes1* type-1-mutant embryos, Dll1 oscillations might be dampened by dampened Hes1 and proneural gene oscillations, and rather sustained Dll1 expression may affect the timing of neurogenesis, because sustained Dll1 overexpression induces neuronal differentiation in a non-cell-autonomous manner (Kawaguchi et al., 2008). Taken together, our results suggest that robust *Hes1* oscillations play an important role in efficient proliferation and maintenance of NPCs and normal brain morphogenesis.

Materials and methods

Animals and generation of *Hes1* type-1 and type-2 mutant mice

All animal experiments were approved by the Institutional Animal Care and Use Committee at Kyoto University. *Hes3*- and *Hes5*-null mice were previously described (Hatakeyama et al., 2004).

For generation of *Hes1* type-1 mutant mice, the Hes1 gene from ATG to the stop codon were replaced with HA sequence fused Hes1 cDNA by BAC recombineering. In the case of *Hes1* type-2 mutant mice, HA-sequence-fused Hes1 cDNA followed by 3'UTR of the Hes1 gene were inserted into between 5'UTR and the first exon of *Hes1* by BAC recombineering. For preparation of the targeting vector of *Hes1* type-1 mutant, a FRT-Neo^R cassette was inserted into the downstream region of 3'UTR. The region from 2.5 kb upstream of HA sequence to 8.0 kb downstream of FRT-Neo^R cassette was retrieved to the pMCS-DTA vector. For preparation of the targeting vector of *Hes1* type-2 mutant, a FRT-Neo^R cassette was inserted into the downstream region of 3'UTR following the Hes1 cDNA sequence, and the region from 1.7 kb upstream of the HA sequence to 6.0 kb downstream of the FRT-Neo^R cassette was retrieved to the pMCS-DTA vector. The linearized targeting vectors were electroporated into TT2 ES cells, and G418-resistant clones were screened by PCR. The homologous recombinant ES clones were injected into eight-cell stage mouse embryos to obtain chimeric mice. Chimeric mice were crossed with pCAG-FLP mice to delete the neo sequence.

Bioluminescence imaging of NPC dissociation culture

NPCs were prepared from the cortex of wild-type embryos and *Hes1* type-1 and type-2 mutant embryos at E12.5 and 14.5. The protocol of dissociation culture by Papain Dissociation System (Worthington) was described previously (Shimojo et al., 2008;

2016). $2.0\text{-}3.0 \times 10^6$ cells/ml of dissociated NPCs were plated into $\phi 12\text{-mm}$ or $\phi 27\text{-mm}$ glass-bottom dishes (IWAKI) with 1 mM luciferin (Nacalai Tesque) in N2/B27 medium (DMEM/F12 supplemented with $1 \times$ N2 [Thermo Fisher Scientific], $1 \times$ B27 [Thermo Fisher Scientific], 1mM N-acetyl-cysteine, 10ng/ml bFGF [Thermo Fisher Scientific]). After one-hour pre-incubation, cells were cultured in medium containing luciferin for measurement of bioluminescence. The dish was placed on the stage of inverted microscope (Olympus IX81) and was maintained at 37°C . Bioluminescence was measured using Olympus objective lens (UPLFLN_40_O) and was transmitted directly to a CCD camera (Princeton Instruments, VersArray 1kb), as previously described (Shimojo et al., 2008; 2016).

Image processing and time-series analysis

Images were processed by Fiji image analysis software. Stack images were applied to Spike Noise Filter to remove signals from cosmic rays and then to Savitzky Golay Temporal Filter to get clear dynamic expression. Period and amplitude of oscillatory expression were measured by single cell tracking (Webb et al., 2016). Detrended fluctuation analysis (DFA) was used for determining self-affinity of a signal, and time-series was detrended by subtracting moving average with 240 min window. To calculate moving averages near edges (initial and late 120 min frames in time-series), data-padding was performed by mirroring. Extreme values were extracted manually, and periods shorter than 60 min and longer than 250 min were discarded. The amplitude was calculated as the average of peak heights relative to the mean (in arbitrary units).

Immunostaining

Embryos and brains were dissected and fixed in 4% paraformaldehyde overnight at 4°C . After being washed in PBS, equilibrated in 25% sucrose/PBS overnight at 4°C and then embedded in OCT compound, tissues were sectioned at $15\text{ }\mu\text{m}$ by cryostat. NPC cultures were fixed in 4% paraformaldehyde for 20 min at room temperature. For immunostaining of Hes1, antigen retrieval was performed in 0.1% Tween20/0.01 M citrate buffer (pH 6.0) using an autoclave (15 min at 105°C). Sections were incubated with primary antibodies (rabbit anti-Hes1, Kobayashi et al., 2009) overnight at 4°C . After being washed in PBS, sections were incubated with secondary antibody (HRP-conjugated anti-rabbit IgG; GE Healthcare) for 90 min at room temperature. After washes in PBS and PBST, color development was enhanced by TSA amplification system (Perkin Elmer) according to the manufacturer's instructions. For immunostaining of Pax6, Tbr2, Tbr1, and Cux1, antigen retrieval was performed in

0.1% Tween20/0.01 M citrate buffer (pH 6.0) using an autoclave (15 min at 105°C). Primary antibodies used were as follows: mouse anti-Pax6 (Abcam), rabbit anti-Tbr2 (Abcam), rabbit anti-Tbr1 (MERCK), rabbit anti-Cux1 (Santa Cruz Biotechnology), mouse anti-Nestin (BD Biosciences), rabbit anti-Tuj1 (Covance), mouse anti-pH3 (MERCK), rabbit anti-casp3 (CST), mouse anti-GAD65 (Chemicon), and rabbit anti-GABA (Sigma). Sections were incubated with primary antibodies overnight at 4°C. The secondary antibodies used were as follows: Alexa 488-conjugated anti-mouse IgG (Molecular Probes), Alexa 594-conjugated anti-rabbit IgG (Molecular Probes), Alexa 488-conjugated anti-rabbit IgG (Molecular Probes), and Alexa 594-conjugated anti-rat IgG (Molecular Probes). Antibodies were listed in Table S1.

In situ hybridization

Preparation of DIG-labeled antisense RNA probes and in situ hybridization using NBT/BCIP (Roche) detection were performed as described previously (Bessho et al., 2001; Shimojo et al., 2008).

PCR analysis

Samples were collected from the telencephalon of each genotype embryo, total RNA was extracted, and then reverse transcription (RT) reaction was performed. Real-time PCR (qPCR) was performed as described previously (Kobayashi et al., 2009). Quantified values of RNA were normalized with those of GAPDH. Primers were listed in Table S2. At least three embryos of each genotype were examined.

Analysis of the distribution of Hes1 protein expression

The intensity of Hes1 immunostaining of individual NPCs in the cortex of each genotype was measured by Fiji software, as previously described (Baek et al., 2006).

Western blot analysis

Western blot analysis of Hes1 was done, as previously described (Hirata et al., 2002).

Measurements of the size of whole body and the telencephalon

The image of embryos was obtained by stereomicroscope (Leica). The length of long axis and the area of whole body and the telencephalon were measured by Fiji software. Quantified values of mutants were compared to those of control mice. At least three embryos of each genotype were examined.

Measurement of the formation of neuronal layers

The area or the number of cells expressing neuronal marker (Tuj1, Tbr2, Cux1, Ctip2, Tbr1) in the cortex of each genotype was measured by Fiji software. For the quantification of neuronal layers, serial 16- μ m sections were cut, and every 20 sections were used. At least three embryos of each genotype were examined.

Measurement of the number of mitotic and apoptotic cells

pH3⁺ mitotic cells in the cortex were classified into apical or basal mitoses (Cárdenas et al., 2018). The numbers of pH3⁺ mitotic and cleaved caspase3⁺ apoptotic cells were counted. At least three embryos of each genotype were examined.

NPC and neurosphere cultures

NPC and neurosphere cultures were performed, as previously described (Ohtsuka et al., 2006; Imayoshi et al., 2013). For cell cycle exit analysis, 10 μ M BrdU was added to NPCs, which were then fixed 24 hours later and subjected to immunostaining for BrdU and Ki67. The proportion of BrdU⁺Ki67⁻ cell number over BrdU⁺ cell number was calculated.

Delay measurement of Hes1 expression

Luc2-fused control, *Hes1* type-1, and *Hes1* type-2 vectors were transfected into C2C12 cells and NPCs. Synchronized Hes1 expression in transfected cells was induced by 5% FBS/DMEM after starvation with 0.2% FBS/DMEM treatment (Harima et al., 2013). The kinetics of Hes1 expression in these cells were measured by photomultiplier tube (CL24B-LIC/B, Churitsu Electric Corp.).

Mathematical simulation

Hes1 oscillations are simulated with the following equations described previously (Hirata et al., 2004; Takashima et al., 2011; Harima et al., 2012)

$$\frac{dp(t)}{dt} = am(t - T_p) - bp(t)$$

$$\frac{dm(t)}{dt} = f(p(t - T_m)) - cm(t)$$

where p(t) and m(t) are the quantities of functional Hes1 protein and *Hes1* mRNA per cell at time t, respectively, and f(p) is the rate of initiation of transcription, which

depends on the amount of the protein, p , present at the time of initiation. a is the rate constant for translation, and b and c are the degradation rate constants for Hes1 protein and *Hes1* mRNA, respectively, which are simply related to the half-lives of the molecules:

$$b = \frac{\ln 2}{\tau_p}, c = \frac{\ln 2}{\tau_m}$$

Because transcription is inhibited by Hes1 protein, which acts as a dimer, we assume

$$f(p) = \frac{k}{1 + (p/p_{\text{crit}})^2}$$

where k is the number of molecules of *Hes1* mRNA synthesized per unit time in the absence of inhibition and p_{crit} is the amount of protein that gives half-maximal inhibition. We set $a = 4.5$ protein molecules per mRNA molecule per min, $p_{\text{crit}} = 40$ molecules per cell, $k = 33$ mRNA molecules per cell per min, and $\tau_m = 3$ min. We assume that the Hes1 protein half-life $\tau_p = 20$ min, $T_p = 8$ min, and $T_m = 29$ min. Under these conditions, oscillatory expression continues. In $T_m = 34$ min (5 min longer), oscillations are maintained. In contrast, when $T_m = 10$ min (19 min shorter) and $T_m = 15.5$ min (13.5 min shorter), oscillations are damped. The figure of Hes1 dynamics was exported from the mathematical model using XPPAUT.

Acknowledgements

We thank Dai Watanabe, Satoshi Yawata, and Chika Nishimura for technical help. This work was supported by Core Research for Evolutional Science and Technology (CREST) (JPMJCR12W2 to R.K.), Grant-in-Aid for Scientific Research on Innovative Areas (16H06480 to R.K.) from Ministry of Education, Culture, Sports, Science, and Technology (MEXT), Japan, and Scientific Research (C) (18K06254 to H.S.) and Research Fellowship for Young Scientists (17J02922 to S.O.) from Japan Society for the Promotion of Science.

References

Albeck, J., Mills, G., and Brugge, J. (2013). Frequency-modulated pulses of ERK activity transmit quantitative proliferation signals. *Mol. Cell* **49**, 249–261.

Aoki, K., Kumagai, Y., Sakurai, A., Komatsu, N., Fujita, Y., Shionyu, C., and Matsuda, M. (2013). Stochastic ERK activation induced by noise and cell-to-cell propagation regulates cell density-dependent proliferation. *Mol. Cell* **52**, 529–540.

Ashall, L., Horton, C.A., Nelson, D.E., Paszek, P., Harper, C.V., Sillitoe, K., Ryan, S., Spiller, D.G., Unitt, J.F., Broomhead, D.S., Kell, D.B., Rand, D.A., Sée, V. and White, M.R.H. (2009). Pulsatile stimulation determines timing and specificity of NF- κ B-dependent transcription. *Science* **324**, 242–246.

Baek, J.H., Hatakeyama, J., Sakamoto, S., Ohtsuka, T., and Kageyama, R. (2006). Persistent and high levels of *Hes1* expression regulate boundary formation in the developing central nervous system. *Development* **133**, 2467–2476.

Bansod, S., Kageyama, R., and Ohtsuka, T. (2017). *Hes5* regulates the transition timing of neurogenesis and gliogenesis in mammalian neocortical development. *Development* **144**, 3156–3167.

Bessho, Y., Sakata, R., Komatsu, S., Shiota, K., Yamada, S., and Kageyama, R. (2001). Dynamic expression and essential functions of *Hes7* in somite segmentation. *Genes Dev.* **15**, 2642–2647.

Bessho, Y., Hirata, H., Masamizu, Y., and Kageyama, R. (2003). Periodic repression by the bHLH factor *Hes7* is an essential mechanism for the somite segmentation clock. *Genes Dev.* **17**, 1451–1456.

Bonev, B., Stanley, P. and Papalopulu, N. (2012). MicroRNA-9 modulates *Hes1* ultradian oscillations by forming a double-negative feedback loop. *Cell Rep.* **2**, 10–18.

Cárdenas, A., Villalba, A., De Juan Romero, C., Picó, E., Kyrousi, C., Tzika, A.C., Tessier-Lavigne, M., Ma, L., Drukker, M., Cappello, S. and Borrell, V. (2018). Evolution of cortical neurogenesis in amniotes controlled by Robo signaling levels. *Cell* **174**, 590–606.

Castro, D.S., Martynoga, B., Parras, C., Ramesh, V., Pacary, E., Johnston, C., Drechsel, D., Lebel-Potter, M., Garcia, L.G., Hunt, C., et al. (2011). A novel function of the proneural factor *Ascl1* in progenitor proliferation identified by genome-wide characterization of its targets. *Genes Dev* **25**, 930-945.

Chen, F., Zhang, C., Wu, H., Ma, Y., Luo, X., Gong, X., Jiang, F., Gui, Y., Zhang, H. and Lu, F. (2017). The E3 ubiquitin ligase SCF^{BXL14} complex stimulates neuronal differentiation by targeting the Notch signaling factor HES1 for proteolysis. *J. Biol. Chem.* **292**, 20100-20112.

Harima, Y., Takashima, Y., Ueda, Y., Ohtsuka, T., and Kageyama, R. (2013). Accelerating the tempo of the segmentation clock by reducing the number of introns in the *Hes7* gene. *Cell Rep.* **3**, 1-7.

Hatakeyama, J., Bessho, Y., Katoh, K., Ookawara, S., Fujioka, M., Guillemot, F., and Kageyama, R. (2004). *Hes* genes regulate size, shape and histogenesis of the nervous system by control of the timing of neural stem cell differentiation. *Development* **131**, 5539-5550.

He, F. and Jacobson, A. (2015). Nonsense-mediated mRNA decay: degradation of defective transcripts is only part of the story. *Annu. Rev. Genet.* **49**, 339-366.

Hirata, H., Yoshiura, S., Ohtsuka, T., Bessho, Y., Harada, T., Yoshikawa, K. and Kageyama, R. (2002). Oscillatory expression of the bHLH factor *Hes1* regulated by a negative feedback loop. *Science* **298**, 840-843.

Hirata, H., Bessho, Y., Kokubu, H., Masamizu, Y., Yamada, S., Lewis, J. and Kageyama, R. (2004). Instability of *Hes7* protein is critical for the somite segmentation clock. *Nat. Genet.* **36**, 750-754.

Hoffmann, A., Levchenko, A., Scott, M.L. and Baltimore, D. (2002). The I κ B–NF- κ B signaling module: temporal control and selective gene activation. *Science* **298**, 1241-1245.

Imayoshi, I., Isomura, A., Harima, Y., Kawaguchi, K., Kori, H., Miyachi, H., Fujiwara, T.K., Ishidate, F. and Kageyama, R. (2013). Oscillatory control of factors

determining multipotency and fate in mouse neural progenitors. *Science* **342**, 1203-1208.

Imayoshi, I. and Kageyama, R. (2014). bHLH factors in self-renewal, multipotency, and fate choice of neural progenitor cells. *Neuron* **82**, 9-23.

Ishibashi, M., Ang, S.-L., Shiota, K., Nakanishi, S., Kageyama, R., and Guillemot, F. (1995). Targeted disruption of mammalian *hairy* and *Enhancer of split* homolog-1 (*HES-1*) leads to up-regulation of neural helix-loop-helix factors, premature neurogenesis and severe neural tube defects. *Genes Dev.* **9**, 3136-3148.

Isomura, A. and, Kageyama, R. (2014). Ultradian oscillators: rhythms and cell fate decisions. *Development* **141**, 3627-3636.

Johnson, H.E. and Toettcher, J.E. (2019). Signaling dynamics control cell fate in the early *Drosophila* embryo. *Dev. Cell* **48**, 361-370.

Jouve, C., Palmeirim, I., Henrique, D., Beckers, J., Gossler, A., Ish-Horowicz, D. and Pourquié O. (2000). Notch signalling is required for cyclic expression of the hairy-like gene *HES1* in the presomitic mesoderm. *Development* **127**, 1421-1429.

Kawaguchi, D., Yoshimatsu, T., Hozumi, K. and Gotoh, Y. (2008). Selection of differentiating cells by different levels of delta-like 1 among neural precursor cells in the developing mouse telencephalon. *Development* **135**, 3849-3858.

Kobayashi, T., Mizuno, H., Imayoshi, I., Furusawa, C., Shirahige, K. and Kageyama, R. (2009). The cyclic gene *Hes1* contributes to diverse differentiation responses of embryonic stem cells. *Genes Dev.* **23**, 1870-1875.

Lahman, I., Bröhl, D., Zyrianova, T., Isomura, A., Zhang, Y., Kapoor, V., Czajkowski, M.T., Griger, J., Ruffault, P.-L., Mademtzoglou, D., Zammit, P.S., Wunderlich, T., Spuler, S., Kühn, R., Preibisch, S., Wolf, J., Kageyama, R. and Birchmeier, C. (2018). Dynamic oscillations of *Hes1* and *MyoD* protein regulate the maintenance of muscle stem cells. *Genes Dev.* **33**, 524-534.

Levine, J.H., Lin, Y. and Elowitz, M.B. (2013). Functional roles of pulsing in genetic

circuits. *Science* **342**, 1193-1200.

Lewis, J. (2003). Autoinhibition with transcriptional delay: a simple mechanism for the zebrafish somitogenesis oscillator. *Curr. Biol.* **13**, 1398-1408.

Manning, C.S., Biga, V., Boyd, J., Kursawe, J., Ymisson, B., Spiller, D.G., Sanderson, C.M., Galla, T., Rattray, M. and Papalopulu, N. (2019). Quantitative single-cell live imaging links HES5 dynamics with cell-state and fate in murine neurogenesis. *Nat. Commun.* **10**, 2835.

Masamizu, Y., Ohtsuka, T., Takashima, Y., Nagahara, H., Takenaka, Y., Yoshikawa, K., Okamura, H. and Kageyama, R. (2006). Real-time imaging of the somite segmentation clock: revelation of unstable oscillators in the individual presomitic mesoderm cells. *Proc. Natl. Acad. Sci. USA* **103**, 1313-1318.

Monk, N.A. (2003). Oscillatory expression of Hes1, p53, and NF-kappaB driven by transcriptional time delays. *Curr. Biol.* **19**, 1409-1413.

Nelson, D.E., Ihekweaba, A.E.C., Elliott, M., Johnson, J.R., Gibney, C.A., Foreman, B.E., Nelson, G., See, V., Horton, C.A., Spiller, D.G., et al. (2004). Oscillations in NF- κ B signaling control the dynamics of gene expression. *Science* **306**, 704-708.

Oates, A.C., Morelli, L.G. and Ares, S. (2012). Patterning embryos with oscillations: structure, function and dynamics of the vertebrate segmentation clock. *Development* **139**, 625-639.

Ohtsuka, T., Imayoshi, I., Shimojo, H., Nishi, E., Kageyama, R. and McConnell, S.K. (2006). Visualization of embryonic neural stem cells using Hes promoters in transgenic mice. *Mol. Cell. Neurosci.* **31**, 109-122.

Ohtsuka, T., Ishibashi, M., Gradwohl, G., Nakanishi, S., Guillemot, F. and Kageyama, R. (1999) *Hes1* and *Hes5* as Notch effectors in mammalian neuronal differentiation. *EMBO J.* **18**, 2196-2207.

Pourquié O. (2011). Vertebrate segmentation: from cyclic gene networks to scoliosis.

Cell **145**, 650-663.

Purvis, J.E. and Lahav, G. (2013). Encoding and decoding cellular information through signaling dynamics. *Cell* **152**, 945-956.

Sang, L., Collier, H.A. and Roberts, J.M. (2008). Control of the reversibility of cellular quiescence by the transcriptional repressor HES1. *Science* **321**, 1095-1100.

Shimojo, H., Ohtsuka, T. and Kageyama, R. (2008). Oscillations in notch signaling regulate maintenance of neural progenitors. *Neuron* **58**, 52-64.

Shimojo, H., Isomura, A., Ohtsuka, T., Kori, H., Miyachi, H. and Kageyama, R. (2016). Oscillatory control of Delta-like1 in cell interactions regulates dynamic gene expression and tissue morphogenesis. *Genes Dev.* **30**, 102-116.

Sueda, R., Imayoshi, I., Harima, Y., and Kageyama, R. (2019). High Hes1 expression and resultant Ascl1 suppression regulate quiescent versus active neural stem cells in the adult mouse brain. *Genes Dev.*, 33, 511-523.

Swinburne, I.A., Miguez, D. G., Landgraf, D. and Silver, P.A. (2008a). Intron length increases oscillatory periods of gene expression in animal cells Email alerting service Intron length increases oscillatory periods of gene expression in animal cells. *Genes Dev.* **22**, 2342–2346.

Swinburne, I.A. and Silver, P.A. (2008b). Intron delays and transcriptional timing during development. *Dev. Cell* **14**, 324–330.

Takashima, Y., Ohtsuka, T., González, A., Miyachi, H. and Kageyama, R. (2011). Intronic delay is essential for oscillatory expression in the segmentation clock. *Proc. Natl. Acad. Sci. USA* **108**, 3300-3305.

Webb, A.B., Lengyel, I.M., Jörg, D.J., Valentin, G., Jülicher, F., Morelli, L.G. and Oates, A.C. (2016). Persistence, period and precision of autonomous cellular oscillators from the zebrafish segmentation clock. *eLife* **5**, e08438.

Yoshioka-Kobayashi, K., Matsumiya, M., Niino, Y., Isomura, A., Kori, H.,

Miyawaki, A., and Kageyama, R. (2020). Control of coupling delay for synchronized oscillations in the segmentation clock. *Nature*, in press.

Yu, X., Alder, J.K., Chun, J.H., Friedman, A.D., Heimfeld, S., Cheng, L. and Civin, C.I. (2006). HES1 inhibits cycling of hematopoietic progenitor cells via DNA binding. *Stem Cells* **24**, 876-888.

Figure legends

Figure 1. Generation of *Hes1* type-1 and type-2 mutant mice. (A,B) Strategies for generating *Hes1* type-1 (A) and type-2 (B) mutant mice. Closed and open squares indicate coding and non-coding regions of *Hes1*, respectively. The targeting vector was introduced into mouse ES cells, and heterozygous mutant mice were obtained. Then, a neomycin resistance gene cassette flanked by FRT sites was removed by crossing with pCAG-FLP mice. Arrowheads indicate the positions of primers. (C) PCR analysis of *Hes1* gene transcription of *Hes1* type-2 mutant mice using the primers indicated in (B).

Figure 2. *Hes1* expression dynamics in *Hes1* type-1 and type-2 mutant mice. (A) Structure of the *Hes1* reporter. Ubiquitinated luciferase (Ub-luc) cDNA was placed under the control of 2.5-kb of the *Hes1* promoter. (B-D) Mice carrying the *Hes1* reporter were crossed with wild-type (B), *Hes1* type-1 mutant (C), and *Hes1* type-2 mutant (D) mice. (E-G) Luciferase activities were monitored in NPCs of (B-D), respectively. (H-J) Quantification of the luciferase activities monitored in (E-G). (K-M) Detrended signals of the luciferase activities monitored in (E-G). (N,O) Average with SEM of the amplitude (N) and period (O) of *Hes1* oscillations. At least 100 cells were examined for each genotype. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, Student's *t*-test.

Figure 3. *Hes1* and *Hes1*-related gene expression in *Hes1* type-1 mutant mice. (A) Immunohistochemistry of *Hes1* in the telencephalon of E10.5 wild-type (WT, upper) and *Hes1* type-1 mutant mice (lower). (B) Distribution of *Hes1* expression levels in NPCs of WT and *Hes1* type-1 mutant mice at E10.5. (C) Box plots of *Hes1* expression levels in NPCs at E10.5. (D) Average levels with SEM of *Hes1* expression in WT and *Hes1* type-1 mutant at E10.5. (E) Western blot analysis of *Hes1* protein expression in WT and *Hes1* type-1 mutant NPCs. $n = 3$ for each genotype. (F,G) In situ hybridization of *Hes1* (F) and *Hes5* (G) in the WT (upper) and *Hes1* type-1 mutant (lower)

telencephalon. (H) Real-time RT-PCR analyses of gene expression in WT and *Hes1* type-1 mutant mice. Values with SEM relative to those at E12.5 are shown. $n = 3$ for each genotype. $*p < 0.05$, Student's *t*-test. Scale bars = 50 μm (A); 300 μm (F,G).

Figure 4. Analyses of neural development in *Hes1* type-1 mutant mice. (A) Images of the whole body at E12.5 and E14.5 and the telencephalon at E14.5 and E16.5 of wild-type (WT) and *Hes1* type-1 mutant mice. (B,C) Quantification of the size (with SEM) of whole body (B) and the telencephalon (C) of WT and *Hes1* type-1 mutant mouse embryos. For WT, $n = 4$ (E10.5), 10 (E12.5), 7 (E14.5), and 4 (E16.5). For *Hes1* type-1 mutant, $n = 3$ (E10.5), 10 (E12.5), 7 (E14.5), and 4 (E16.5). (D,E) Immunohistochemistry of Nestin, Tuj1, Pax6, and Tbr2 in the cortex of WT and *Hes1* type-1 mutant mice at E10.5 and E12.5. (F) Quantification of the thickness (with SEM) of Tuj1⁺ and Tbr2⁺ regions in the cortex of WT and *Hes1* type-1 mutant mice. (G,H,J,K) *In situ* hybridization of *Pax6* (G), *Tbr2* (H), *Ascl1* (J), and *Neurog2* (K) in the telencephalon of WT and *Hes1* type-1 mutant mice at E10.5 and E12.5. (I,L) Real-time RT-PCR analyses of *Pax6* (I), *Tbr2* (I), *Ascl1* (L), and *Neurog2* (L) mRNA expression levels (with SEM) in NPCs in the telencephalon of WT and *Hes1* type-1 mutant mice at E12.5 and E14.5. (M,N) Immunohistochemistry of Cux1, Ctip2, and Tbr1 in the cortex of WT and *Hes1* type-1 mutant mice at postnatal day 0 (P0). (O) Quantification of the thickness (with SEM) of Cux1⁺, Ctip2⁺, and Tbr1⁺ regions in the cortex of WT and *Hes1* type-1 mutant mice at P0. (P) Immunohistochemistry of GAD65 and GABA in the lateral (LGE) and medial ganglionic eminence (MGE). (Q) Quantification of the area (with SEM) of GAD45⁺ and GABA⁺ regions in the LGE of WT and *Hes1* type-1 mutant mice at P0. At least 3 samples were examined for quantification. $*p < 0.05$, Student's *t*-test; a.u. = arbitrary units. Scale bars = 3 mm (A); 400 μm (D left, E left, M left, N left); 300 μm (G,H,J,K); 200 μm (D right, E right); 100 μm (M right, N right, P); 50 μm (D middle, E middle).

Figure 5. Analyses of cell proliferation and cell death in *Hes1* type-1 mutant mice. (A) Immunohistochemistry of phospho-Histone H3 (pH3), cleaved caspase3 (casp3), and Ki67 in the cortex of WT and *Hes1* type-1 mutant mice at E10.5, E12.5, and E14.5. Scale bars = 100 μm . (B-F) Quantification of apical (B) and basal mitotic cells (C), total number of mitotic cells (D), and apoptotic cells (E) in the cortex and the ganglionic eminence (F) of wild-type and *Hes1* type-1 mutant mice at E10.5, E12.5, and E14.5. Average values with SEM are shown. Scale bars = 500 μm .

Figure 6. Analyses of neurogenesis in *Hes1* type-1-mutant;*Hes5*-null mice. (A) Images of the whole body at E12.5 and the brain at E16.5 of *Hes5*-null and *Hes1* type-1 mutant;*Hes5*-null mice. (B) Quantification of the whole body and telencephalon sizes (with SEM) of *Hes5*-null mice and *Hes1* type-1-mutant;*Hes5*-null mice. For *Hes5*(-/-), n = 6 (E12.5) and 6 (E16.5). For *Hes1* type-1 mutant;*Hes5*(-/-), n = 7 (E12.5) and 7 (E16.5). (C,E,G,I) Immunohistochemistry of Nestin (C), Tuj1 (C), Tbr2 (E), Ctjp2 (G,I), and Cux1 (I) in the cortex of *Hes5*-null and *Hes1* type-1-mutant;*Hes5*-null mice. (D,F,H,J) Quantification of the proportions (with SEM) of Tuj1⁺ (D), Tbr2⁺ (F), Ctjp2⁺ (H,J), and Cux1⁺ (J) regions in the cortex of *Hes5*-null mice and *Hes1* type-1-mutant;*Hes5*-null mice. At least 3 samples were examined for quantification. *p < 0.05, **p < 0.01, ***p < 0.001, Student's *t*-test. Scale bars = 2 mm (A); 400 μm (C); 200 μm (E,G); 100 μm (I).

Figure 7. Analyses of neurogenesis in *Hes1* type-1-mutant;*Hes3*(-/-);*Hes5*(-/-) mice. (A) Images of the whole body at E12.5 of *Hes3*(-/-);*Hes5*(-/-) and *Hes1* type-1 mutant;*Hes3*(-/-);*Hes5*(-/-) mice. (B,D) Size quantification (with SEM) of the whole body (B), telencephalon (B,D), Cortex (CX, D), and ganglionic eminence (GE, D) of *Hes3*(-/-);*Hes5*(-/-) and *Hes1* type-1 mutant;*Hes3*(-/-);*Hes5*(-/-) mice at E12.5. (C,F) Immunohistochemistry of Nestin (C), Tuj1 (C), cleaved caspase-3 (F), and phospho-Histone H3 (pH3) (F) in the telencephalon and spinal cord of *Hes3*(-/-);*Hes5*(-/-) and *Hes1* type-1 mutant;*Hes3*(-/-);*Hes5*(-/-) mice at E12.5. (E,G) Quantification of the proportions (with SEM) of Tuj1⁺ (E) and the number of cells expressing cleaved caspase-3 (G) in the CX and GE of *Hes3*(-/-);*Hes5*(-/-) and *Hes1* type-1 mutant;*Hes3*(-/-);*Hes5*(-/-) mice at E12.5. n = 3 for each genotype. *p < 0.05, **p < 0.01, Student's *t*-test. Scale bars = 2 mm (A); 400 μm (C,F).

Figure 1.






Figure 2.




Figure 3.




Figure 4.






Figure 5.


A




B




C



D



E



F




Figure 6.





Figure7.



Supplemental Information




Figure S1. Mathematical simulation. We used a mathematical model described before (Hirata et al., 2004). (A) Oscillatory expression depends on delayed negative feedback. When the transcriptional delay (T_m) is 29 min, oscillation continues. (B,C) When T_m is shortened, oscillation quickly dampens and ceases. (D) When T_m is lengthened, oscillation slows but continues. See Materials and Methods for more details.




Figure S2. Measurement of transcriptional delays. (A) Schematic structures of the three reporter genes, wild-type (WT) *Hes1* (control), intronless *Hes1* (type-1 mutant), and *Hes1* with *Hes1* cDNA knocked-in into the first exon (type-2 mutant), each of which contained Luc2 cDNA fused in-frame at the 5' end of the coding region. (B,C) Each reporter was introduced into C2C12 myoblast cells (B) and NS cells (C), and *Hes1* promoter was activated with serum treatment at time = 0 to measure the expression kinetics of *Hes1* reporters.




Figure S3. Characterization of pHes1-Ub-luc. (A) Schematic structure of pHes1-Ub-luc. (B) pHes1-Ub-luc was introduced into NPCs with the mCherry expression vector. Luciferase activities (white) in NPC cultures carrying pHes1-Ub-luc (red) were first measured, and Hes1 immunostaining (green) was performed. Among NPCs carrying pHes1-Ub-luc (mCherry⁺), Hes1⁺ cells are negative for the luciferase activity (arrows), while Hes1⁻ cells are positive for the luciferase activity (arrowhead). (C) Quantification of luciferase activities and Hes1 protein levels in individual NPCs carrying pHes1-UbLuc. n = 97. These data indicate that the luciferase activity and the endogenous Hes1 protein expression exhibited an inverse correlation. Because Hes1 protein expression and Hes1 gene transcription also exhibit an inverse correlation due to negative feedback, these data suggest that the pHes1-Ub-luc reporter activity matches the endogenous Hes1 gene transcription.






Figure S4. *Hes1* expression dynamics in *Hes1* type-1 and type-2 mutant mice. Luciferase activities were monitored in NPCs carrying pHes1-Ub-luc of wild-type (A), *Hes1* type-1 mutant (B), and *Hes1* type-2 mutant (C) mice.

A



B



C




Figure S5. The expression patterns of the *Hes1* type-2 mutant mic. (A) *In situ* hybridization analyses of the telencephalon of *Hes1* type-2-mutant mice. The expression patterns of the *Hes1* type-2 mutant mice were not significantly different from those of the WT from E10.5 to E16.5. (B,C) Real-time RT-PCR analyses of gene expression in WT and *Hes1* type-2 mutant mice. Values with SEM relative to the WT are shown. n = 3 for each genotype. *p < 0.05, Student's *t*-test.




Figure S6. Neurogenesis and cell cycle analyses of NPC cultures derived from wild-type and *Hes1* type-1 mutant embryos. (A,C) Staining of Tuj1, Ki67, BrdU, and DAPI. (B) Quantification of Tuj1⁺ cell number (with SEM). (D) Proportions (with SEM) of cell cycle exit. (E,F) Quantification of the number (E) and size (F) of neurospheres. Average values with SEM are shown. Scale bars: 100 μ m.




Figure S7. Hes1 expression dynamics in *Hes1*(+/-) NPCs. (A-E) Luciferase activities were monitored in NPCs carrying pHes1-Ub-luc of wild-type (WT) and *Hes1*(+/-) NPCs. The period (A), period distribution (B), relative amplitude (C), and amplitude/mean (D) were quantified, based on traces of *Hes1* oscillations in *Hes1*(+/-) NPCs (E). (F) Western blot analysis of Hes1 protein expression in WT and *Hes1*(+/-) NPCs. Average values with SEM are shown. n = 3.




Figure S8. Analyses of neural development of *Hes1*(+/-) and *Hes1*(+/-);*Hes5*(-/-) embryos. (A,C) Images of whole body (A) at E14.5 and anterior portions (C) at E16.5 of WT, *Hes1*(+/-), *Hes1*(-/-), and *Hes1*(+/-);*Hes5*(-/-) embryos. (B,D) Quantification of

the size (with SEM) of whole body at E14.5 (B) and at E16.5 (D). (E) Real-time RT-PCR analyses of Notch-related gene expression in WT and *Hes1*(+/-) embryos at E12.5 and E14.5. n = 3 for each sample. (F-H) Immunostaining of Tbr2 (F), Tbr1 (G), and Ctip2 (H) in the cortex of *Hes5*(-/-) and *Hes1*(+/-);*Hes5*(-/-) embryos at E10.5 and E12.5. (I) Real-time RT-PCR analyses of *Tbr2*, *Tbr1*, and *Ctip2* in the brains of *Hes5*(-/-) and *Hes1*(+/-);*Hes5*(-/-) embryos at E10.5 and E12.5. At least 3 samples were examined for each genotype. (J) Immunohistochemistry of phospho-Histone H3 (pH3) and cleaved caspase3 (casp3) in the cortex of *Hes5*(-/-) and *Hes1*(+/-);*Hes5*(-/-) embryos at E10.5 and E12.5. (K-N) Quantification of apical (K) and basal mitotic cells (L), total number of mitotic cells (M), and apoptotic cells (N) in the cortex of *Hes5*(-/-) and *Hes1*(+/-);*Hes5*(-/-) embryos at E10.5 and E12.5. Average values with SEM are shown. At least 3 samples were examined for each genotype. *p < 0.05, ***p < 0.001, Student's *t*-test. Scale bars: 4 mm (A,C); 100 μ m (F-H,J).

Antigen	Host	Manufacturers, references	Antigen Retrieval
Hes1	Rabbit	Kobayashi et al. 2009	○
Nestin	Mouse	BD Biosciences	
Tuj1	Rabbit	Covance	
Pax6	Mouse	Abcam	○
Tbr2	Rabbit	Abcam	○
Tbr1	Rabbit	MERCK	○
Ctip2	Rat	Abcam	
Cux1	Rabbit	Santa Cruz Biotechnology	○
phosphorylated histone H3	Mouse	MERCK	
cleaved-caspase 3	Rabbit	CST	
GAD65	Mouse	Chemicon	
GABA	Rabbit	Sigma	

Table S1. Antibody list.

Gene	Forward primer	Reverse primer
Hes1	TGAAGGATTCCAAAATAAAATTCT CTGGG	CGCCTCTTCTCCATGATAGGCTTTGA TGAC
Hes1 F	TAAAAAGTTACTTTTTGTAGAGAGC	
Hes1 R1		AGCCTTCACTCTTTTATTATATTTTC TC
Hes1 R2		AGGAATTTTCTCCATTATATCAGC
Hes3	CCCTGCTTAGCACTGCTGAGA	CAGGGCTCAGAAGGCACTAAA
Hes5	AAGTACCGTGGCGGTGGAGATGC	CGCTGGAAGTGGTAAAGCAGCTT
Dll1	TCAGATAACCCTGACGGAGGC	AGGTAAGAGTTGCCGAGGTCC
Neurog2	TCGCCAGGGACTGTATCT	CTGTGAAGTGGAGTCCG
Ascl1	GCCACCAGAATGACTTCAGCAC	AAGGCAACCTATGGGAACCAAC

GAPDH	TGGGTGTGAACCACGA	AAGTTGTCATGGATGACCTT
-------	------------------	----------------------

Table S2. Primer sequences.