

Absolute embeddings in Hausdorff spaces

筑波大学大学院数理物質科学研究科 川口 慎二 (Shinji Kawaguchi)

Graduate School of Pure and Applied Sciences, University of Tsukuba

Arhangel'skiĭ and Tartir [3] characterized compactness by some relative separation property and posed the following problem; *characterize Tychonoff spaces X , for which there is a Tychonoff space Y containing disjoint closed copies X_1 and X_2 of X such that these copies cannot be separated in Y by open subsets*. Answering this question, Bella and Yaschenko [4] proved the following theorem. We note that this theorem also follows from Matveev, Pavlov and Tartir [6, Theorem 2.3].

Theorem 1 (Bella-Yaschenko [4]; see also [6]). *For a Tychonoff space X , the following conditions are equivalent.*

- (a) X is Lindelöf.
- (b) *If a Tychonoff space Y contains two disjoint closed copies X_1 and X_2 of X , then these copies can be separated in Y by open subsets.*

As another type of absolute embeddings, Bella and Yaschenko [4] also obtained the following characterization of absolute weak C -embeddings; recall that a subspace Y of a space X is *weakly C -embedded* in X if every continuous real-valued function f on Y has an extension over X which is continuous at every point of Y ([1]). A Tychonoff space X is *almost compact* if $|\beta X \setminus X| \leq 1$, where βX denotes the Stone-Čech compactification of X .

Theorem 2 (Bella-Yaschenko [4]). *A Tychonoff space X is weakly C -embedded in every larger Tychonoff space if and only if X is almost compact or Lindelöf.*

Concerning Theorem 2, Arhangel'skiĭ [2] posed the following problem; *when is a Hausdorff (Tychonoff) space Y weakly C -embedded in every larger Hausdorff space X ?* Yamazaki [9] answered this problem as follows.

Theorem 3 (Yamazaki [9]). *A Hausdorff space X is weakly C -embedded in every larger Hausdorff space if and only if either X is compact or every continuous real-valued function on X is constant.*

In view of these results, it is natural to consider a characterization of spaces X satisfying the condition (b) of Theorem 1 in the realm of Hausdorff spaces. We give a characterization of such spaces as follows.

Theorem 4. *For a Hausdorff space X , the following conditions are equivalent.*

- (a) X is compact.
- (b) *If a Hausdorff space Y contains two disjoint closed copies X_1 and X_2 of X , then these copies can be separated in Y by open subsets.*

For the detail of the proof, see [5].

Remark 5. Using [6, Theroem 2.3], we obtain the regular case of Theorem 1 as follows; *for a regular space X , the following conditions are equivalent.*

- (a) X is Lindelöf.
- (b) *If a regular space Y contains two disjoint closed copies X_1 and X_2 of X , then these copies can be separated in Y by open subsets.*

Remark 6. Yajima [7] proved that the following condition (c) is equivalent to the conditions (a) and (b) in Theorem 1; (c) *For every compactification αX of X , any two disjoint closed copies of X in $(X \times \alpha X) \cup (\alpha X \times X)$ are completely separated in it.*

Remark 7. It was proved in [8]; *for a Tychonoff space X , the following conditions are equivalent.*

- (a) X is compact.
- (b) *If a Tychonoff space Y contains two disjoint closed copies X_1 and X_2 of X , then these copies can be completely separated in Y .*

How about the corresponding case of regular (Hausdorff) spaces? Indeed, for a non-empty regular (respectively, Hausdorff) space X , we can construct a regular (respectively, Hausdorff) space Y contains two disjoint closed copies X_1 and X_2 of X such that these copies cannot be completely separated in Y ([5]).

References

- [1] A.V. Arhangel'skiĭ, *Relative topological properties and relative topological spaces*, Topology Appl., **70** (1996), 87–99.
- [2] A.V. Arhangel'skiĭ, *From classic topological invariants to relative topological properties*, Sci. Math. Japon., **55** (2002), 153–201.
- [3] A.V. Arhangel'skiĭ and J. Tartir, *A characterization of compactness by a relative separation property*, Questions Answers Gen. Topology, **14** (1996), 49–52.

- [4] A. Bella and I.V. Yaschenko, *Lindelöf property and absolute embeddings*, Proc. Amer. Math. Soc., **127** (1999), 907–913.
- [5] S. Kawaguchi, *Compactness and absolute embeddings in Hausdorff spaces*, to appear in Questions Answers Gen. Topology.
- [6] M.V. Matveev, O.I. Pavlov and J. Tartir, *On relatively normal spaces, relatively regular spaces, and on relative property (a)*, Topology Appl., **93** (1999), 121–129.
- [7] Y. Yajima, *Characterizations of paracompactness and Lindelöfness by the separation property*, Proc. Amer. Math. Soc., **131** (2002), 1297–1302.
- [8] K. Yamazaki, *A proof for the Blair-Hager-Johnson theorem on absolute z -embedding*, Comm. Math. Univ. Carolinae, **43** (2002), 175–179.
- [9] K. Yamazaki, *Absolute weak C -embedding in Hausdorff spaces*, Topology Appl., **131** (2003), 273–279.

Doctoral Program in Mathematics,
Graduate School of Pure and Applied Sciences,
University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan
E-mail: shinji@math.tsukuba.ac.jp