

To BCI from Subtractive Algebra

井関 清志 (Kiyoshi ISEKI)

After introducing the cocepts BCK, BCI(1966), many mathematicians have tried the generalizations of both systems. The final one(1994) is due to A.Ursini(Univ. of Siena). It is a **subtractive algebra**[32]. This is an algebra with a constant 0 and a binary operation $*$ satisfying the following two axioms:

$$x * x = 0. \quad x * 0 = x.$$

In our discussion, there is another important identity, which is very often used. Its identity is

$$0 * x = 0.$$

On the other hand, in BCI, the following facts hold.

1. $x * 0 = x, \quad x * x = 0,$
2. $x \leq y \Rightarrow z * y \leq z * x, \quad x * z \leq y * z,$
3. $(x * y) * z = (x * z) * y,$ (permutation rule)
4. $(0 * x) * (0 * y) = 0 * (x * y),$
5. The minimal element of a branch with x is given by $0 * (0 * x).$
6. If x is a minimal element, then $0 * (0 * x) = x.$
7. $x * y = y * x = 0 \Rightarrow x = y.$ (quasi-identity)
8. $0 \leq x$ for all x in BCK, but not in BCI.

We have the atlas of proper BCI of order $n \leq 5$ (J.Hao[8], B.Hu[10]). J.Meng, Y.B.Jun and E.H.Roh[21] gave all proper BCI of order 6.

From now, we omit $*$, so $x * y$ is denoted by xy

The first generalization **BCH** of both systems is due to Q.Hu. BCH is defined by the following conditions.

- (1) $xx = 0$,
- (2) $(xy)z = (xz)y$,
- (3) $xy = yx = 0 \Rightarrow x = y$.

The class of BCH does not make a variety, because it has the quasi-identity (3). The exact proof is not simple. Many useful and important properties of BCH are published in his book [11, 369-456]. Q,Hu pointed out that proper BCH of order ≤ 3 do not exist. J.Hao[9] proved that there exist exactly five proper BCH of order 4.

Remark In this paper, the class of each system with the quasi-identity (3) is never a variety. Moreover, many identities appear in this paper. An ordinary identity has same letters(variables) in both sides. For example, see the identity (2). On the other hand, some variables in one side disappear in another side. For example identity (1). A.Kolmogorov pointed out the importance of the later case in his course of algorithm(Yu.M.Movsisyan [22]). For example, we have Kolmogorov identity $(xy)(yz) = xz$ in semigroup theory.

Y.Komori(1984) introduced the concept of **BCC-algebra**:

1. $0x = 0$,
2. $x0 = x$,
3. $((xy)(zy))(xz) = 0$.
4. $xy = yx = 0 \Rightarrow x = y$.

This syestem was extensively studied by W.A.Dudek([3],[4],[5] and [6]).

Remark $((xy)(xz))(zy) = 0$ and $x0 = x \Rightarrow (x(xy))y = 0$.

From the first identity, $((x0)(xz))(z0) = 0$, Hence $x0 = x \Rightarrow (x(xz))z = 0$. The identity obtained is one of the axioms of BCK, BCI.

Y.B.Jun, E,H,Roh and H.S.Kim (1998) introduced the concept of **BH-algebra**([17],[32]):

1. $xx = 0$,
2. $x0 = x$,
3. $xy = yx = 0 \Rightarrow x = y$.

J.Neggers and H.S.Kim(2001) gave interesting examples of a BH-algebra. Let $\mathbb{R}$ be the set of real numbers.

$$xy = \begin{cases} 0, & \text{if } x = 0, \\ \frac{(x-y)^2}{x}, & \text{otherwise.} \end{cases}$$

It is easy check that $(\mathbb{R},., 0)$ is a BH-algebra satisfying $0x = 0$, but not a BCH-algebra. We know an example of BH in which $0x = 0$ does not hold.

.Neggers and H.S.Kim(1999) gave an algebra which is called a **d-algebra**[25].

1. $xx = 0$,
2. $0x = 0$.
3. $xy = 0 = yx \Rightarrow x = y$.

The system includes a quasi-identity.

There is a simple example of a d-algebra. Let X be the set of non-negative real numbers. Define

$$xy = \max\{0, x - y\}.$$

Then X is a d-algebra with $x0 = x$.

Remark If a d-algebra is associative, it is trivial. $0 = 0x = (xx)x = x(xx) = x0$ Hence $x = 0$. So the d-algebra is trivial. J.Negger and H.S.Kim [25].

They(2002) introduce a new concept of **B-algebra**[30].

The axioms are

1. $xx = 0$,
2. $x0 = x$,
3. $(xy)z = x(z(0y))$.

There is an interesting example of B-algebra. Let X be the set of all real numbers except for nonnegative $-n$, Define xy by

$$xy = \frac{n(x - y)}{n + y}$$

Then X is a B-algebra. On the other hand, define $xy = xy^{-1}$ in a given group G . Then G is a B-algebra. M.Kondo and Y.B.Jun[21] proved that the class of B-algebra is same with the class of groups.

However, J.Neggers and H.S.Kim work is never worthless(For example, A,Walendziak [34]).

C.B.Kim and H.S.Kim defined **BM algebra** with a constant 0 and a binary operation $*$ satisfying

1. $00 = 0$,
2. $x0 = x$,
3. $(zx)(zy) = yx$.

Coxeter algebra is introduced by H,S, Kim, Y.H.Kim and J.Neggers(2004)

1. $xx = 0$,
2. $x0 = x$,
3. $(xy)z = (xz)y$.

K.Iseki, J.Neggers and H.S.Kim defined **J-algebra**.

1. $x * 0 = x$,
2. $x * (x * (y * (y * x))) = y * (y * (x * (x * y)))$.

Remark. In set theory, there is an identity:

$$A - (A - (B - (B - C))) = A \cap B \cap C.$$

Axiom 2 is a special case of the above identity.

S.M.Hong, Y.B.Jun and M.Aliöztürk(2003) defined an algebra which is called **gBCK-algebra**.

1. $x0 = x$,
2. $xx = 0$,
3. $(xy)z = (xz)y$,
4. $(xy)z = (xz)(yz)$,

A positive implicative BCK is gBCK-algebra, The following relations are true:

- a) $0x = 0$,
- b) $(xy)x = 0$,
- c) $xy = 0 \Rightarrow (xz)(yz) = 0$.

Proof,

$$0 = 00 = (xx)(xx) = (xx)x = 0x.$$

$$(xy)x = (xx)y = 0y = 0.$$

$$xy = 0 \Rightarrow (xz)(yz) = (xy)z = 0z = 0.$$

In 2005, Y.B.Jun and H.S.Kim introduced the concept of a **subtraction algebra**. This is defined by the following axioms:

- 1. $x(yx) = x$,
- 2. $x(xy) = y(yx)$,
- 3. $(xy)z = (xz)y$.

The algebra does not include a constant 0. However, xx acts as a constant 0.

I introduced two systems in our early works. One of them is **I-algebra** which is formulated as follows:

- 1. $x0 = x$,
- 2. $0x = 0$,
- 3. $((xy)(xz))(zy) = 0$,
- 4. $(xy)(x(yx)) = 0$,
- 5. $xy = yx = 0 \Rightarrow x = y$.

Until now, I mentioned algebras and systems with only one binary operation. On the hand, various algebras and systems with several binary operations are in the literature.

I introduced a system in our researches(1967). It is **Griss-algebra**. The system has two operations $*$ and $+$. A constant 0 is included in Griss system. $x \leq y$ means $xy = 0$. Griss system is formulated as follows:

1. $0x = 0$,
2. $x + x = x$,
3. $x; = y + x$,
4. $(x + y)(x + z) \leq zy$,
5. $x \leq x + y$,
6. $xy = yx = 0 \Rightarrow x = y$.

J.M.Neggers and H.S.Kim(2002) defined an algebra with a constant 0 and two binary operations satisfying the following three axioms.

1. $x0 = x$,
2. $(0x) + x = 0$,
3. $(xy)z = x(y + z)$.

The algebra is called β -**algebra**[32].

There are other important generalizations of BCK and BCI due to Bucarest research group(under G.Georgescu). They defined the concepts of pseudo-MV, pseudo-BL, pseudo-BCK, Iseki algebra, and pseudo-Iseki algebras and a pseudo t-norm. A.Iorgulescu(2003) defined many systems([12],[13]), for example, left-Iseki(RP)

algebra,..., left Hájek(R)-algebra. Among them, I mention only definition of pseudo-BCK is given. A **pseudo-BCK** is a system $\mathbf{X} = (X, \leq, *, \diamond, 0)$, where $\leq$ is a binary relation, $*$, $\diamond$ are binary operations, and 0 is a constant, and is given by the following axiom system:

1. $(x * y) \diamond (x * z) \leq z * y, (x \diamond y) * (x \diamond z) \leq z \diamond y,$
2. $x * (x \diamond y) \leq y, x \diamond (x * y) \leq y,$
3. $x \leq x,$
4. $0 \leq x.$
5. $x \leq y \wedge y \leq x \Rightarrow x = y,$
6. $x \leq y \Leftrightarrow x * y = 0 \Leftrightarrow x \diamond y = 0.$

References

- [1] M.Abe and K.Iseki, A survey of BCK and BCI algebras, Congresso de logica Aplicada a Tecnologia, LAPTEC 2000. 431-443.
- [2] J.Allen, J.Eggers and H.S.Kim, L-up and mirror algebras, Math. Japonica, 58(2004),381-388.
- [3] W.A.Dudek, The number of subalgebras of finite BCC-algebras, Bll. inst. Math. Academia Sinica. 20(1992), 129-136.
- [4] W.A.Dudek, X.H.Zhang, On ideals and congruences in BCC-algebras, Czech. Math. J.48(123)(1998), 21-29.
- [5] W,A,Dudek and X.H.Zhang, On atoms in BCC-algebras, Diss. Math. 15(1995), 81-85.

- [6] W.A.Dudek, K.H.Kim and Y.B.Jun, Fuzzy BCC-subalgebras of BCC-algebras with respect to a t-norm, *Sci. Math.* 4(2000), 99-106.
- [7] T.Y.Guo, The structure of a class of BCI-algebras, *Math. Japonica*, 38(1993), 703-711.
- [8] J.Hao, Atlas of proper BCI-algebras of order $n \leq 5$, *Math. Japonica*, 38(1993), 589-591.
- [9] J.Hao, Complete classification of proper BCH-algebras of order four, *Sci Marg.*, 2(1999), 205-213.
- [10] B.Hu, Extension by one point and quasi-commutativity of BCH-algebra (in Chinese), *J. of Wuhan Univ.* 24(1991), 508-516.
- [11] Q.Hu, BCI-algebra (in Chinese), Shanxi Science-Technique Publishing House, Xian, 1987.
- [12] A.Iorgulescu, Some direct ascendants of Warfield and MV algebras, *Math. Japonica*, 57(2003), 583-647.
- [13] A.Iorgulescu, On pseudo-BCK algebra and PORMs, *Math. Japonica*, 60(2004), 293-305.
- [14] K.Iseki, A way to BCK and related systems, *Math. Japonica*, 52(2000), 163-179.
- [15] K.Iseki, Some fundamental theorems on BCK, Words, Semigroups & Transductions, *Festschrift in Honor of Gabriel Therrin*, World Sci.(2001), 231-238.
- [16] K.Iseki, A way to BCI, *Annales Societatis Math. Polonae, Seria I, Commentationes Math., Tomus Specialis in Honorem Julian Musielak.*(2004), 97-103

- [17] Y.B.Jun, H.R.Roh and H.S.Kim, On BH-algebras, *Sci. Math.*, 1(1998), 347-354.
- [18] Y.B.Jun, Characterization of pseudo-BCK algebras, *Math. Japonica*, 57(2003), 225-230.
- [19] K.H.Kim, On subtraction semigroups, *Sci. Math. Japonicae*, 62(2005), 273-280.
- [20] Y.Komori, The class of BCC is not a variety, *Math. Japonica*, 29(1984), 391-394.
- [21] M.Kondo and Y.B.Jun, The class of B-algebras coincides with the class of groups, *Math. Japonica*, 57(2003), 175-177.
- [22] J.Meng, Y.B.Jun and E.H.Roh, BCI-algebras of order 6, *Math. Japonica*, 47(1998), 33-43.
- [23] Yu.M.Movsisyan, Introduction to the theory of algebras with hyperidentities (in Russian), Erevan Univ. Press, 1986.
- [24] Yu.M.Movsisyan, Hyperidentities and hypervarieties in algebras (in Russian), Erevan Univ. Press, 1990.
- [25] J.Neggers and H.S.Kim, On d-algebras, *Math. Slovaca* 49(1999)19-26.
- [26] J.Neggers and H.S.Kim, On analytic T-algebras, *Math. Japonica*, 53(2001), 157-163.
- [27] B.Schein, Difference semigroups, *Comm. in Algebra* 20(1992), 2153-2169.
- [28] J.Neggers and H.S.Kim, A fundamental theorem of B-homomorphism for B-algebras, *Int. Math. Jour*, 2(2002), 207-214.

- [29] J.Neggers, Y.B.Jun and H.S.Kim, On d-ideals in d-algebras, Math. Slovaca, 49(1999), 243-251.
- [30] J.Neggers and H.Kim, On B-algebras, Mat. Vesnik 54(2002), 21-29.
- [31] J.Neggers and H.S.Kim, On β -algebras, Math. Slovaca, 52(2002), 517-530.
- [32] J.Neggers and H.S.Kim, On analytic T-algebras, Math. Japonica, 53(2001), 157-163.
- [33] A.Ursini, On subtractive varieties, I - IV, Alg. Universalis, 31(1994), 204-222, 36(1996), 222-259, 37(1997), 296-333. 38(1997), 355-389.6 [34] A.Walendziak, A Note on normal subalgebras in B-algebras, Sci. Math. Japonicae, 62(2005), 49-53.