

Monoids with finite (regular) complete presentation

東邦大理 小林美治 (Yuji Kobayashi)

京都産業大理 勝良昌司 (Masashi Katura)

If a monoid is defined by a finite complete rewriting system, then it has some good properties such as solvability of the word problem (see [1]). So, it is an important problem to find good conditions for monoids to have finite complete presentation.

A rewriting system over a (finite) alphabet Σ is a subset R of $\Sigma^* \times \Sigma^*$. An element (u, v) of R is called a rule and written $u \rightarrow v$. A word $x \in \Sigma^*$ is rewritten to $y \in \Sigma^*$ under R , if $x = x_1 u x_2$, $y = x_1 v x_2$ for some $x_1, x_2 \in \Sigma^*$ and some $u \rightarrow v \in R$. In this case we write $x \rightarrow_R y$. The reflexive transitive closure and the reflexive symmetric transitive closure of the relation $\rightarrow_R$ is denoted by $\rightarrow_R^*$ and $\leftrightarrow_R^*$, respectively. The quotient monoid $M = \Sigma^* / \leftrightarrow_R^*$ is a monoid presented by (Σ, R) .

A subset S of Σ^* is called s-closed, if any subword of an element of S is also in S , equivalently, if the complement of S forms an ideal of Σ^* . S is s-closed if and only if S is expressed as

$$S = \Sigma^* - \Sigma^* T \Sigma^* \quad (1)$$

with a subset T of Σ^* . A set S is finitely s-closed if S is expressed as (1) with a finite set T , or equivalently, S is the complement of a finitely generated ideal.

Proposition 1. For an s-closed subset S of Σ^* the following are equivalent.

- (1) S is finitely s-closed.
- (2) There is a positive n such that any subword of length $\leq n$ of $x \in \Sigma^*$ is in S , then x is in S .

Corollary. A finite s-closed set is finitely s-closed.

Let $R \subset \Sigma^* \times \Sigma^*$ and let $M = M(\Sigma, R)$ be the monoid presented by (Σ, R) . A subset S of Σ^* is a transversal for M (or for (Σ, R)), if S forms a complete set of representatives for M , that is, for any $x \in \Sigma^*$ there is a unique $\hat{x} \in S$ with $x \leftrightarrow^* \hat{x}$.

M has a complete presentation, if M is presented by (Σ, R) such that R is a complete (i.e. noetherian and confluent) rewriting system.

Proposition 2. If a monoid has a finite complete presentation, then it has a finitely s-closed transversal.

We are interested in the converse problem of this results.

Problem 1. If a monoid has a finitely s-closed transversal, does it admit a finite complete presentation ?

Proposition 3. Let $S = \Sigma^* - \Sigma^* T \Sigma^*$ be an s-closed transversal for M , and let $R(T) = \{u \rightarrow \hat{u} \mid u \in T\}$. If $R(T)$ is noetherian, then R is a complete system defining M .

Unfortunately, the system $R(T)$ given above is not always noetherian even when S is finitely s-closed and T is chosen minimally.

Example 1. Let $\Sigma = \{a, b\}$ and $M = M(\Sigma, E)$, where

$$E = \{(baab, aba)\} \cup \{(x, y) \in \Sigma^* \times \Sigma^* \mid |x| = |y| = 4\}.$$

Then M is a finite monoid with finite s-closed transversal

$$S = \{x \in \Sigma^* \mid |x| \leq 3, x \neq aba\} \cup \{baab\}.$$

A set T with which S is expressed as (1) must contain the rule

aba. Hence, $R = R(T)$ contains the rule $aba \rightarrow baab$. Thus

$$abaa \rightarrow_R baaba \rightarrow_R babaab \rightarrow_R bbaabab \rightarrow_R bbabaabb \rightarrow_R \dots,$$

and R is not noetherian.

If M has a transversal $S = \Sigma^* - \Sigma^* T^*$ with T a singleton, then we expect that the system $R(T)$ would be noetherian, and M would admit a complete one-rule system.

Problem 2. Suppose M has a transversal S of the form

$$S = \Sigma^* - \Sigma^* v \Sigma^*, \quad v \in \Sigma^*.$$

Is the one-rule system $\{v \rightarrow \hat{v}\}$ noetherian?

A system R is regular, if the left hand sides of the rules from R forms a regular set (see [2]).

Theorem 1. Suppose M is presented by a regular complete system (Σ, R) . Then,

- (1) M has a regular sets of representative.
- (2) $\text{Irr}(R)$ grows either exponentially or polynomially.
- (3) If $\text{Irr}(R)$ grows exponentially, then the monoid $M = M(\Sigma, R)$ also grows exponentially and contains a free submonoid of rank 2.

Problem 3. Does M have a regular complete presentation, provided M has a regular s-closed transversal?

A system (Σ, R) is polynomially mild, if there is a polynomial f such that $|y| \leq f(|x|)$ for any $x \in \Sigma^*$ and y such that $x \rightarrow^* y$.

Theorem 2. If (Σ, R) is a polynomially mild regular complete system such that $\text{Irr}(R)$ grows polynomially, then the monoid $M = M(\Sigma, R)$ grows polynomially.

In theorem 2, the mildness of R cannot be removed. In fact, M can grows exponentially in general, even if $\text{Irr}(R)$

grows polynomially.

Example 2. Let $\Sigma = \{a, b\}$ and $R_1 = \{abb \rightarrow ba\}$, $R_2 = \{ba \rightarrow abb\}$. Then, the both R_1 and R_2 are complete and define the same monoid. However, $\text{Irr}(R_1)$ grows exponentially, while $\text{Irr}(R_2)$ grows polynomially. So, the system R_2 is not mild.

There is a monoid with a very simple presentation which never admits a finite complete presentation.

Example 3. Let $\Sigma = \{a, b\}$ and $R = \{b^2a \rightarrow ab^2, ab^3 \rightarrow b^3, b^3a \rightarrow b^3\}$, then $M = M(\Sigma, R)$ has no finitely s -closed transversal. So, M has no finite complete presentation.

References

- [1] K. Madlener and F. Otto, About the descriptive power of certain classes of finite string-rewriting systems, Theoret. Comp. Sci. **67**, (1989), 143-172.
- [2] Y. Kobayashi, A finitely presented monoid which has solvable word problem but has no regular complete presentation, to appear in Theoret. Comp. Sci.