

Note on blocks of p -solvable groups with same Brauer category

熊本大学理学部 渡辺アツミ (Atumi Watanabe)
Department of Mathematics, Faculty of Science
Kumamoto University

1

Let p be a prime and let $\mathcal{O}$ be a complete discrete valuation ring with an algebraically closed residue field k of characteristic p . Let G be finite group and b be a block of G with maximal (G, b) -subpair (P, e_P) where b is a block idempotent of $\mathcal{O}G$. For any subgroup Q of P , let (Q, e_Q) be a unique (G, b) -subpair contained in (P, e_P) . Following Kessar, Linckelmann and Robinson [4], we denote by $\mathcal{F}_{(P, e_P)}(G, b)$ the category whose objects are subgroups of P and for $Q, R \leq P$, whose set of morphisms from Q to R are the set of group homomorphisms $\varphi : Q \rightarrow R$ such that there exists $x \in G$ such that ${}^x(Q, e_Q) \subseteq (R, e_R)$ and $\varphi(u) = xux^{-1}$ for all $u \in Q$. We call $\mathcal{F}_{(P, e_P)}(G, b)$ the Brauer category of b . Let $\mathbf{B}_G(b)$ be the Brauer category of b in the sense of Thévenaz [10], § 47. The categories $\mathcal{F}_{(P, e_P)}(G, b)$ and $\mathbf{B}_G(b)$ are equivalent. Let R be a normal subgroup of P such that $N_G(P) \subseteq N_G(R)$ and c be the Brauer correspondent of b in $N_G(R)$, that is, c is a unique block of $N_G(R)$ such that $\text{Br}_P(c) = \text{Br}_P(b)$ where Br_P is the Brauer homomorphism from $(\mathcal{O}G)^P$ onto $kC_G(P)$. Set $N = N_G(R)$. The notations R, c and N are fixed. Thus $b = c^G$ and (P, e_P) is a maximal (N, c) -subpair. The arguments in the proof of Theorem in Kessar-Linckelmann [5] imply the following.

Theorem 1 *Assume that G is p -solvable. With the above notations, suppose that $\mathcal{F}_{(P, e_P)}(G, b) = \mathcal{F}_{(P, e_P)}(N, c)$. Then there is an indecomposable $\mathcal{O}Gb$ - $\mathcal{O}Nc$ -bimodule M which satisfies the following.*

- (i) M and its $\mathcal{O}$ -dual M^* induce a Morita equivalence between $\mathcal{O}Gb$ and $\mathcal{O}Nc$.
- (ii) As an $\mathcal{O}(G \times N)$ -module M has a vertex ΔP and an endo-permutation $\mathcal{O}(\Delta P)$ -module as a source where $\Delta P = \{(u, u) \mid u \in P\}$.

Let $H_{(P, e_P)}^*(G, b)$ be the cohomology ring of b in the sense of Linckelmann [6], [7], that is, $H_{(P, e_P)}^*(G, b)$ is the subring of $H^*(P, k)$ consisting of $\zeta \in H^*(P, k)$ satisfying $\text{res}_Q \zeta = {}^g \text{res}_Q \zeta$ for all $Q \leq P$ and, for all $g \in N_G(Q, e_Q)$. We prove the following.

Theorem 2 *Assume that G is p -solvable. With the above notations, if $H_{(P, e_P)}^*(G, b) = H_{(P, e_P)}^*(N, c)$, then $\mathcal{F}_{(P, e_P)}(G, b) = \mathcal{F}_{(P, e_P)}(N, c)$.*

2

We prove Theorem 1 using the following.

Lemma 1 (Harris-Linckelmann [3], Lemma 4.2) *Assume that G is p -solvable. For any p -subgroup Q of G , we have $O_{p'}(N_G(Q)) = O_{p'}(G) \cap N_G(Q) = O_{p'}(G) \cap C_G(Q) = O_{p'}(C_G(Q))$.*

Proposition 1 (Harris-Linckelmann [2], Proposition 3.1 (iii)) *Let G be a p -solvable group and b be a block of G such that b covers a G -invariant block of $O_{p'}(G)$. Then b is of principal type, that is, for any p -subgroup Q of G , $\text{Br}_Q(b)$ is a block of $kC_G(Q)$.*

Proposition 2 (Fong[1]; Puig[9]) *Let G be a p -solvable group and b be a block of G with defect group P . Then the following holds.*

(i) *There is a subgroup H of G and an H -invariant block e of $O_{p'}(H)$ such that $O_{p'}(G)P \subseteq H$ and $\mathcal{O}Gb \cong \text{Ind}_H^G(\mathcal{O}He)$ as interior G -algebras.*

(ii) *P is a Sylow p -subgroup of H and P is a defect group of e as a block of H . Moreover let (P, e'_P) be a maximal (H, e) -subpair and let $e_P = \text{Tr}_{CH(P)}^{CG(P)}(e'_P)$. Then (P, e_P) is a maximal (G, b) -subpair.*

Note that in the above proposition $\mathcal{F}_{(P, e_P)}(G, b) = \mathcal{F}_{(P, e'_P)}(H, e)$ since $\mathcal{O}Gb \cong \text{Ind}_H^G(\mathcal{O}He)$ as interior G -algebras.

Proposition 3 ([5], Proposition 6) *With the notations in the above proposition, let R be a subgroup of P such that $N_G(P) \subseteq N_G(R)$. Denote by c the Brauer correspondent of b in $N_G(R)$, and by f the Brauer correspondent of e in $N_H(R)$. Then f is an $N_H(R)$ -invariant block of $O_{p'}(N_H(R))$ and $\mathcal{O}N_G(R)c \cong \text{Ind}_{N_H(R)}^{N_G(R)}(\mathcal{O}N_H(R)f)$ as interior $N_G(R)$ -algebras.*

The following is shown in the proof of Theorem in [5].

Theorem 3 (Kessar-Linckelmann) *Let G be a p -solvable group and b be a block of G with defect group P . Let R be a subgroup of P such that $N_G(P) \subseteq N_G(R)$ and let c be the Brauer correspondent of b in N where we set $N = N_G(R)$. If b covers a G -invariant block of $O_{p'}(G)$ and if $G = O_{p'}(G)N$, then there is an indecomposable $\mathcal{O}Gb$ - $\mathcal{O}Nc$ -bimodule M which satisfies the following.*

(i) *M and its $\mathcal{O}$ -dual M^* induce a Morita equivalence between $\mathcal{O}Gb$ and $\mathcal{O}Nc$.*

(ii) *As an $\mathcal{O}(G \times N)$ -module M has a vertex ΔP and an endo-permutation $\mathcal{O}(\Delta P)$ -module as a source.*

Proof of Theorem 1. We prove by induction on $|G|$. Let H , e , e'_P and e_P be as in Proposition 2, and let f be as in Proposition 3. We may assume that e_P 's in Theorem 1 and Proposition 2 are equal by replacing H , e , e'_P and f , by H^x , e^x , $(e'_P)^x$ and f^x respectively for some $x \in N_G(P)$ if necessary. By Proposition 2,

$$\mathcal{F}_{(P, e_P)}(G, b) = \mathcal{F}_{(P, e'_P)}(H, e).$$

By Proposition 3, (P, e'_P) is a maximal $(N_H(R), f)$ -subpair and

$$\mathcal{F}_{(P, e_P)}(N, c) = \mathcal{F}_{(P, e'_P)}(N_H(R), f).$$

So by the assumption we have $\mathcal{F}_{(P, e'_P)}(H, e) = \mathcal{F}_{(P, e'_P)}(N_H(R), f)$. Since $\mathcal{O}Gb \cong \text{Ind}_H^G(\mathcal{O}He)$ as interior G -algebras, the $\mathcal{O}Gb$ - $\mathcal{O}He$ -bimodule $b\mathcal{O}Ge = \mathcal{O}Ge$ and the $\mathcal{O}He$ - $\mathcal{O}Gb$ -bimodule $e\mathcal{O}G$ induce a Morita equivalence between $\mathcal{O}Gb$ and $\mathcal{O}He$. Similarly the $\mathcal{O}Nc$ - $\mathcal{O}N_H(R)f$ -bimodule $\mathcal{O}Nf$ and the $\mathcal{O}N_H(R)f$ - $\mathcal{O}Nc$ -bimodule $f\mathcal{O}N$ induce a Morita equivalence between $\mathcal{O}Nc$ and $\mathcal{O}N_H(R)f$. Suppose that $H < G$. By the induction hypothesis for H and e , there is an indecomposable $\mathcal{O}He$ - $\mathcal{O}N_H(R)f$ -bimodule M_0 such that M_0 and M_0^* induce a Morita equivalence between $\mathcal{O}He$ and $\mathcal{O}N_H(R)f$, and that M_0 as an $\mathcal{O}(H \times N_H(R))$ -module has a vertex ΔP and an endo-permutation $\mathcal{O}(\Delta P)$ -module as a source. Set $M = b\mathcal{O}G \otimes_{\mathcal{O}He} M_0 \otimes_{\mathcal{O}N_H(R)f} \mathcal{O}Nc \cong M_0^{G \times N}$. Then M satisfies (i) and (ii) in Theorem 1. Therefore we may assume that $H = G$. Then $b = e$.

Let $Y = O_{p', p}(G)$. Then b is a G -invariant block of Y because $Y/O_{p'}(G)$ is a p -group. Furthermore we have $Y = O_{p'}(G)(Y \cap P)$. Set $Q = P \cap Y$. Then Q is a defect group of b as a block of Y . Now since G is constrained, $C_Y(Q) = C_G(Q)$. Therefore we see that (Q, e_Q) is a maximal (Y, b) -subpair. By the Frattini argument and the assumption that $\mathcal{F}_{(P, e_P)}(G, b) = \mathcal{F}_{(P, e_P)}(N, c)$,

$$G = N_G(Q, e_Q)Y \subseteq N_N(Q)C_G(Q)Y \subseteq NY \subseteq NO_{p'}(G).$$

So we have $G = NO_{p'}(G)$. This and Theorem 3 complete the proof.

Proof of Theorem 2. We prove by induction on $|G|$. Let H , e , e'_P and e_P be as in Proposition 2, and let f be as in Proposition 3. We may assume that e_P 's in Theorem 2 and Proposition 2 are equal as in the proof of Theorem 1. Since $\mathcal{F}_{(P, e_P)}(G, b) = \mathcal{F}_{(P, e'_P)}(H, e)$ and $\mathcal{F}_{(P, e_P)}(N, c) = \mathcal{F}_{(P, e'_P)}(N_H(R), f)$ we have

$$H_{(P, e_P)}^*(G, b) = H_{(P, e'_P)}^*(H, e).$$

$$H_{(P, e_P)}^*(N, c) = H_{(P, e'_P)}^*(N_H(R), f).$$

From the assumption, we have $H_{(P, e'_P)}^*(H, e) = H_{(P, e'_P)}^*(N_H(R), f)$. Suppose that $H < G$. Then by the induction hypothesis, $\mathcal{F}_{(P, e'_P)}(H, e) = \mathcal{F}_{(P, e'_P)}(N_H(R), f)$, and hence $\mathcal{F}_{(P, e_P)}(G, b) = \mathcal{F}_{(P, e_P)}(N, c)$. Therefore we may assume that $H = G$. Then b covers a G -invariant block of $O_{p'}(G)$ and P is a Sylow p -subgroup of G . Note that the element $b \in \mathcal{O}O_{p'}(G)$.

From Proposition 1, b is of principal type. On the other hand, by Lemma 1, $\text{Br}_R(b)$ is an N -invariant block idempotent of $kO_{p'}(N)$ and c is a lifting of $\text{Br}_R(b)$ to $\mathcal{O}N$. So by Proposition 1, c is also of principal type. So we may assume that b is a principal block. Therefore by a theorem of Mislin [8], we obtain $\mathcal{F}_{(P, e_P)}(G, b) = \mathcal{F}_{(P, e_P)}(N, c)$. This completes the proof.

References

- [1] P. Fong, On the characters of p -solvable groups, Trans. Amer. Math. Soc. **98**(1961). 263-284.
- [2] M.E. Harris and Linckelmann, Splendid derived equivalences for blocks of finite p -solvable groups, J. London Math. Soc. (2) **62**(2000). 85-96.
- [3] M.E. Harris and Linckelmann, On the Glauberman and Watanabe correspondences for blocks of finite p -solvable groups. Trans. Amer. Math. Soc. **354**(2002). 3435-3453.
- [4] R. Kessar, M. Linckelmann and G.R. Robinson, Local control in fusion systems of p -blocks of finite groups, J. Algebra **257**(2002). 393-413.
- [5] R. Kessar and M. Linckelmann, On blocks of strongly p -solvable groups, D. Benson: Groups, Representations and Cohomology Preprint Archive.
- [6] M. Linckelmann, Transfer in Hochschild cohomology of blocks of finite groups. Algebr. Represent. Theory **2** (1999). 107-135.
- [7] M. Linckelmann, Varieties in block theory. J. Algebra **215**(1999), 460-480.
- [8] G. Mislin, On group homomorphisms inducing mod p -cohomology isomorphism. Comment. Math. Helv. **65**(1990). 454-461.
- [9] L. Puig, Local block theory in p -solvable groups. Proceedings of Symp. Pure Math. **37**(1980). 385-388.
- [10] J. Thévenaz, " G -algebras and modular representation theory", Oxford Sci. Publ., Clarendon Press, Oxford, 1955.