

學位申請論文



白井三平

論 文 内 容 の 要 旨

報 告 番 号	甲 第 号	氏 名	白 井 三 平
論文調査担当者	主査 永 田 雅 宜 土方 弘 明 ・ 戸 田 宏		
(論 文 題 目) Local Torelli theorem for some non-singular weighted complete intersections. (ある種の非特異 weighted complete intersections に対する局所 Torelli 定理)			
(論文内容の要旨) P. A. Griffiths は Periods of integrals on algebraic manifolds と題する論文において, local Torelli theorem についての結果を整理し, 解明すべき問題を提起した。 すなわち, コンパクト ケーラー多様体について, その deformations のパラメーター空間で定義された period map なるものを考え, それが infinitesimal deformations をうまく分離するかどうかということが local Torelli theorem の成否で判定されるという問題設定を行ない, 従って, どのような場合にそれが成立するかという形の問題提起をし, 特に, 単連結な代数曲面で canonical bundle が ample である場合についてはどうかと問うた。			

申請者はこの問題について、まず参考論文(その2)において、 d 次元($d \geq 2$)の non-singular complete intersection であるような多様体(基礎体の標数0)で canonical bundle が ample であるものについて、local Torelli theorem が成立するという大変すぐれた結果を与えた。

さらに、主論文において、weighted complete intersection の場合に、若干の条件の下でやはり local Torelli theorem が成り立つことを示し、また他方、証明の適用と、仮定した条件との関連を、例を挙げて吟味し、今後の更なる発展のための資料としている。

1976, 5月

主論文

LOCAL TORELLI THEOREM FOR SOME NON-SINGULAR
WEIGHTED COMPLETE INTERSECTIONS

by SAMPEI USUI

Introduction.

For a compact Kähler manifold X , P. A. Griffiths constructed and studied a "period map" from the parameter space of deformations of X to the "Griffiths domain" which parametrizes the Hodge structures on the underlying differentiable manifold of X ([2]). As one of these studies, he formulated the local Torelli problem in the form (P_1) in §1, which asks whether a period map separates infinitely near points (cf. [2] and also [7]). After examining this problem in some examples, Griffiths presented in his article [3] a problem :

"(7.1). PROBLEM*. Find method to treat the local Torelli theorem. In particular, decide whether it is true or false for simply-connected canonical surfaces (i.e. surfaces with ample canonical bundle)."

The purpose of the present paper is to give a proof of the local Torelli theorem for some non-singular "weighted complete intersections" (see Theorem 2.1 below).

In the author's previous paper [7], he gave a proof of the local Torelli theorem for non-singular complete intersections with ample canonical line bundle. On the other hand, Sigefumi Mori introduced and studied a family of varieties so called "weighted complete intersections" ([5]). Weighted complete

intersections have several properties ^{similar} to those of complete intersections (cf. [5] and also §1), so that our method in [7] is naturally applicable for some weighted complete intersections. However, we need some assumptions (the assumptions i) and ii) in Theorem 2.1) in order to let our method work well.

§3 includes the discussions on these assumptions and some examples for which our method breaks down.

The author wishes to thank Professor S. Mori for the valuable suggestions.

§1. Preliminaries.

a. Local Torelli problem. In this subsection, we recall the formulation of the local Torelli problem (for the background of this problem see [2] and also [7]).

Let X be a compact ["]Kähler manifold. The local Torelli problem is the question whether the period map of Griffiths separates infinitely near points at the point in the parameter space of the Kuranishi family of deformations of X , corresponding to X .

The local Torelli problem in r -th cohomology is formulated as follows.

(P₁) Let $H^1(X, T_X) \otimes H^{r-i+1, i-1}(X) \longrightarrow H^{r-i, i}(X)$ be the pairing induced from $T_X \otimes \Omega_X^{r-i+1} \longrightarrow \Omega_X^{r-i}$ ($1 \leq i \leq r$). Is $\tau \in H^1(X, T_X)$ zero if $\tau \cdot \alpha_i$ is zero in $H^{r-i, i}(X)$ for every $\alpha_i \in H^{r-i+1, i-1}(X)$ ($1 \leq i \leq r$)?

b. Weighted complete intersections. S. Mori generalized the notion of complete intersection and studied so called "weighted complete intersections" ([5]).

The following definitions and results are found in [5].

Definition 1.1. Let $n, e_0, \dots, e_n$ be positive integers. We set

$$m = \text{l.c.m.} \{e_0, \dots, e_n\}, \text{ and}$$

$$r(e) = r(e_0, \dots, e_n) = \min_{p: \text{prime}} \# \{1 \mid 0 \leq i \leq n, p \nmid e_i\}.$$

Let K be a field.

$Q(e_0, \dots, e_n)$, or simply $Q(e)$ is the scheme $\text{Proj}(K[X_0, \dots, X_n])$, where

the gradation of $K[X_0, \dots, X_n]$ is defined as follows;

$$\deg X_i = e_i \quad (0 \leq i \leq n), \text{ and } \deg a = 0 \quad (a \in K).$$

For an integer a , $O_{Q(e)}(a)$, or simply $O_Q(a)$ is the coherent $O_{Q(e)}$ -module corresponding to the homogeneous $K[X_0, \dots, X_n]$ -module $K[X_0, \dots, X_n](a)$.

For a positive integer h , S_h is the closed subset of $Q(e)$, defined by the ideal generated by $\{x_i \mid h \nmid e_i\}$.

It is easy to see that the singular locus of $Q(e)$ is $\bigcup_{1 \leq h} S_h$.

Definition 1.2. With the notations in Definition 1.1, $P(e_0, \dots, e_n)$, or simply $P(e)$ is the open subscheme $Q(e) - \bigcup_{1 \leq h} S_h$ of $Q(e)$. We call the scheme $P(e)$ a weak projective space of size $(e_0, \dots, e_n)$, or simply of size (e) . We define $\mathcal{O}_P(a) = \mathcal{O}_Q(a)|_{P(e)}$ for every integer a .

Definition 1.3. With the notations in Definitions 1.1 and 1.2, let $c, a_1, \dots, a_c$ be positive integers. We consider $\text{Proj}(K[X_0, \dots, X_n]/(F_1, \dots, F_c))$ for homogeneous elements $F_1, \dots, F_c$ in the graded ring $K[X_0, \dots, X_n]$ with $\deg F_j = a_j$ ($1 \leq j \leq c$), satisfying the following conditions.

- i) $F_1, \dots, F_c$ form a regular sequence of $K[X_0, \dots, X_n]$.
- ii) $V_+(F_1, \dots, F_c) \cap \bigcup_{1 \leq h} S_h = \emptyset$.

An algebraic K -scheme X is called a weighted complete intersection in $P(e)$ of type $(a_1, \dots, a_c)$, if X is isomorphic to such a K -scheme

$\text{Proj}(K[X_0, \dots, X_n]/(F_1, \dots, F_c))$.

In this case, for an arbitrary integer a , we denote by $O_X(a)$ the O_X -module induced from $O_P(a)$.

Results. (1) We see that $\text{codim}_{Q(e)} \bigcup_{1 \leq h} S_h = r(e)$, and the maximal dimension of complete subschemes of $P(e)$ is given by $r(e) - 1$.

(2) $O_P(1)$ is an invertible sheaf on $P(e)$, and we have a natural isomorphism $O_P(1)^{\otimes a} \xrightarrow{\sim} O_P(a)$ for every integer a .

(3) For a weighted complete intersection X , $O_X(1)$ is an ample invertible sheaf. Moreover, if $\dim X \geq 3$, then we have $\text{Pic } X \approx \mathbb{Z}$ and $O_X(1)$ generates $\text{Pic } X$.

(4) Let $X = \text{Proj}(K[X_0, \dots, X_n]/(F_1, \dots, F_c))$ be a weighted complete intersection of dimension ≥ 1 in $P(e)$ of type $(a_1, \dots, a_c)$. Then we have

$$(K[X_0, \dots, X_n]/(F_1, \dots, F_c))_a \xrightarrow{\sim} H^0(X, O_X(a)) \quad (a \in \mathbb{Z}),$$

$$H^j(X, O_X(a)) = 0 \quad (a, j \in \mathbb{Z}, \quad 0 < j < n - c = \dim X), \quad \text{and}$$

$$\omega_X = O_X\left(\sum_{j=1}^c a_j - \sum_{i=0}^n e_i\right) \quad \text{where } \omega_X \text{ denotes the dualizing sheaf of } X.$$

(5) We have the following exact sequence.

$$0 \longrightarrow O_P \longrightarrow \bigoplus_{i=0}^n O_P(e_i) \longrightarrow T_P \longrightarrow 0.$$

Restricting this sequence to a weighted complete intersection X in $\mathbb{P}(e)$,

we have

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \bigoplus_{i=0}^n \mathcal{O}_X(e_i) \longrightarrow T_{\mathbb{P}}|_X \longrightarrow 0.$$

(6) If X is a non-singular weighted complete intersection in $\mathbb{P}(e)$ of type $(a_1, \dots, a_c)$, then the normal bundle $N_{X/\mathbb{P}}$ of X in $\mathbb{P}(e)$ is isomorphic to $\bigoplus_{j=1}^c \mathcal{O}_X(a_j)$, and we have the following exact sequence.

$$0 \longrightarrow T_X \longrightarrow T_{\mathbb{P}}|_X \longrightarrow N_{X/\mathbb{P}} \longrightarrow 0.$$

2. Local Torelli theorem.

In this section, we shall apply the method used in the previous paper [7] for non-singular weighted complete intersections.

We use the same notations as in §1. We take $K = \mathbb{C}$ the field of complex numbers.

The following theorem is our main result.

Theorem 2.1. Let $X = \text{Proj}(\mathbb{C}[X_0, \dots, X_n]/(F_1, \dots, F_c))$ be a non-singular weighted complete intersection in $\mathbb{P}(e_0, \dots, e_n)$ of type $(a_1, \dots, a_c)$ of dimension

$d \geq 2$. Assume that $a_j > 1$ ($1 \leq j \leq c$), and that the canonical line bundle K_X is ample. Assume, furthermore,

- i) $e_0 = e_1 = 1$;
- ii) for some integer b with $2 \leq b \leq n$, $F_1, \dots, F_c, X_0, X_1, X_b$ form a regular sequence of $\mathbb{C}[X_0, \dots, X_n]$.

Then the local Torelli theorem in the d -th cohomology holds for X .

Before proving the above theorem, we first reformulate the local Torelli problem for such a non-singular weighted complete intersection X as in Theorem 2.1.

In addition to the notations in §1, we use the following notations;

$R = \mathbb{C}[X_0, \dots, X_n]$, I is the homogeneous ideal of R generated by $\{F_1, \dots, F_c\}$, and k is the integer with $O_X(k) = K_X$ (cf. (4) in §1).

From the exact sequence (5) in §1, we have the exact sequence of cohomology groups:

$$\bigoplus_{i=0}^n H^1(X, O_X(e_i)) \longrightarrow H^1(X, T_{\mathbb{P}}^1 X) \longrightarrow H^2(X, O_X) \longrightarrow \bigoplus_{i=0}^n H^2(X, O_X(e_i)).$$

By (4) in §1, we see that $\bigoplus_{i=0}^n H^1(X, O_X(e_i)) = 0$, and that $H^2(X, O_X) = 0$ in

case $d > 2$ and $H^2(X, \mathcal{O}_X) \longrightarrow \bigoplus_{i=0}^n H^2(X, \mathcal{O}_X(e_i))$ is injective in case $d = 2$.

Indeed, since $k > 0$, the last statement follows from the fact that the dual of this morphism $\bigoplus_{i=0}^n H^0(X, \mathcal{O}_X(k-e_i)) \longrightarrow H^0(X, \mathcal{O}_X(k))$ is surjective (cf. (4) in §1).

Hence we have $H^1(X, T_{\mathbb{P}}|X) = 0$. Therefore, from the exact sequences (5)

and (6) in §1, we have the following diagram.

$$\begin{array}{ccccccc}
 \bigoplus_{i=0}^n H^0(X, \mathcal{O}_X(e_i)) & & & & & & \\
 \downarrow & & & & & & \\
 H^0(X, T_{\mathbb{P}}|X) & \longrightarrow & \bigoplus_{j=1}^c H^0(X, \mathcal{O}_X(a_j)) & \longrightarrow & H^1(X, T_X) & \longrightarrow & 0, \\
 \downarrow & & & & & & \\
 0 & & & & & &
 \end{array}$$

where the row and the column are exact. Tensoring K_X to the exact sequences

(5) and (6) in §1, we obtain a similar diagram.

$$\begin{array}{ccccccc}
 \bigoplus_{i=0}^n H^0(X, \mathcal{O}_X(e_i+k)) & & & & & & \\
 \downarrow & & & & & & \\
 H^0(X, T_{\mathbb{P}}(k)|X) & \longrightarrow & \bigoplus_{j=1}^c H^0(X, \mathcal{O}_X(a_j+k)) & \longrightarrow & H^1(X, T_X(k)), & & \\
 \downarrow & & & & & & \\
 0 & & & & & &
 \end{array}$$

where the row and the column are exact. Note that the composite homomorphism

$$\bigoplus_{i=0}^n H^0(X, \mathcal{O}_X(e_i+l)) \longrightarrow H^0(X, T_{\mathbb{P}}(l)|X) \longrightarrow \bigoplus_{j=1}^c H^0(X, \mathcal{O}_X(a_j+l)) \quad (l = 0, k) \text{ is}$$

given by the Jacobian matrix

$$\begin{pmatrix} \partial_0^{F_1} & \cdots & \partial_n^{F_1} \\ \cdots & \cdots & \cdots \\ \partial_0^{F_c} & \cdots & \partial_n^{F_c} \end{pmatrix},$$

where $\partial_i^{F_j}$ denotes $\partial^{F_j}/\partial X_i$ ($1 \leq j \leq c$, $0 \leq i \leq n$). Therefore, in order to prove (P_1) for $r = d$, it is enough to prove the following (P_2) .

(P_2) Let $\tilde{\Phi}_j$ be homogeneous elements in R of degree a_j ($1 \leq j \leq c$).

Suppose that, for each homogeneous element G in R of degree k , there exist homogeneous elements $A_0, \dots, A_n$ in R satisfying

$$G \begin{pmatrix} \tilde{\Phi}_1 \\ \vdots \\ \tilde{\Phi}_c \end{pmatrix} \equiv \begin{pmatrix} \partial_0^{F_1} & \cdots & \partial_n^{F_1} \\ \cdots & \cdots & \cdots \\ \partial_0^{F_c} & \cdots & \partial_n^{F_c} \end{pmatrix} \begin{pmatrix} A_0 \\ \vdots \\ A_n \end{pmatrix} \pmod{I}.$$

Then there exist homogeneous elements $B_0, \dots, B_n$ in R satisfying

$$\begin{pmatrix} \tilde{\Phi}_1 \\ \vdots \\ \tilde{\Phi}_c \end{pmatrix} \equiv \begin{pmatrix} \partial_0^{F_1} & \cdots & \partial_n^{F_1} \\ \cdots & \cdots & \cdots \\ \partial_0^{F_c} & \cdots & \partial_n^{F_c} \end{pmatrix} \begin{pmatrix} B_0 \\ \vdots \\ B_n \end{pmatrix} \pmod{I}.$$

In the statement (P_2) above, we take G to be a monomial only in X_0, X_1 .

Then, by splitting G into a product of X_ν ($\nu = 0, 1$) and another monomial

and by multiplying the latter to $\tilde{\Phi}_j$'s, we modify (P_2) a little stronger assertion as follows.

(P_3) Let $\tilde{\Phi}_j$ be homogeneous elements in R of degree $\leq a_j + k - 1$ ($1 \leq j \leq c$). Suppose that, for each ν ($\nu = 0, 1$), there exist homogeneous elements $A_{\nu 0}, \dots, A_{\nu n}$ in R with $\deg A_{\nu i} \leq k + e_i$ ($0 \leq i \leq n$) satisfying

$$(*_{\nu}) \quad X_{\nu} \begin{bmatrix} \tilde{\Phi}_1 \\ \vdots \\ \tilde{\Phi}_c \end{bmatrix} \equiv \begin{bmatrix} \partial_0^{F_1} & \dots & \partial_n^{F_1} \\ \dots & \dots & \dots \\ \partial_0^{F_c} & \dots & \partial_n^{F_c} \end{bmatrix} \begin{bmatrix} A_{\nu 0} \\ \vdots \\ A_{\nu n} \end{bmatrix} \pmod{I}.$$

Then there exist homogeneous elements $B_0, \dots, B_n$ in R satisfying

$$\begin{bmatrix} \tilde{\Phi}_1 \\ \vdots \\ \tilde{\Phi}_c \end{bmatrix} \equiv \begin{bmatrix} \partial_0^{F_1} & \dots & \partial_n^{F_1} \\ \dots & \dots & \dots \\ \partial_0^{F_c} & \dots & \partial_n^{F_c} \end{bmatrix} \begin{bmatrix} B_0 \\ \vdots \\ B_n \end{bmatrix} \pmod{I}.$$

It is easy to see that, by the inductive use of (P_3) , we obtain (P_2) .

Next we prove some lemmas which play an essential roll in the proof of

Theorem 2.1.

Lemma 2.2. With the notations in Theorem 2.1, we have

$$H^0(X, \Omega_X^{d-1}(\ell)) = 0 \quad (\ell \leq 1).$$

Proof. We prove this in two steps.

$$\text{Step 1. } H^0(X, \Omega_{\mathbb{P}}^{d-1}(\ell)|_X) \xrightarrow{\sim} H^0(X, \Omega_X^{d-1}(\ell)) \quad (\ell \leq 1).$$

Indeed, we have the following exact sequence dual to the exact sequence (6)

in §1.

$$(1) \quad 0 \longrightarrow \check{N} \longrightarrow \Omega_{\mathbb{P}}^1|_X \longrightarrow \Omega_X^1 \longrightarrow 0,$$

where $\check{N}$ is the conormal bundle of X in $\mathbb{P}(e)$. By (6) in §1, we have

$$\check{N} = \bigoplus_{j=1}^c \mathcal{O}_X(-a_j). \quad \text{The exact sequence (1) induces a filtration of } \Omega_{\mathbb{P}}^{d-1}|_X.$$

Tensoring $\mathcal{O}_X(\ell)$, we have the filtration

$$\Omega_{\mathbb{P}}^{d-1}(\ell)|_X = E_0 \supset E_1 \supset \dots \supset E_{d-1} \supset E_d = 0$$

of $\Omega_{\mathbb{P}}^{d-1}(\ell)|_X$, with successive quotients

$$G_i = E_i/E_{i+1} \approx (\bigwedge^i \check{N}) \otimes \Omega_X^{d-1-i}(\ell).$$

Hence we have the following short exact sequences.

$$(2) \quad 0 \longrightarrow E_{i+1} \longrightarrow E_i \longrightarrow G_i \longrightarrow 0 \quad (0 \leq i \leq d-1).$$

Since $a_j > 1$ ($1 \leq j \leq c$), $(\bigwedge^i \check{N}) \otimes \mathcal{O}_X(\ell)$ ($i > 0$, $\ell \leq 1$) are direct sums of

negative line bundles. Hence, from the Nakano vanishing theorem, we have

$$H^j(X, G_i) = 0 \quad (i > 0, \quad j = 0, 1).$$

Therefore, observing the cohomology sequences of the exact sequences (2), we

obtain the desired results.

Step 2. $H^0(X, \Omega_{\mathbb{P}}^{d-1}(\ell)|_X) = 0 \quad (\ell \leq 1).$

We have the following exact sequence dual to the exact sequence (5) in §1.

$$(3) \quad 0 \longrightarrow \Omega_{\mathbb{P}}^1|_X \longrightarrow \bigoplus_{i=0}^n \mathcal{O}_X(-e_i) \longrightarrow \mathcal{O}_X \longrightarrow 0.$$

In case $d = 2$, tensoring $\mathcal{O}_X(\ell)$ to (3), we have

$$(4) \quad 0 \longrightarrow \Omega_{\mathbb{P}}^1(\ell)|_X \longrightarrow \bigoplus_{i=0}^n \mathcal{O}_X(\ell - e_i) \longrightarrow \mathcal{O}_X(\ell) \longrightarrow 0.$$

For $\ell \leq 0$, we see that $H^0(X, \mathcal{O}_X(\ell - e_i)) = 0$. Hence $H^0(X, \Omega_{\mathbb{P}}^1(\ell)|_X) = 0$. For

$\ell = 1$, we see that

$$H^0(X, \mathcal{O}_X(1 - e_i)) \cong \begin{cases} R_0 & \text{if } e_i = 1, \\ 0 & \text{if } e_i > 1, \end{cases}$$

and that

$$H^0(X, \mathcal{O}_X(1)) \cong R_1.$$

Hence we have

$$\begin{array}{ccc} \bigoplus_{i=0}^n H^0(X, \mathcal{O}_X(1 - e_i)) & \longrightarrow & H^0(X, \mathcal{O}_X(1)) \\ \uparrow \wr & & \uparrow \wr \\ R_0 \oplus \dots \oplus R_0 & \xrightarrow{\sim} & R_1. \end{array}$$

Therefore, by the exact sequence (4), we obtain the desired results in this case.

In case $d \geq 3$, observe the following exact sequence induced from (3).

$$0 \longrightarrow \Omega_{\mathbb{P}}^{d-1}(\ell)|_X \longrightarrow \left(\bigwedge^{d-1} \left(\bigoplus_{i=0}^n \mathcal{O}_X(-e_i) \right) \right) \otimes \mathcal{O}_X(\ell).$$

Since $H^0(X, (\bigwedge^{d-1} (\bigoplus_{i=0}^n \mathcal{O}_X(-e_i))) \otimes \mathcal{O}_X(l)) = 0$ ($l \leq 1$), we obtain the desired results. Q.E.D.

By virtue of Lemma 2.2, we obtain the following key lemma.

Lemma 2.3. Let A_i be homogeneous elements in R with $\deg A_i \leq k + e_i + 1$ ($0 \leq i \leq n$). Assume that

$$\begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \equiv \begin{bmatrix} \partial_0^{F_1} & \cdots & \partial_n^{F_1} \\ \cdots \cdots \cdots \\ \partial_0^{F_c} & \cdots & \partial_n^{F_c} \end{bmatrix} \begin{bmatrix} A_0 \\ \vdots \\ A_n \end{bmatrix} \pmod{I}.$$

Then there exists a homogeneous element C in R , which is independent of suffix i , such that

$$A_i \equiv e_i X_i C \pmod{I} \quad (0 \leq i \leq n).$$

Proof. From the exact sequences (5) and (6) in §1, we have the following diagram:

$$\begin{array}{c}
0 \\
\downarrow \\
H^0(X, \mathcal{O}_X(\ell')) \\
\downarrow \\
\bigoplus_{i=0}^n H^0(X, \mathcal{O}_X(e_i + \ell')) \\
\downarrow \alpha \\
\begin{array}{ccccccc}
0 & \longrightarrow & H^0(X, T_X(\ell')) & \longrightarrow & H^0(X, T_P(\ell')|_X) & \xrightarrow{\beta} & H^0(X, N_{X/P}(\ell')) \\
& & & & \downarrow & & \\
& & & & 0 & &
\end{array}
\end{array}$$

Since the composite homomorphism $\beta \cdot \alpha$ is given by the Jacobian matrix

$$\begin{pmatrix}
\partial_0^{F_1} & \dots & \partial_n^{F_1} \\
\vdots & & \vdots \\
\partial_0^{F_c} & \dots & \partial_n^{F_c}
\end{pmatrix},$$

the statement of Lemma 2.3 is equivalent to

$$(5) \quad H^0(X, T_X(\ell')) = 0 \quad (\ell' \leq k+1).$$

Moreover, since $T_X \otimes K_X \xrightarrow{\sim} \Omega_X^{d-1}$, that is, $T_X(k) = \Omega_X^{d-1}$, (5) is equivalent to Lemma 2.2. Q.E.D.

Finally we prove Theorem 2.1.

Poof. It is enough to prove the assertion (P_3) . Let $X_0(*_1)$ (resp. $X_1(*_0)$) be the equation obtained by multiplying X_0 (resp. X_1) to the equation $(*_1)$

(resp. $(*_0)$) in (P_3) . By subtracting the equation $X_1(*_0)$ from the equation $X_0(*_1)$, we have that

$$\begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \equiv \begin{bmatrix} \partial_0^{F_1} & \dots & \partial_n^{F_1} \\ \dots & \dots & \dots \\ \partial_0^{F_c} & \dots & \partial_n^{F_c} \end{bmatrix} \begin{bmatrix} X_0 A_{10} - X_1 A_{00} \\ \vdots \\ X_0 A_{1n} - X_1 A_{0n} \end{bmatrix} \pmod{I}.$$

Applying Lemma 2.3 for $A_i = X_0 A_{1i} - X_1 A_{0i}$, we obtain C in R such that

$$(6) \quad X_0 A_{1i} - X_1 A_{0i} \equiv e_i X_i C \pmod{I} \quad (0 \leq i \leq n).$$

We now use the assumption ii) repeatedly in order to cancel X_b, X_1, X_0 .

Observing the above formula (6) for $i = b$ modulo $(F_1, \dots, F_c, X_0, X_1)$, we

can cancel X_b and we have

$$C \equiv X_0 D + X_1 E \pmod{I}$$

for some D and E in R . Substituting this formula into (6), we have

$$X_0 (A_{1i} - e_i X_i D) - X_1 (A_{0i} + e_i X_i E) \equiv 0 \pmod{I}.$$

Observing this formula modulo $(F_1, \dots, F_c, X_0)$, we can cancel X_1 and we obtain

$$(7) \quad A_{0i} + e_i X_i E \equiv X_0 B_i \pmod{I}$$

for some B_i in R ($0 \leq i \leq n$). Substituting this formula (7) into $(*_0)$ in

(P_3) , we have

$$X_0 \begin{bmatrix} \tilde{\Phi}_1 \\ \vdots \\ \tilde{\Phi}_c \end{bmatrix} \equiv X_0 \begin{bmatrix} \partial_0^{F_1} & \dots & \partial_n^{F_1} & B_0 \\ \dots & \dots & \dots & \vdots \\ \partial_0^{F_c} & \dots & \partial_n^{F_c} & B_n \end{bmatrix} - E \begin{bmatrix} \partial_0^{F_1} & \dots & \partial_n^{F_1} \\ \dots & \dots & \dots \\ \partial_0^{F_c} & \dots & \partial_n^{F_c} \end{bmatrix} \begin{bmatrix} e_0 X_0 \\ \vdots \\ e_n X_n \end{bmatrix} \pmod{I}.$$

The last term in this equation vanishes by the Euler's formula. Observing this equation modulo $(F_1, \dots, F_c)$, we can cancel X_0 and we obtain the desired conclusion. Q.E.D.

§3. On the assumptions i) and ii) in Theorem 2.1.

In this section we discuss the additional assumptions i) and ii) in Theorem 2.1.

We use the same notations as in §1.

In the following cases, the assumptions i) and ii) in Theorem 2.1 are fulfilled.

Proposition 3.1. Let $X = \text{Proj}(K[X_0, \dots, X_n]/(F_1, \dots, F_c))$ be a weighted complete intersection of dimension ≥ 2 in $\mathbb{P}(e)$. If one of the following conditions is satisfied, then the assumption ii) in Theorem 2.1 is fulfilled.

i) $e_0 = e_1 = 1$, and that $\dim X$ is the maximal dimension of the complete subschemes of $\mathbb{P}(e)$.

ii) $e_0 = e_1 = \dots = e_{n-1} = 1$. In this case, we may have to change

$X_0, \dots, X_{n-1}$ by a linear transformation if necessary.

Proof. Case i). Let S_h be a maximal dimensional component of the singular locus $\bigcup_{1 \leq h} S_h$ of $Q(e)$. We see that $S_h = V_+(\{X_i \mid h \nmid e_i\})$ and that $\#\{X_i \mid h \nmid e_i\} = r(e)$ by definition. We have $\dim X = r(e) - 1$ by virtue of (1) in §1. Hence we have that

$$\#(\{F_1, \dots, F_c\} \coprod \{X_i \mid h \nmid e_i\}) = n + 1.$$

On the other hand, by the definition of weighted complete intersection, we see that $X \cap S_h = \emptyset$. Therefore, $\{F_1, \dots, F_c\} \coprod \{X_i \mid h \nmid e_i\}$ form a regular sequence of $K[X_0, \dots, X_n]$.

Case ii) In this case, the fixed points of the linear system $K[X_0, \dots, X_n]_1$ is $V_+(X_0, \dots, X_{n-1})$, which is the only singular point on $Q(e)$. Therefore, by the definition of weighted complete intersection, we get the desired assertion.

Q.E.D.

The following example also satisfies the assumptions i) and ii) in Theorem 2.1.

Example 3.2. Let $n, q, e_0, \dots, e_n, e_{n+1}, \dots, e_{n+q}$ be positive integers

with $n \geq 2$ and with $e_0 = \dots = e_n = 1$. Let $K[X_0, \dots, X_n, X_{n+1}, \dots, X_{n+q}]$ be a graded ring whose gradation is given by $\deg X_i = e_i$ ($0 \leq i \leq n+q$). Let c be a non-negative integer with $n - c \geq 2$. Take homogeneous elements $F_1, \dots, F_c, F_{c+1}, \dots, F_{c+q}$ in $K[X_0, \dots, X_{n+q}]$ of degree > 1 , which satisfy the following conditions (1) and (2).

(1) $F_j \in K[X_0, \dots, X_n]$ ($1 \leq j \leq c$), and $F_1, \dots, F_c$ form a regular sequence of $K[X_0, \dots, X_n]$.

(2) $F_{c+j} = F'_{c+j} + X_{c+j}^{b_j}$ where $F'_{c+j} \in K[X_0, \dots, X_{n+j-1}]$ and b_j is a positive integer ($1 \leq j \leq q$).

Put $Y_j = \text{Proj}(K[X_0, \dots, X_{n+j}]/(F_1, \dots, F_{c+j}))$ ($0 \leq j \leq q$) and put $Y = Y_0$ and $X = Y_q$. Then it is easy to see that the following properties (3) and (4) hold.

(3) Y is a complete intersection of dimension ≥ 2 , and Y_j is a weighted complete intersection ($1 \leq j \leq q$).

(4) The natural morphism $Y_j \longrightarrow Y_{j-1}$, corresponding to the inclusion $K[X_0, \dots, X_{n+j-1}] \hookrightarrow K[X_0, \dots, X_{n+j}]$, is a b_j -sheeted branched cyclic covering ($1 \leq j \leq q$).

Changing variables $X_0, \dots, X_n$ by a linear transformation if necessary, we

see that the above example X satisfies the assumptions i) and ii) in

Theorem 2.1.

Remark 3.3. Let Y be a complete intersection of dimension ≥ 3 , and let

$$X = Y_q \longrightarrow Y_{q-1} \longrightarrow \cdots \longrightarrow Y_1 \longrightarrow Y_0 = Y$$

be a sequence of branched cyclic coverings over Y , that is, $Y_j \longrightarrow Y_{j-1}$ is a branched cyclic covering ($1 \leq j \leq q$).

Then, by virtue of (3) in §1, we see that, as in Example 3.2, such X becomes a weighted complete intersection (cf. Corollary to Theorem 1.2 in [8]).

In case $\dim Y = 2$, the above statement is false in general (e.g. take $Y = \mathbb{P}^1 \times \mathbb{P}^1$).

Next we show some examples for which we can not apply our method.

Before showing examples, we prove a lemma.

Lemma 3.4. Let Y be a non-singular projective algebraic variety over a field K and let Z be a non-singular closed subscheme of Y defined over K . Let $i : Y \hookrightarrow \mathbb{P}^N$ be a projective embedding. Assume that $\dim Z < \frac{1}{2} \dim Y$.

Then there exists a hypersurface H in $\mathbb{P}^N$ with the following properties.

- i) $Y \cdot H$ is non-singular, and
- ii) $Y \cdot H$ contains Z .

Moreover, we may assume that, for such a hypersurface H , $\deg H$ is large enough.

Proof. Let I_Z be the $\mathcal{O}_Y$ -ideal of Z . Let $\tilde{Y} = \text{Proj}(I_Z \oplus (I_Z/I_Z^2)) \xrightarrow{\pi} Y$ be the blowing-up of Y with the center Z . Let $L = i^* \mathcal{O}_{\mathbb{P}}(1)$ be the induced line bundle on Y . Then it is easy to see that, for a large integer a , $\pi^*(L^{\otimes a}) \otimes \mathcal{O}_{\tilde{Y}}(1)$ is very ample. Let $j : \tilde{Y} \hookrightarrow \mathbb{P}^{N'}$ be the projective embedding by the global sections of $\pi^*(L^{\otimes a}) \otimes \mathcal{O}_{\tilde{Y}}(1)$. Note that each fibre of π is embedded linearly by j . We denote by B the set $\{H' \mid H' \text{ is a hyperplane of } \mathbb{P}^{N'}, \text{ which contains some fibres of } \pi^{-1}(Z) \longrightarrow Z\}$. B can be considered as a closed subscheme of $\text{Gr}(N', N'-1)$. Since $\pi^{-1}(Z) \longrightarrow Z$ is a $\mathbb{P}^r$ -bundle, where $r = \text{codim}_Y Z - 1$, Z can be considered as a closed subscheme of $\text{Gr}(N', r)$. Now consider the following diagram.

$$\begin{array}{ccc}
 & \text{Fl}(N'; N'-1, r) & \\
 p \swarrow & & \searrow q \\
 \text{Gr}(N', N'-1) \supset B & & \text{Gr}(N', r) \supset Z.
 \end{array}$$

By definition, we see that $B = p_* q^{-1}(Z)$. Since q is a $\mathbb{P}^{N'-(r+1)}$ -bundle, we see that

$$\begin{aligned} \dim p_* q^{-1}(Z) &\leq \dim q^{-1}(Z) = N' - (r+1) + \dim Z \\ &= N' - (\operatorname{codim}_Y Z - \dim Z) < N', \end{aligned}$$

where the last inequality follows from the assumption $\dim Z < \frac{1}{2} \dim Y$. Hence there is a hyperplane H' of $\mathbb{P}^{N'}$ which satisfies the following conditions.

1') $\tilde{Y} \cdot H'$ is non-singular, and

1'') H' does not contain any fibres of $\pi^{-1}(Z) \longrightarrow Z$.

Since $\mathcal{O}_Y(\pi(H')) = L^{\otimes a}$, there is a hypersurface H of $\mathbb{P}^N$ of degree a with $Y \cdot H = \pi(H')$. It is easy to see that this hypersurface H satisfies the required properties. Q.E.D.

The following example shows that if we admit only the assumption 1) in Theorem 2.1, our argument in proving the local Torelli theorem breaks down.

Example 3.5. Let e_0, e_1, e_2, e_3, e_4 be positive integers. Assume that

(1) $e_0 = e_1 = 1$, and that

(2) e_2, e_3, e_4 are greater than 1 and any two of them is mutually prime.

Let $R = \mathbb{C}[X_0, \dots, X_4]$ be the graded ring with $\deg X_i = e_i$ ($0 \leq i \leq 4$). Put

$m = \text{l.c.m.}\{e_0, \dots, e_4\} = e_2 e_3 e_4$, and let

$$F_1 = X_0^m + X_1^m + e_2 X_2^{m/e_2} + e_3 X_3^{m/e_3} + e_4 X_4^{m/e_4}$$

be a homogeneous element in R of degree m . Put $Y = \text{Proj}(R/(F_1))$ and

$Z = \text{Proj}(R/(F_1, X_0, X_1))$. Then it is easy to see by the Jacobian criterion that

Y and Z are non-singular weighted complete intersections in $\mathbb{P}(e_0, \dots, e_4)$ of type (m) and of type $(m, 1, 1)$ respectively.

Applying Lemma 3.4 for Y , Z and $L = \mathcal{O}_Y(m)$, we have a homogeneous element F_2 in R of degree am for a large enough integer a , satisfying the following conditions.

(3) $X = \text{Proj}(R/(F_1, F_2))$ is a non-singular weighted complete intersection in $\mathbb{P}(e_0, \dots, e_4)$ of type (m, am) .

(4) X contains Z .

For this X , it is easy to see the follows.

(5) $H^0(X, \Omega_X^1(l)) \neq 0$ ($l = e_2, e_3, e_4, e_3 e_4, e_4 e_2, e_2 e_3$) (cf. the proof of Lemma 2.2).

(6) F_1, F_2, X_0, X_1 do not form a regular sequence of R .

The fact (5) shows that if we want to prove the local Torelli theorem for this X by our method, we shall have to begin cancelling by using X_0 and X_1 .

But it is impossible because of the fact (6) (cf. the proof of Theorem 2.1).

The next example shows that we can not apply our method, if we remove the assumption 1) in Theorem 2.1.

Example 3.6. Let $e_0 = 5$, $e_1 = 7$, $e_2 = 8$ and $e_3 = 9$. Let X be a non-singular weighted hypersurface in $\mathbb{P}(e_0, e_1, e_2, e_3)$ (such a X exists). For this X , it is easy to see the following.

$$(1) \quad H^0(X, \Omega_X^1(\mathcal{L})) \neq 0 \quad (\mathcal{L} = e_i e_j, \quad i \neq j) \quad (\text{cf. the proof of Lemma 2.2}).$$

Hence, by using any pair of $\{X_i \mid 0 \leq i \leq 3\}$, we can not begin cancelling (cf. the proof of Theorem 2.1).

References

- [1] P. Deligne, Cohomologie des intersections complètes, exp. XI in S.G.A. 7 II, Springer Lecture Notes 340 (1973).
- [2] P. A. Griffiths, Periods of integrals on algebraic manifolds I, II, III, Amer. J. Math. 90 (1968), 568-626; 805-865; Publ. Math. I.H.E.S. 38 (1970).

- [3] P. A. Griffiths, Periods of integrals on algebraic manifolds: Summary of main results and discussion of open problems, Bull. Amer. Math. Soc. 75 (1970).
- [4] P. A. Griffiths and W. Schmid, Recent developments in Hodge theory: a discussion of techniques and results, Proc. Symp. at Bombay in 1973, Oxford Univ. Press (1975).
- [5] S. Mori, On a generalization of complete intersections, J. Math. Kyoto Univ. 15-3 (1975).
- [6] C. Peters, The local Torelli theorem of some cyclic branched coverings, to appear.
- [7] S. Usui, The local Torelli theorem for non-singular complete intersections, to appear in Japanese J. Math.
- [8] J. J. Wavrik, Deformations of branched coverings of complex manifolds, Amer. J. Math. 90 (1968).