

RELATIVE COMPACTIFICATION OF SEMIABELIAN NÉRON MODELS

I. NAKAMURA
HOKKAIDO UNIVERSITY

ABSTRACT. Let R be a complete discrete valuation ring, $k(\eta)$ its fraction field, $S = \operatorname{Spec} R$, $(G_\eta, \mathcal{L}_\eta)$ a polarized abelian variety over $k(\eta)$ with $\mathcal{L}_\eta$ symmetric ample cubical and $\mathcal{G}$ the Néron model of G_η over S . Suppose that $\mathcal{G}$ is semiabelian over S . Then there exists a *unique* relative compactification $(P, \mathcal{N})$ of $\mathcal{G}$ such that (α) P is Cohen-Macaulay with $\operatorname{codim}_P(P \setminus \mathcal{G}) = 2$ and (β) $\mathcal{N}$ is ample invertible with $\mathcal{N}|_{\mathcal{G}}$ cubical and $\mathcal{N}_\eta = \mathcal{L}_\eta^{\otimes n}$ for some positive integer n . We study the totally degenerate case in [MN24], while we study in [N24] the partially degenerate case and then the case where R is a Dedekind domain. Most remarkable is that the compactification satisfying the conditions (α) and (β) is uniquely determined by $(G_\eta, \mathbf{Z}\mathcal{L}_\eta)$.

CONTENTS

1. Introduction	2
2. Construction of a relatively complete model	3
3. Construction of P	5
4. Voronoi polytopes	6
5. The properties of P and P_0	7
6. Examples	8
References	10

The author was partially supported by the Grants-in-aid for Scientific Research (C) (No. 17K05188, 22K03261), JSPS. This work was partially supported also by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

2000 *Mathematics Subject Classification*. Primary 14K05; Secondary 14J10, 14K99.

Key words and phrases. Abelian varieties, Néron model, relative compactification.

1. INTRODUCTION

This report is based in part on a joint work with K. Mitsui. The preprints [MN24] and [N24] on the same subject appeared in arXiv.

Let R be a complete discrete valuation ring (abbr. CDVR), $k(\eta)$ (resp. $k(0)$) the fraction (resp. residue) field of R , $S := \operatorname{Spec} R$, and η (resp. 0) the generic (resp. closed) point of S .

Let $(G_\eta, \mathcal{L}_\eta)$ be a polarized abelian variety over $k(\eta)$ with $\mathcal{L}_\eta$ symmetric ample cubical, $\mathcal{G}$ the Néron model of G_η and $G := \mathcal{G}^0$ the identity component of $\mathcal{G}$. We say that G is semiabelian if G_0 is an extension of an abelian variety by a torus.

By subdividing fans associated with torus embeddings over S , [K98] proved that there are relative compactifications of a semiabelian Néron model $\mathcal{G}$ which are regular, but rarely satisfy the condition (b) of Theorem 1.1 below. Those constructed by [K98] are not codimension two compactifications in general.

In contrast with it, our compactification (which we wish to call a *cubical compactification*) of $\mathcal{G}$ is Cohen-Macaulay but not regular in general. However it is a codimension two compactification of $\mathcal{G}$ and perhaps its most remarkable feature is the uniqueness as in Theorem 1.2.

Thus we have a very simple picture:

- for a given $(G_\eta, \mathcal{L}_\eta)$, there exists a *unique Néron model* $\mathcal{G}$ of it;
- if G is semi-stable (=semi-abelian), then there exists a *unique cubical compactification* of $\mathcal{G}$.

A triple $(P, i, \mathcal{N})$ is called a *relative compactification of $\mathcal{G}$* (abbr. *compactification of $\mathcal{G}$*) (extending $(G_\eta, \mathcal{L}_\eta)$) if

- (rc1) P is an irreducible proper flat S -scheme;
- (rc2) $i : \mathcal{G} \hookrightarrow P$ is an open immersion with $P_\eta = i(\mathcal{G}_\eta) = i(G_\eta)$;
- (rc3) $\mathcal{N}$ is an ample invertible $\mathcal{O}_P$ -module with $i^*\mathcal{N}_\eta \in \mathbf{Z}_{>0}\mathcal{L}_\eta$.

We prove the following theorems in [MN24] and [N24]:

Theorem 1.1. [MN24]+[N24] *If $\mathcal{G}$ is semiabelian over S , then there exists a relative compactification $(P, i, \mathcal{N})$ of $\mathcal{G}$ such that*

- (a) P is Cohen-Macaulay;
- (b) $i(\mathcal{G}) = P \setminus \operatorname{Sing}(P_0)$ with $\operatorname{codim}_P \operatorname{Sing}(P_0) \geq 2$ where $\operatorname{Sing}(P_0)$ denotes the singular locus of P_0 ;
- (c) $i^*\mathcal{N}$ is cubical with $i_*i^*\mathcal{N} = \mathcal{N}$;
- (d) G acts on P so that $i|_G$ is G -equivariant.

Theorem 1.2. [N24] *If $(P', i', \mathcal{N}')$ is another relative compactification of $\mathcal{G}$ subject to (a)-(c), then $P \simeq P'$.*

To be more precise, we prove:

Theorem 1.3. [N24] *Let $(G, \mathcal{L})$ be a semi-abelian S -scheme, $\mathcal{G}$ its Néron model and $(P, i, \mathcal{N})$ a cubical compactification of $\mathcal{G}$ extending $(G_\eta, \mathcal{L}_\eta)$. Then*

1. $P \simeq \text{Proj } A(\mathcal{G}, i^*\mathcal{N})$ with $i_*i^*\mathcal{N} = \mathcal{N}$;
2. P is uniquely determined by $(G_\eta, \mathbf{Z}\mathcal{L}_\eta)$;
3. $\mathcal{N}$ is uniquely determined by $(G_\eta, \mathcal{L}_\eta)$ up to positive multiples;
4. any $k(\eta)$ -automorphism h_η of G_η with $h_\eta^*\mathcal{L}_\eta \simeq \mathcal{L}_\eta$ extends to an S -automorphism g of P with $g(\mathcal{G}) = \mathcal{G}$ and $g^*\mathcal{N} = \mathcal{N}$.

The following is an over-Dedekind-domain version of the above:

Theorem 1.4. [N24] *Let D be a Dedekind domain, K the fraction field of D and $S = \text{Spec } D$. Let $(G, \mathcal{L})$ be a semiabelian S -scheme over S , $\mathcal{L}$ a symmetric ample cubical invertible sheaf on G and $\mathcal{G}$ a Néron model of G . Then there exists a relative compactification $(P, i, \mathcal{N})$ of $\mathcal{G}$ extending $(G_K, \mathcal{L}_K)$ such that*

- (a) P is Cohen-Macaulay;
- (b) $i(\mathcal{G}) = P \setminus \text{Sing}(P/S)$ with $\text{codim}_P \text{Sing}(P_S) \geq 2$ where $\text{Sing}(P/S)$ denotes the union of the singular loci of closed fibers of P over S ;
- (c) $i^*\mathcal{N}$ is ample cubical invertible;
- (d) G acts on P so that $i|_G$ is G -equivariant.

Moreover the relative compactification $(P, i, \mathcal{N})$ satisfying (a)-(c) is unique up to isomorphism in the sense of Theorem 1.2.

2. CONSTRUCTION OF A RELATIVELY COMPLETE MODEL

2.1. Set-up. Let R be a CDVR, $S = \text{Spec } R$, and η (resp. 0) the generic point (resp. closed point) of S . $k(\eta)$ the fraction field of R and $k(0)$ the residue field of R by the maximal ideal I of R . Suppose that we are given a g -dimensional polarized abelian variety $(G_\eta, \mathcal{L}_\eta)$ over $k(\eta)$ with $\mathcal{L}_\eta$ ample cubical and symmetric. Let G be the connected Néron model of G_η . By choosing a suitable finite extension of $k(\eta)$, we may assume that G_0 is an extension of a g' -dimensional abelian variety A_0 over $k(0)$ by a split torus $\mathbf{G}_{m,k(0)}^{g''}$.

Throughout this report we assume $g = g''$, $A_0 = 0$ and $G_0 \simeq \mathbf{G}_{m,k(0)}^g$.

2.2. Degeneration data.

Theorem 2.1. [FC90] *Suppose that $A_0 = 0$ and G_0 is a split $k(0)$ -torus T_0 . Let X be a free $\mathbf{Z}$ -module of rank g such that $T_0 \simeq \text{Spec } k(0)[X]$. Then there exist a submodule Y of X of finite index, a function $a : Y \rightarrow k(\eta)^\times$ and a bilinear function $b : Y \times X \rightarrow k(\eta)^{\times-1}$ such that*

¹To be exact, $b(y + y', x) = b(y, x)b(y', x)$ and $b(y, x + x') = b(y, x)b(y, x')$.

- (i) $a(0) = 1, a(y) = a(-y) \ (\forall y \in Y)$;
- (ii) $b(y, z) = b(z, y) = a(y + z)a(y)^{-1}a(z)^{-1} \ (\forall y, z \in Y)$;
- (iii) $b(y, y) \in I \ (\forall y \in Y \setminus \{0\})$, and for every $n \geq 0$, $a(y) \in I^n$ for all but finitely many $y \in Y$;
- (iv) $\Gamma(G_\eta, \mathcal{L}_\eta)$ is identified with the $k(\eta)$ -vector subspace of formal Fourier series $\theta = \sum_{x \in X} \sigma_x(\theta)w^x$ such that
$$\sigma_{x+y}(\theta) = a(y)b(y, x)\sigma_x(\theta), \ \sigma_x(\theta) \in k(\eta) \ (\forall x \in X, \forall y \in Y).$$

Remark 2.2. Furthermore by taking a finite extension of $k(\eta)$ and the integral closure of R in $k(\eta)$ if necessary, we may assume:

there exist a function $a : X \rightarrow k(\eta)^\times$ and a bilinear function $b : X \times X \rightarrow k(\eta)^\times$ extending a and b in Theorem 2.1, which satisfy instead of (i)-(ii)

- (i') $a(0) = 1, a(y) = a(-y) \ (\forall y \in X)$;
- (ii') $b(y, z) = b(z, y) = a(y + z)a(y)^{-1}a(z)^{-1} \ (\forall y, z \in X)$.

2.3. Construction. We start with the stronger version of Theorem 2.1 with (i') and (ii'). Then we can construct a flat projective S -scheme P extending $(G_\eta, \mathcal{L}_\eta)$ as follows [AN99]:

- we define a graded algebra $R^\sharp := R[a(x)w^x\vartheta; x \in X]$;
- $\tilde{P}^\sharp :=$ the normalization of $\text{Proj}(R^\sharp)$ is a torus embedding over S locally of finite type;
- $\tilde{P}^{\sharp, \wedge}$: the formal completion of $R^\sharp$;
- the quotient $\tilde{P}^{\sharp, \wedge}/Y$ is algebraizable, where Y acts on $R^\sharp$ by

$$S_y^*(a(x)w^x\vartheta) = a(x + y)w^{x+y}\vartheta;$$

- the algebraization $P^\sharp$ of $\tilde{P}^{\sharp, \wedge}/Y$ is flat projective with $P^\sharp \supset G$;
- $(G_\eta, \mathcal{L}_\eta) \simeq (P_\eta^\sharp, \mathcal{O}_{P^\sharp}(1)_\eta)$.
- but $\mathcal{G} \not\subset P^\sharp$ in general.

See [AN99]. The above $(P, \mathcal{L})$ is the normalization of a projectively stable quasi-abelian scheme over S in [N99].

2.4. The complex case. Over the complex number field $\mathbf{C}$, $a(x)$ and $b(y, x)$ in the above are familiar objects:

- $D = \{s \in \mathbf{C}; |s| < 1\}$, $G = \mathbf{C}^g \times D/\mathbf{Z}^g + \mathbf{Z}^g T$;
- $a(x) = \mathbf{e}(x^t T x/2)$, $b(x, y) = \mathbf{e}(x^t T y)$;
- let $T = T_0(s) + B \log s/(2\pi i)$, with ${}^t T = T$, where $T_0(s)$ holomorphic on D , $B = \text{val}_s(b) \in M_{g \times g}(\mathbf{Z})$ even positive definite;
- if $T_0 \equiv 0$, then $a(x) = s^{B(x, x)/2}$, $b(x, y) = s^{B(x, y)}$;
- we choose $R = \mathbf{C}[[s]]$ the formal power series ring of one variable to formulate the problem in the algebraic manner.

3. CONSTRUCTION OF P

3.1. Definitions. Let $\mathcal{G}$ be the Néron model $\mathcal{G}$ of G . Next we construct a new relatively complete model $\tilde{P}$ so that P : the algebraization of $\tilde{P}^\wedge/Y$ may contain $\mathcal{G}$. Let $\Phi := \mathcal{G}_0/G_0$ be the component group of $\mathcal{G}$. Recall

$$\Phi \simeq X^\vee/\beta(Y)$$

where $\beta : Y \rightarrow X^\vee$ defined by $\beta(y)(x) = B(x, y)$. We choose $0 \in \Psi \subset X^\vee$ an arbitrary set of representatives of Φ . Let $N := |\Phi|$.

Definition 3.1. We define

$$\begin{aligned} \theta_+ &:= \prod_{u \in \Psi} \vartheta_u, \quad \theta_- := \prod_{u \in \Psi} \vartheta_{-u}, \\ \theta &:= \theta_+ \theta_-. \end{aligned}$$

Lemma 3.2. Let $\mu := \beta^{-1} \det(B) : X^\vee \rightarrow Y$, and $N = |\Phi|$. By taking a finite extension of $k(\eta)$ if necessary, there exists an extension b^e of b to $X^\vee \times X$ such that $b^e(\beta(y), x) = b(y, x)$ ($\forall y \in Y, \forall x \in X$). Set $\epsilon(u) := b^e(u, \mu(u))$ ($u \in X^\vee$). Then

- $\delta_u^* \theta = \epsilon(u) w^{2\mu(u)} \theta$;
- $\epsilon(u)$ is independent of the choice of Ψ ;
- $\epsilon(u+v) = \epsilon(u) \epsilon(v) b^e(u, \mu(v))^2$ ($\forall u, v \in X^\vee$);
- $\delta_{\beta(y)} = S_y$ ($\forall y \in Y$); this definition induces an isomorphism:

$$\{\delta_v; v \in X^\vee\} / \{S_y; y \in Y\} \simeq X^\vee/\beta(Y) \simeq \Phi.$$

Definition 3.3. Now we define an algebra

$$(1) \quad R^\dagger := R[\epsilon(u) b^e(u, \alpha) w^{\alpha+2\mu(u)} \theta; \alpha \in \Sigma, u, v \in X^\vee],$$

where Σ is the set of integral points of some bounded convex polytope (Voronoi polytope) of $X_{\mathbf{R}}$ with integral vertices. See § 4.

3.2. Construction of P . Our construction of P proceeds as follows:

- $\tilde{P} :=$ the normalization of $\text{Proj}(R^\dagger)$ is a torus embedding over S locally of finite type;
- $\tilde{P}^\wedge$: the formal completion of $\tilde{P}$;
- $X^\vee$ acts on $R^\dagger$ by

$$\delta_v^*(\epsilon(u) b^e(u, \alpha) w^{\alpha+2\mu(u)} \theta) = \epsilon(u+v) b^e(u+v, \alpha) w^{\alpha+2\mu(u+v)} \theta;$$

- $\delta_{\beta(y)} = S_y$ ($\forall y \in Y$);
- the quotient $\tilde{P}^\wedge/Y$ is algebraizable;
- the algebraization P of $\tilde{P}^\wedge/Y$ is flat projective;
- $(G_\eta, \mathcal{L}_\eta^{\otimes 2N}) \simeq (P_\eta, \mathcal{O}_P(1)_\eta)$.

To achieve $P \supset \mathcal{G}$, we need another step in § 4 of choosing a suitable multiple of $\mathcal{O}_P(1)$.

Remark 3.4. The algebra $R^\sharp$ represents the sheaf $\mathcal{L}$, while the algebra $R^\dagger$ may represent the tensor product over $P^\wedge$:

$$\bigotimes_{u \in \Psi} \delta_v^*(\mathcal{L}^\wedge)^\times.$$

4. VORONOI POLYTOPES

4.1. Voronoi polytopes. The relatively complete model $\tilde{P}$ is an S -scheme of locally of finite type, which is a torus embedding over S described by a certain fan consisting of infinitely many cones. It appears in the standard manner, which we do not write down here. See [MN24, 7.4]. Instead, we will define Voronoi polytopes, which helps us to understand the S -scheme $\tilde{P}$ to some extent.

By Th. 2.1 (ii'), we have a symmetric positive definite bilinear form

$$B(x, y) = \text{val}_s b(x, y) \quad (x, y \in X)$$

associated with a totally degenerate semiabelian S -scheme $(G, \mathcal{L})$.

Definition 4.1. Let $l \in \mathbf{Z}_{>0}$. We define a distance $\|z\| = \sqrt{B(z, z)}$ on $X_{\mathbf{R}}$. For any $c \in X^\vee$, we define a Voronoi polytope $\Sigma_l(c)$ by

$$\Sigma_l(c) = \{x \in X_{\mathbf{R}}; \|x - 2l\mu(u)\| \geq \|x - 2l\mu(c)\| \ (\forall u \in X^\vee)\}.$$

4.2. Properties. We summarize the properties of Voronoi polytopes:

- for $(G, \mathcal{L}^{\otimes l})$, we have lB and $\Sigma_l(c)$;
- $\Sigma_l(c)$ is a bounded convex polytope with $\Sigma_{l'}(0) = l'\Sigma_l(0)$;
- $\Sigma_l(c) = \Sigma_l(0) + 2l\mu(c)$;
- $X_{\mathbf{R}} = \bigcup_{c \in X^\vee} \Sigma_l(c)$;
- $\Sigma_l(c)$ ($c \in X^\vee$) and their faces form a poly. decomp. of $X_{\mathbf{R}}$;
- $\Sigma_l := \Sigma_l(0) \cap X$ is a finite subset of X with $\Sigma_l + 2l\mu(X^\vee) = X$;
- there exists $l_0 \in \mathbf{N}$ such that $\Sigma_{l_0}(0)$ is integral (Definition 4.2);
- if $B = E_8$, then $l_0 = 30$;
- if $g = 2$, then $l_0 = 1$.

Definition 4.2. A convex polytope Δ in $X_{\mathbf{R}}$ is said to be *integral* if

- (i) Δ is the convex closure of $\Delta \cap X$ and $\Delta = -\Delta \ni 0$;
- (ii) $\Delta \cap X$ contains a basis of X .

Note that $l_0 = 1$ iff $\Sigma_1(0)$ is integral.

If Δ is integral, then $\Delta \cap X$ is a star of [M72, 2.2].

4.3. Cubical compactifications.

Theorem 4.3. *Let $R^\dagger$ be the algebra (1) with $\Sigma := \Sigma_1(0) \cap X$. If $\Sigma_1(0)$ is integral, then by choosing $\Sigma := \Sigma_1(0) \cap X$, P is a cubical compactification of $(G_\eta, \mathcal{L}_\eta)$ such that $\mathcal{G} \subset P$.*

If $\Sigma_i(0)$ is integral, then we take $\mathcal{L}^{\otimes l}$ for $\mathcal{L}$ and define the algebra $R_l^\dagger$ for $R^\dagger$ with due modification. See [MN24, 9.1] for the detail.

Lemma 4.4. *Let P be a cubical compactification of $(G_\eta, \mathcal{L}_\eta)$. Then P_0 is the union of normal torus embeddings, each of which is (isomorphic to) the one associated with the same Voronoi polytope $\Sigma_i(0)$.*

See [N24, 9.10-9.12].

Remark 4.5. In general, we need pullbacks of P in § 2 and faithfully flat descents of P in § 3 in order to construct cubical compactifications in Theorems 1.1-1.4. See [MN24, §§ 9/11-12] and [N24, §§ 6/8].

5. THE PROPERTIES OF P AND P_0

5.1. The properties. Assume $\Sigma_1(0)$ is integral. The following are some standard properties of P and P_0 proved in [N24, §§ 9-10]:

- P_0 is reduced, Cohen-Macaulay with trivial dualizing sheaf;
- the semi-universal covering $\tilde{P}_0$ of P_0 is stratified into closed subschemes $\bar{Z}(\Delta)$, each $\bar{Z}(\Delta)$ being associated with a (unique) closed face Δ of some $\Sigma(c)$ ($c \in X^\vee$) and $\dim_k \bar{Z}(\Delta) = \dim_{\mathbf{R}} \Delta$;
- $H^q(P, \mathcal{O}_P)$ is an R -free module of rank $\binom{g}{q}$;
- $H^q(P, \mathcal{L}^{\otimes m}) = H^q(P_0, \mathcal{L}_0^{\otimes m}) = 0$ ($\forall q > 0, \forall m > 0$);
- $\Gamma(P, \mathcal{L}^{\otimes m})$ is a free R -module of rank $(2Nm)^g |X/Y|$;
- $\mathcal{L}^{\otimes m}$ is very ample on P for $m \geq 4g$.

5.2. The closed subscheme $\bar{Z}(\Delta)$.

Definition 5.1. Let Fan be the fan in the $(g+1)$ -dimensional Euclidean space $\mathbf{R}_{\geq 0} f_0 \oplus X_{\mathbf{R}}^\vee$ associated with the torus embedding $\tilde{P}$ over S , where f_0 is a $\mathbf{Z}$ -basis of $\mathbf{Z} \subset \mathbf{R}$. Fan consists of cones τ_Δ where Δ ranges over the set of all faces of Voronoi polytopes $\Sigma(c)$ ($c \in X^\vee$). Let $Z(\Delta) (\simeq \mathbf{G}_{m,k(0)}^{\dim \Delta})$ be the relative interior of $\bar{Z}(\Delta)$ of P_0 .

Now let us summarize the properties of $\bar{Z}(\Delta)$:

- $H^q(\bar{Z}(\Delta), \mathcal{L}_{\bar{Z}(\Delta)}^{\otimes m}) = 0$ ($\forall q \geq 1, \forall m \geq 1$);
- $H^0(\bar{Z}(\Delta), \mathcal{L}_{\bar{Z}(\Delta)}^{\otimes m})$ is spanned by monomials w^x ($x \in m\Delta \cap X$);
- $\mathcal{L}_0^{\otimes m}$ is very ample on $\bar{Z}(\Delta)$ if $m \geq \dim \Delta$.

5.3. Limits of K -rational points.

Definition 5.2. Let f_0 be a $\mathbf{Z}$ -basis of $\mathbf{Z}$, f_i ($i \in [1, g]$) a $\mathbf{Z}$ -basis of $X^\vee$, and m_i the $\mathbf{Z}$ -basis of X dual to f_i . We define

$$\text{Cut}(\tau_\Delta) := -f_0 + (\tau_\Delta \cap (f_0 \times X_{\mathbf{R}}^\vee)).$$

For example, $\Delta = \Sigma(c)$, then $\text{Cut}(\tau_\Delta) = -c$. If $\Delta = \bigcap_{c \in \Lambda} \Sigma(c)$ (Λ : maximal), then $\text{Cut}(\tau_\Delta) =$ the convex closure of $-c$ ($c \in \Lambda$).

Definition 5.3. Let s be a uniformizer of R , v_s the valuation of R with $v_s(s) = 1$ and define an *absolute value* $|\cdot|$ of $k(\eta)$ by $|a| = e^{-v_s(a)}$ for $a \in k(\eta)$. Then v_s and $|\cdot|$ can be uniquely extended to $\Omega :=$ the algebraic closure of $k(\eta)$. Note that $|a| \geq 0$, and $|a| = 0$ iff $a = 0$.

Definition 5.4. Let $Q \in \tilde{P}(\Omega)$ and $\log |w^x(Q)| := \log |Q^*(w^x)| \in \mathbf{Q}$ ($x \in \tilde{X}$). Since $w^{m_0}(Q) = s$, we define

$$\begin{aligned} \log(Q) &= -\sum_{i=0}^g (\log |w^{m_i}(Q)|) f_i \in \tilde{X}_{\mathbf{Q}}^\vee, \\ \text{cutlog}(Q) &= -f_0 + \log(Q) = -\sum_{i=1}^g (\log |w^{m_i}(Q)|) f_i \in X_{\mathbf{Q}}^\vee. \end{aligned}$$

Let K be an algebraic extension of the field $k(\eta)$. The following describes the limits of K -rational points Q [N24, § 9.6].

Theorem 5.5. *Let K be a finite extension of $k(\eta)$ and $Q \in G(K)$. Then $\lim Q \in Z(\Delta)$ iff $\text{cutlog}(Q) \in \text{Cut}(\tau_\Delta)^0$.*

Note that Δ is uniquely determined by Q in Theorem 5.5.

6. EXAMPLES

6.1. Two dimensional case. We recall known compactifications in dimension two. We consider the following case:

- X is a union of two $\mathbf{P}^1$ with 3 double points, which is a stable curve of genus two embedded in $\mathcal{X}$;
- $\mathcal{X}$ is a proper regular surface over S with $\mathcal{X}_0 = X$;
- $(G_\eta, \mathcal{L}_\eta)$ is the Jacobian variety of $\mathcal{X}_\eta$;
- $\mathcal{G}$ is the Néron model G_η , and $G := \mathcal{G}^0$;
- $G_0 \simeq \mathbf{G}_{m,k(0)}^2$ is the generalized Jacobian of X .

Let P be a relative compactification of $\mathcal{G}$. Then P_0 is

$$\begin{cases} 3\nabla & [\text{N77}] \text{ over } \mathbf{C}/\text{Non-cubical}, \\ 2\Delta & [\text{N75}] [\text{Namikawa76}][\text{AN99}], \\ 2\Delta + \nabla & [\text{M72}], \\ 2\Delta, 2\Delta + \nabla & [\text{OS79}], \\ 3\nabla & [\text{MN24}][\text{N24}]/\text{Cubical} \end{cases}$$

where $\Delta = \mathbf{P}^2$, $\nabla = Q_{p_1, p_2, p_3}(\mathbf{P}^2)$.

6.2. A smallest cubical compactification $P^\natural$. Let $X = \mathbf{Z}m_1 + \mathbf{Z}m_2 \simeq \mathbf{Z}^2$, $m_3 = -(m_1 + m_2)$, $\Sigma := \{0, \pm m_i; i \in [1, 3]\}$, $\Psi := \{0, \pm m_1\}$ and $N := |\Psi| = 3$. We define $R^\natural$ as follows:

$$R^\natural = R[\epsilon_+(u)b^e(u, \alpha)w^{\alpha+\mu(u)}\theta_+; u \in X^\vee, \alpha \in \Sigma]$$

where

$$\begin{aligned} \theta_+ &:= \prod_{y \in \Psi} \vartheta_y, \quad \epsilon_+(u) := s^{u_1^2 + u_1 u_2 + u_2^2}, \quad b^e(u, y) = s^{u_1 y_1 + u_2 y_2}, \\ \beta(y) &:= (2y_1 - y_2)f_1 + (-y_1 + 2y_2)f_2, \\ \mu(u) &:= (2u_1 + u_2)m_1 + (u_1 + 2u_2)m_2, \\ E_+(u) &:= v_s \epsilon_+(u) = u(\mu(u))/2 = u_1^2 + u_1 u_2 + u_2^2, \\ u &= u_1 f_1 + u_2 f_2, \quad y = y_1 m_1 + y_2 m_2, \\ \Sigma(0) &:= \{x \in X_{\mathbf{R}}; E_+(u) + u(x) \geq 0 \ (\forall u \in X^\vee)\}. \end{aligned}$$

It turns out

$$\Sigma(0) = \text{the convex closure of } \Sigma, \quad \Sigma = \Sigma(0) \cap X.$$




FIGURE 1. Σ and $\Sigma(0)$

The S -scheme $P^\natural$ is a cubical compactification of $\mathcal{G}$ which is constructed by starting from $\tilde{P}^\natural := \text{Proj } R^\natural$ through the process in § 3.2. By § 5, $\bar{Z}(\Sigma(c))$ is the torus embedding associated with the Voronoi polytope $\Sigma(c)$, which is ∇ . Thus we see that $P_0^\natural$ is the union of 3 copies of ∇ .

REFERENCES

- [AN99] V. Alexeev and I. Nakamura, On Mumford's construction of degenerating abelian varieties, *Tôhoku Math. J.* **51** (1999), 399-420.
- [FC90] G. Faltings, C.-L. Chai, Degenerations of abelian varieties, *Ergebnisse der Mathematik Band 22*, Springer-Verlag (1990), ix + 280 pp.
- [K98] K. Künnemann, Projective regular models for abelian varieties, semistable reduction, and the height pairing, *Duke Math. J.* **95** (1998), 161-212.
- [M72] D. Mumford, An analytic construction of degenerating abelian varieties over complete rings, *Compositio Math.* **24** (1972), 239-272.
- [MN24] K. Mitsui, I. Nakamura, Relative compactification of semiabelian Néron models, I (arXiv:2021.08113).
- [N75] I. Nakamura, *On moduli of stable quasi abelian varieties*, *Nagoya Math. J.* **58** (1975), 149-214.
- [N77] I. Nakamura, Relative compactification of Néron models and its application, *Complex analysis and algebraic geometry*, edited by Shioda and Baily, Iwanami and Cambridge Univ. press (1977), 207-225.
- [N99] I. Nakamura, Stability of degenerate abelian varieties, *Invent. Math.* **136** (1999), 659-715.
- [N24] I. Nakamura, Relative compactification of semiabelian Néron models, II (arXiv:2024.09172).
- [Namikawa76] Y. Namikawa, *A new compactification of the Siegel space and degenerations of abelian varieties*, I, II, *Math. Ann.* **221** (1976), 97-141, 201-241.
- [OS79] T. Oda, C. S. Seshadri, Compactifications of the generalized Jacobian variety, *Trans. Amer. Math. Soc.* **253** (1979), 1-90.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO, 060,
JAPAN

E-mail address: nakamura@math.sci.hokudai.ac.jp