

# On controlling the dynamical degrees of automorphisms of complex simple abelian varieties

Yutaro Sugimoto, The University of Tokyo

**Classification of abelian varieties**  
Let  $X$  be an (projective) abelian variety over  $\mathbb{C}$  and define  $\text{End}(X) := \{f : X \rightarrow X : \text{morphism} \mid f \text{ preserves the group structure}\}$ . If  $X$  is a simple abelian variety, then the structure of  $\text{End}_{\mathbb{Q}}(X) := \text{End}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$  is classified as below. (cf. [1, Chapter 5.5])

| Classification of $\text{End}_{\mathbb{Q}}(X)$ |                                                |              |                   |                       |
|------------------------------------------------|------------------------------------------------|--------------|-------------------|-----------------------|
|                                                | $B = \text{End}_{\mathbb{Q}}(X)$               | $K$          | $d   e_0$         | restriction           |
| Type I                                         | $K$                                            | totally real | $1   e$           | $e   g$               |
| Type II                                        | totally indefinite quaternion algebra over $K$ | totally real | $2   e$           | $2e   g$              |
| Type III                                       | totally definite quaternion algebra over $K$   | totally real | $2   e$           | $2e   g$              |
| Type IV                                        | division ring with center $K$                  | CM-field     | $d   \frac{e}{2}$ | $\frac{d^2 e}{2}   g$ |

**Remark**  
 $B := \text{End}_{\mathbb{Q}}(X)$  is a division ring,  $\phi : B \rightarrow B$  is a Rosati involution,  $K$  is a center of  $B$ ,  $K_0 := \{x \in K \mid \phi(x) = x\}$ ,  $[B : K] = d^2$ ,  $[K : \mathbb{Q}] = e$  and  $[K_0 : \mathbb{Q}] = e_0$ .  
For each type of division ring  $B$ , for almost all cases, there is a construction of a complex simple abelian variety  $X$  such that  $\text{End}_{\mathbb{Q}}(X) = B$  (see below).

**Dynamical degrees of endomorphisms of abelian varieties**  
Fix  $f \in \text{End}(X)$  for a  $g$ -dimensional complex abelian variety  $X$ . The dynamical degrees of  $f$  are defined as  $\lambda_k(f) := \lim_{n \rightarrow +\infty} \|(f^n)^* : H^{k,k}(X) \rightarrow H^{k,k}(X)\|^{\frac{1}{n}}$  for  $0 \leq k \leq g$ , where  $\|\cdot\|$  is a norm for linear transformations and  $H^{k,k}(X)$  is the  $(k, k)$ -Dolbeault cohomology.  
Writing  $X = \mathbb{C}^g / \Lambda$  and  $f$  induces  $\rho_f : \mathbb{C}^g \rightarrow \mathbb{C}^g$ . Under this notation, the dynamical degrees of  $f$  can be calculated as  $\lambda_k(f) := \prod_{i=1}^k |\rho_i|$  where  $\{\rho_i\}_{1 \leq i \leq g}$  are eigenvalues of  $\rho_f$  which are renumbering as  $|\rho_1| \geq |\rho_2| \geq \dots \geq |\rho_g| > 0$ . By using the Lefschetz fixed point theorem, this value can be calculated more concretely.

**Theorem (cf. [1, Chapter 13])**  
Assume  $X$  is simple and denote  $B := \text{End}_{\mathbb{Q}}(X)$  and define  $d, e, e_0$  as above.  
Then,  $\prod_{i=1}^g (1 - \rho_i)(1 - \bar{\rho}_i) = \text{Nrd}_{B/\mathbb{Q}}(\text{id}_X - f)^{\frac{2g}{e}}$ , where  $\text{Nrd}_{B/\mathbb{Q}}$  is the reduced norm.  
By applying this theorem for  $n_X - f \in \text{End}(X)$  ( $n \in \mathbb{Z}$ ), we get the next theorem.

**Theorem (cf. [5, Section 3], [3, Section 2.2])**  
Assume  $X$  is simple and  $f \in \text{End}(X)$ . Let  $P(x) \in \mathbb{Z}[x]$  be the minimal polynomial of  $f$  in  $\text{End}_{\mathbb{Q}}(X)$ . Then, each root of  $P(x)$  appears the same number of times in the set  $\{\rho_1, \dots, \rho_g, \bar{\rho}_1, \dots, \bar{\rho}_g\}$ .  
Thus, the calculation of the dynamical degree of endomorphisms on simple abelian varieties is returned to considering  $f \in B$  as the element of the division ring.

**Constructing simple abelian varieties from division ring**  
Let  $(B, \mathcal{O}, \phi)$  be a triple of a division ring  $B$  of any type in the above table, a  $\mathbb{Z}$ -order  $\mathcal{O}$  for  $B$ , and a positive anti-involution  $\phi : B \rightarrow B$ , which fixes  $K_0$  in  $K$ . The construction of a complex abelian variety  $X$  with  $B \hookrightarrow \text{End}_{\mathbb{Q}}(X)$  such that  $\mathcal{O} \hookrightarrow \text{End}(X)$  and  $\phi$  is compatible with the Rosati involution of  $X$ , is explained in [1, Chapter 9]. If certain conditions for  $d, e$ , and  $e_0$  hold, then  $B \hookrightarrow \text{End}_{\mathbb{Q}}(X)$  can be an isomorphism, and now  $X$  is a simple abelian variety.

**Realization of Salem number**  
A Salem number is a real algebraic integer  $\lambda$  greater than 1 where its other conjugates have modulus at most equal to 1, at least one having a modulus equal to 1.  
There is a relation between the first dynamical degree and the Salem number.

**Theorem ([2, Theorem 5.1(1)])**  
Let  $X$  be a compact Kähler surface and  $f : X \dashrightarrow X$  be a bimeromorphic map with  $\rho(f^*) > 1$  for the operator  $f^*$  on  $H^{1,1}(X)$  where  $\rho$  is the spectral radius.  
Then the operator  $f^*$  has exactly one eigenvalue  $\lambda \in \mathbb{R}_{>0}$ , of modulus  $|\lambda| > 1$ , and in fact  $\lambda = \rho(f^*)$ .

The realization of a Salem number as the first dynamical degree is one of the topics, but for the case of 2-dimensional abelian varieties, realizable Salem numbers have a degree at most 4. For higher-dimensional cases, a realization of a part of Salem numbers is proved in [3, Theorem 2.1 (i)].  
We get all Salem numbers are realizable as the first dynamical degree of an automorphism of a simple abelian variety. It proceeds as follows (cf. [5, Section 5]).  
Let  $\lambda$  be a Salem number of degree  $g$ , and let  $f(x) \in \mathbb{Z}[x]$  be the minimal polynomial of  $\sqrt{\lambda}$ . Let  $\gamma$  be one of the roots of  $f(X)$  which has modulus 1, and define  $L := \mathbb{Q}(\gamma)$ ,  $K := \mathbb{Q}(\gamma + \frac{1}{\gamma})$ . Then, there exists an algebraic integer  $a \in \mathcal{O}_K$  such that  $K(\sqrt{a}) = L$ , and a prime number  $p$  such that the quaternion algebra  $(\frac{a,p}{K})$  is divisional (cf. [3, Theorem 2.5, Lemma 2.11]). Existence of a positive anti-involution  $\phi$  over  $B := (\frac{a,p}{K})$  is reduced to the existence of  $c \in B \setminus K$  such that  $c^2$  is totally negative (cf. [1, Theorem 5.5.3]), and this can be achieved.  
Let  $1, i, j, ij$  be the basis of  $(\frac{a,p}{K})$  such that  $i^2 = a, j^2 = p, ij = -ji$ , and then

$\mathcal{O} := \mathcal{O}_L \oplus \mathcal{O}_L j$  is a  $\mathbb{Z}$ -order of  $(\frac{a,p}{K})$ . Thus, we get a suitable complex simple abelian variety  $X$  (of Type II) from the triple  $(B, \mathcal{O}, \phi)$  as above, and  $\text{End}(X)$  contains  $\gamma$ . The minimal polynomial of  $\gamma$  is  $f(x)$ , and the maximal absolute value of its roots is  $\sqrt{\lambda}$ . Thus, the first dynamical degree of the automorphism, which corresponds to  $\gamma \in \text{End}(X)$ , is  $\sqrt{\lambda^2} = \lambda$ .

**Small first dynamical degrees with fixing dimension**  
For a  $g$ -dimensional simple abelian variety  $X$  and its automorphism, we consider the minimal polynomial of automorphism in endomorphism algebra  $\text{End}_{\mathbb{Q}}(X)$ . Now the first dynamical degree of the automorphism is the square of the maximal absolute value of the roots of the minimal polynomial. Since the degree of the minimal polynomial is at most  $2g$  by the above table, there are only finitely many candidates of the minimal polynomial when considering the first dynamical degree is less than  $1 + \epsilon$  for any  $\epsilon$ . Thus, the smallest first dynamical degree of automorphisms, which are larger than 1, exists for each  $g$ , and it can be calculated by dividing by cases from Type I to Type IV. We denote the minimum value by  $m(g)$ , and the result for small  $g$  and the prime cases are as below.

| Theorem ([6, Theorem B]) |                                  |
|--------------------------|----------------------------------|
| dimension $g$            | minimum value $m(g)$             |
| 1                        | (all 1)                          |
| 2                        | $4\cos^2(\pi/5) = 2.6180 \dots$  |
| 3                        | $4\cos^2(\pi/7) = 3.2469 \dots$  |
| 4                        | $2\cos(\pi/5) = 1.6180 \dots$    |
| 5                        | $4\cos^2(\pi/11) = 3.6825 \dots$ |
| 6                        | $2\cos(\pi/7) = 1.8019 \dots$    |
| 7                        | $4.0333 \dots$                   |
| 8                        | $2\cos(\pi/5) = 1.6180 \dots$    |
| 9                        | $1.1509 \dots^2 = 1.3247 \dots$  |
| 10                       | $1.1762 \dots^2 = 1.3836 \dots$  |

**Theorem ([6, Theorem A])**  
For a prime number  $p$ ,  
$$\begin{cases} m(p) = 4\cos^2(\frac{\pi}{2p+1}) & (\text{if } 2p+1 \text{ is also a prime}) \\ 4 + 4^{-2p-3} < m(p) < 4.3264 = (2.08)^2 & (\text{otherwise}) \end{cases}$$

**Small first dynamical degrees without fixing dimension**  
Without fixing the dimension of simple abelian varieties, there exists an automorphism, whose dynamical degree is as close to 1 as possible, but not 1. It proceeds as follows (cf. [5, Section 8]). Let  $p > 3$  be a prime number and define  $F := \mathbb{Q}(\zeta_p, \sqrt[p]{\zeta_p + \frac{1}{\zeta_p}})$ ,  $K = \mathbb{Q}(\zeta_p)$  and  $K_0 = \mathbb{Q}(\zeta_p + \frac{1}{\zeta_p})$ . Now  $F/K$  is a cyclic extension of degree  $p$ ,  $K_0$  is a totally real number field, and its field extension  $K$  is a CM-field. There exists an element  $u \in K^\times / N_{F/K}(F^\times)$  whose order is exactly  $p$ , and define  $B = \bigoplus_{i=0}^{p-1} u^i F$ . Let  $\sigma$  be the generator of  $\text{Gal}(F/K)$  and define the multiplication on  $B$  by  $e \cdot u = u \cdot \sigma(e)$  for  $e \in F$ . By [4, Chapter 15.1, Corollary d],  $B$  is a division ring, which is of Type IV in the table.  
Define  $a := \zeta_p + \frac{1}{\zeta_p}$  and then the map

$$\phi : B \rightarrow B$$
$$u^i \zeta_p^j \sqrt[p]{a^k} \mapsto u^i \zeta_p^{ik-j} \sqrt[p]{a^k} \quad (0 \leq i, k \leq p-1, 0 \leq j \leq p-2)$$

is an anti-involution over  $\mathbb{Q}$  with fixing  $K_0$  in  $K$ . By [1, Theorem 5.5.6], there exists a positive anti-involution on  $B$ .  
Now by defining  $\mathcal{O} = \bigoplus_{i=0}^{p-1} u^i \mathcal{O}_F$ , the triple  $(B, \mathcal{O}, \phi)$  induces a complex simple abelian variety  $X$  (of Type IV, dimension  $\frac{p^2(p-1)}{2}$ ) with  $\sqrt[p]{\zeta_p + \frac{1}{\zeta_p}} \in \text{End}(X)$ . The minimal polynomial of  $\sqrt[p]{\zeta_p + \frac{1}{\zeta_p}}$  over  $\mathbb{Q}$  is  $\prod_{k=1}^{\frac{p-1}{2}} (x^p - (\zeta_p^k + \frac{1}{\zeta_p^k}))$  and so the first dynamical degree of the corresponding automorphism is  $\sqrt[p]{\left|\zeta_p^{\frac{p-1}{2}} + \frac{1}{\zeta_p^{\frac{p-1}{2}}}\right|^2}$ . By  $p \rightarrow \infty$ , this value converges to 1.

**References**

- [1] Birkenhake, C., Lange, H. : *Complex abelian varieties*, second ed., Grundlehren der mathematischen Wissenschaften, 302, Springer-Verlag, Berlin, 2004.
- [2] Diller, J., Favre, C. : *Dynamics of bimeromorphic maps of surfaces*, American Journal of Mathematics, 123(2001), No. 6, 1135-1169.
- [3] Dang, N.-B., Herrig, T. : *Dynamical degrees of automorphisms on abelian varieties*, Advances in Mathematics, 395(2022), 108082.
- [4] Pierce, R.S. : *Associative Algebras*, Graduate Texts in Mathematics, 88, Springer, New York, 1982.
- [5] Sugimoto, Y. : *Dynamical degrees of automorphisms of complex simple abelian varieties and Salem numbers*, 2023., arXiv:2302.02271.
- [6] Sugimoto, Y. : *The minimum value of the first dynamical degrees of automorphisms of complex simple abelian varieties with fixing dimensions*, 2023., arXiv:2306.09271.